Global Journal of Mathematical Analysis, 7 (2) (2019) 15-18

Global Journal of Mathematical Analysis

Website: www.sciencepubco.com/index.php/GJMA

Research paper

New algorithm method for solving the variational inequality problem in Hilbert space

Zena Hussein Maibed*

Department of Mathematics, College of Education for Pure Science, Ibn Al-Haithem , University of Baghdad
**Corresponding author E-mail: a.hussein@yahoo.com*

Abstract

The purpose of this paper is to introduce a concept of generalized non_spreading and define a new algorithm for infinite families of generalized non_spreading and finite families of resolvent mappings. Also, We study the existence solution of variational inequality to a common fixed point in Hilbert spaces. The main results in this paper extend and generalized of many known results in the literature.

Keywords: *Resolven; Mapping; Non-Spreading; Mapping; Common; Fixed Point; Strong Convergence.*

1. Introduction

Let K be a nonempty bounded closed convex subset of a Hibert space X. A mapping $\mathcal{P}$: K $\rightarrow$ K is said to be nonexpansive if $\|\mathcal{P}\,x - \mathcal{P}\,y\| \leq \|x - y\|$, for all $x, y \in K$. On the other hand anyimultivalued mapping $\mathcal{G}$ is called monotone if: $< z_1 - z_2 , w_1 - w_1 > \geq 0\ \forall z_i \in D(\mathcal{G})$, for all $w_i \in \mathcal{G}(z_i)$. And it is called maximal monotone, (M,.M), if for all $(z, h) \in X \times X, \langle z - w, h - k\rangle \geq 0$ andifor all $(w, k) \in gph(\mathcal{G})$ then we get, $h \in \mathcal{G}(z)$. The Monotone mappings play important role in optimization Theory see the books([1]-[5]). It has been shown that if X is uniformly convex then every nonexpansive mapping $\mathcal{P}$: K $\rightarrow$ K has a fixed point (see Browder [6], cf. also Kirk [7]). In 1974, Ishikawa [8] introduced a new iteration procedure for approximating fixed points of pseudo-contractive compact mappings in Hilbert spaces Note that the normal Mann iteration procedure [9], is a special case of the Ishikawa one. For a comparison of the two iterative processes in the one-dimensional case we referithe reader to Rhoades [10]. For more details and literature on the convergence of Ishikawa and Mann iterates we refer to [11–18]. Recently, Sastry and Babu [19] introduced the analogs of Mann and Ishikawa iteratesifor multivalued mappings and proved convergence theorems for nonexpansive mappings whose domain is a compact convex subset of a Hilbert space. The convergence of the iteration processes are studied by many researchers, see([20]-[29]). In this paper, we generalize results of Sastry and Babu to uniformly convex Banach spaces. We also introduce both of the iteration processes in a new sense and prove a convergence theorem of Mann iterates for a mapping defined on a non compact domain. Now, we recall some definitions and lemmas which will used in the proofs. Lemma (1.1): [7]
If $\langle a_n\rangle$ be a sequenceiof non-negativeireal numberisuch that:
$a_{n+1} \leq (1 - \gamma_n)a_n + S_n , n \geq 0$, where $\langle\gamma_n\rangle$ is a sequence in (0, 1) and $\langle S_n\rangle$ is a sequence in $\mathbb{R}$, $\sum_{n=0}^{\infty}\gamma_n = \infty$ and $\lim\sup_{n\to\infty}\frac{S_n}{\gamma_n} \leq 0$ or $\sum_{n=1}^{\infty}|S_n| < \infty$. Then $a_n \rightarrow 0$ as $n \rightarrow \infty$.

Lemma 1.2: *[15] If $\langle a_n\rangle$ be a sequence nonnegative real numbers such that:*

$a_{n+1} \leq (1 - \gamma_n)a_n + \gamma_n S_n + \beta_n , n \geq 0$, where $\langle\gamma_n\rangle, \langle\beta_n\rangle$ and $\langle S_n\rangle$ are satisfies the following:

1) $\gamma_n \subset [0,1]$; $\sum_{n=1}^{\infty}\gamma_n = \infty$
2) $\lim\sup_{n\to\infty} S_n \leq 0$ or $\sum_{n=1}^{\infty}|\gamma_n S_n| < \infty$
3) $\beta_n \geq 0$ for each $n \geq 0$ such that $\sum_{n=0}^{\infty}\beta_n < \infty$. Then $\lim_{n\to\infty} a_n = 0$.

Lemma 1.3: *[16] Let C be a nonempty convex closed subset of real Hilbert space $X$and $\mathcal{P}$ is non-expansive multivalued operator with $Fix(\mathcal{P}) \neq \emptyset$.If $\langle x_n\rangle$ sequence in C such that $x_n \rightharpoonup x$ and $(I - \mathcal{P})x_n \rightarrow y, x, y \in C$ .Then we have $(I - \mathcal{P})x = y$.*

2. Main results

In this section, we define a new iterations for sequence of generalized non_spreading mapping and then study the convergence for these schemes.

Copyright © Zena Hussein. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Definition 2.1: *A mapping $\mathcal{P}$ is generalizedinon_spreading mapping if for each x, y in $D(\mathcal{P})$, there exists positive real sequence $\langle z_n \rangle$ then the following inequality holds*

$$||\mathcal{P}x - \mathcal{P}y||^2 \leq ||x - y||^2 + z_n < x - \mathcal{P}x, y - \mathcal{P}y >$$

Lemma 2.1: *Let $\mathcal{G}_1, \mathcal{G}_2, ,,, \mathcal{G}_m$; are M.M mappings, $\langle f_n \rangle$ be a sequence of generalized non_spreading mapping on C. If $\sum \alpha_{n,i} = 1$ and $\mathcal{P}^i_{\sigma_n} = \sigma_n f_n(x) + (1 - \sigma_n) \sum_{i=1}^m \alpha_{n,i} J^i_{r_n,i}(x)$, where $\langle \alpha_{n,i} \rangle$ be a sequences in (0, 1] , $\langle \sigma_n \rangle$ $\langle \sigma_n \rangle$ be a sequence of positive real numbers and $\langle J^i_{r_n,i} \rangle$ be a sequence of resolvent mapping. Then $\mathcal{P}^i_{\sigma_n}$ are also generalized non_spreading on C for all $i = 1,2, \ldots\ldots\ldots, m$.*

Proof
For all $x, y \in C$

$$\left\|\mathcal{P}^i_{\sigma_n}(x) - \mathcal{P}^i_{\sigma_n}(y)\right\|^2 = \left\| \sigma_n f_n(x) + (1 - \sigma_n) \sum_{i=1}^m \alpha_{n,i} J^i_{r_n,i}(x) - \sigma_n f_n(y) - (1 - \sigma_n) \sum_{i=1}^m \alpha_{n,i} J^i_{r_n,i}(y) \right\|^2$$

$$\left\|\mathcal{P}^i_{\sigma_n}(x) - \mathcal{P}^i_{\sigma_n}(y)\right\|^2 \leq \left\|\sigma_n \left(f_n(x) - f_n(y)\right) + (1 - \sigma_n) \sum \alpha_{n,i} \left(J^i_{r_n,i}(x) - J^i_{r_n,i}(y)\right\|^2 \right. \leq \sigma_n \|f_n(x) - f_n(y)\|^2 + (1 - \sigma_n) \sum_{i=1}^m \alpha_{n,i} \left\|J^i_{r_n,i}(x) - J^i_{r_n,i}(y)\right\|^2$$

$$= \sigma_n \|x - y\|^2 + \sigma_n z_n < x - \mathcal{P}x, y - \mathcal{P}y > + (1 - \sigma_n)\|x - y\|^2 = \|x - y\|^2 + \sigma_n z_n < x - \mathcal{P}x, y - \mathcal{P}y >$$

Theorem 2.2: *let $\mathcal{G}_1, \mathcal{G}_2, ,,, \mathcal{G}_m$ are M.M mappings, $\langle f_n \rangle$ be a sequence of generalized non_spreading mapping on C and $(\cap_{i=1}^{\infty} F(f_n)) \cap \left(\cap_{i=1}^m \mathcal{G}_i^{-1}(0)\right) \neq \emptyset$ for all $i = 1,2, \ldots\ldots\ldots, m$. If the iteration process $\langle x_n \rangle$ defined as the following $x_{n+1} = \sigma_n f_n(x_n) + (1 - \sigma_n) \sum_{i=1}^m \alpha_{n,i} J^i_{r_n,i}\left(x_{\sigma_n}\right)$, where $\langle \sigma_n \rangle$ be a sequence of positive real numbers convergent to 0.Then there exist a solution of variational inequality*

$$\langle (I - f_n)\left(x_{n_{k+1}}\right), x_{n_k} - \tilde{x} \rangle \leq 0 \leq 0$$

Proof

Let $v \in (\cap_{i=1}^{\infty} F(f_n)) \cap \left(\cap_{i=1}^m \mathcal{G}_i^{-1}(0)\right)$

$$\|x_{n+1} - v\|^2 \leq \sigma_n \|f_n(x_t) - v\|^2 + (1 - \sigma_n) \sum_{i=1}^m \alpha_{n,i} \left\|J^i_{r_n,i}(x_k) - v\right\|^2$$

$$\leq \sigma_n \|x_k - v\|^2 + \sigma_n z_n \langle x_k - f_n x, v - f_n v \rangle + (1 - \sigma_n) \sum_{i=1}^m \alpha_{n,i} \|x_k - v\|^2$$

$$\|x_{k+1} - v\|^2 \leq \sigma_n \|x_k - v\|^2 + (1 - \sigma_n)\|x_k - v\|^2 + \sigma_n z_n \langle x - f_n x, v - f_n v \rangle$$

$$\|x_{k+1} - v\|^2 \leq \|x_k - v\|^2 + \sigma_n z_n \langle x - f_n x, v - f_n v \rangle$$

Hence, $\|x_{k+1} - v\|^2 \leq \|x_k - v\|^2$

So, $\langle x_k \rangle$ is bounded sequence.

$$\left\|x_{k+1} - \sum_{i=1}^m \alpha_{n,i} J^i_{r_n,i} x_k\right\| = \left\|\sigma_n f_n(x_k) + (1 - \sigma_n) \sum_{i=1}^m \alpha_{n,i} J^i_{r_n,i}(x_k) - \sum_{i=1}^m \alpha_{n,i} J^i_{r_n,i}(x_k)\right\|$$

$$\leq \sigma_n \left\|f_n(x_k) - \sum_{i=1}^m \alpha_{n,i} J^i_{r_n,i}(x_k)\right\|$$

Since $\langle f_n(x_k) \rangle$ and $\langle J^i_{r_n,i}(x_k) \rangle$ are bounded and $\langle \sigma_n \rangle$ be a sequence of positive real numbers convergent to 0, then as $n \longrightarrow 0$ we get

$\left\|x_k - \sum_{i=1}^m \alpha_{n,i} J^i_{r_n,i}(x_k)\right\| \longrightarrow 0$, then we get

$\left\|x_{k+1} - J^i_{r_n,i} x_k\right\| \longrightarrow 0$, for all $i = 1,2, \ldots\ldots\ldots, m$

Now, since $\langle x_k \rangle$ is bounded sequence then there exists $\langle x_{kn} \rangle$ subsequence of $\langle x_k \rangle$ such that

$x_{kn} \rightharpoonup \check{x}$. By lemma (1.3) $\Longrightarrow \check{x} \in \mathcal{G}_i^{-1}(0)$ $\forall$ $i = 1,2, \ldots\ldots\ldots, m$

$x_{k+1} - \check{x} = k(f_n(x_k) - \check{x}) + (1 - k) \sum_{i=1}^m \alpha_{n,i} (J^i_{r_n,i} x_k - \check{x})$

Using $x_k - \check{x}$ to make inner product, we get

$$\|x_k - \check{x}\|^2 = \sigma_n \langle f_n(x_k) - \check{x}, x_k - \check{x} \rangle + (1 - \sigma_n) \sum_{i=1}^m \alpha_{n,i} \langle J^i_{r_n,i} x_k - \check{x}, x_k - \check{x} \rangle$$

$$\leq \sigma_n \langle f_n(x_k) - \check{x}, x_k - \check{x} \rangle + (1 - \sigma_n) \sum_{i=1}^m \alpha_{n,i} \langle x_k - \check{x}, x_k - \check{x} \rangle$$

$$= \sigma_n \langle f_n(x_k) - \check{x}, x_k - \check{x} \rangle + (1 - \sigma_n)\|x_k - \check{x}\|^2.$$

$\sigma_n \ \|x_{k+1} - \breve{x}\|^2 \leq \sigma_n \langle f_n(x_k) - \breve{x}, x_k - \breve{x} \rangle$

$\|x_{k+1} - \breve{x}\|^2 \leq \langle f_n(x_k) - \breve{x}, x_k - \breve{x} \rangle$

$\leq \langle f_n(x_k) - f_n(\breve{x}), x_k - \breve{x} \rangle + \langle f_n(\breve{x}) - \breve{x}, x_k - \breve{x} \rangle$

$\|x_{k+1} - \breve{x}\| \leq \|x_k - \breve{x}\|^2 + z_n \|x_k - \breve{x}\| . \langle x_k - f_n x_k, \breve{x} - f_n \breve{x} \rangle + \langle f_n(\breve{x}) - \breve{x}, x_k - \breve{x} \rangle$

In particular,

$\left\|x_{(k+1)n} - \breve{x}\right\|^2 \leq \|x_{kn} - \breve{x}\|^2 + z_n \|x_{kn} - \breve{x}\| . \langle x_{kn} - f_n x_{kn}, \breve{x} - f_n \breve{x} \rangle + \langle f_n(\breve{x}) - \breve{x}, x_{kn} - \breve{x} \rangle$

But $x_{kn} \rightharpoonup \breve{x}$, then$||x_{kn} - \breve{x}||^2 \longrightarrow 0$ as $n \longrightarrow \infty$

Now, since

$x_{k+1} - f_n(x_{k+1}) = \sigma_n f_n(x_k) - f_n(x_k) + f_n(x_k) + f_n(x_{k+1}) + (1 - \sigma_n) \sum_{i=1}^{m} \alpha_{n,i} J_{r_n,i}^{i} x_k$

Hence,

$x_{n+1} - f_n(x_n) = (f_n(x_k) + f_n(x_{k+1})) + (\sigma_n - 1) f_n(x_n) + (1 - \sigma_n) \sum_{i=1}^{m} \alpha_{n,i} J_{r_n,i}^{i}(x_n)$

$x_{n+1} - f_n(x_n) = -(1 - \sigma_n)(f_n - \sum_{i=1}^{m} \alpha_{n,i} J_{r_n,i}^{i}) x_k + f_n(x_k) + f_n(x_{k+1})$

$For\ all\ z \in (\cap_{i=1}^{\infty} F(f_n)) \cap \left(\cap_{i=1}^{m} \mathcal{G}_i^{-1}(0) \right)$

$\langle x_{n+1} - f_n(x_n), x_k - z \rangle = -(1 - \sigma_n) \langle f_n - \sum_{i=1}^{m} \alpha_{n,i} J_{r_n,i}^{i}) x_k + (f_n(x_k) + f_n(x_{k+1})), x_k - z \rangle \leq 0$

(As $(f_n(x_k) + f_n(x_{k+1}) + (f_n - \sum_{i=1}^{m} \alpha_{n,i} J_{r_n,i}^{i})(x_k))$ is monotone mapping. It is true for any subsequence of $\langle x_k \rangle$.
And hence, as $n_k \to \infty$ we get, $\tilde{x}$ is a solution of variational inequality

$\langle (I - f_n)(x_{nk+1}), x_{nk} - \tilde{x} \rangle \leq 0$, as n $\to \infty$.

References

[1] K.P.R. Sastry, G.V.R. Babu, Convergence of Ishikawa iterates for a multi-valued mapping with a fixed point, Czechoslovak Math. J. 55 (2005) 817–826. https://doi.org/10.1007/s10587-005-0068-z.

[2] F.E. Browder, Nonlinear operators and nonlinear equations of evolution in Banach spaces, in: Proc. Symp. Pure Math., vol. 18, Amer. Math. Soc., Providence, RI, 1976.

[3] W.A. Kirk, A fixed point theorem for mappings which do not increase distance, Amer. Math. Monthly 72 (1965) 1004–1006. https://doi.org/10.2307/2313345.

[4] S. Ishikawa, Fixed points by a new iteration method, Proc. Amer. Math. Soc. 44 (1974) 147–150. https://doi.org/10.1090/S0002-9939-1974-0336469-5.

[5] W.R. Mann Mean value methods in iteration, Proc. Amer. Math. Soc. 4 (1953) 506–510. https://doi.org/10.1090/S0002-9939-1953-0054846-3.

[6] J.M. Borwein and J.D. Vanderwerff, Convex Functions, Cambridge University Press, 2010.

[7] R.S. Burachik and A.N. Iusem, Set-Valued Mappings and Enlargements of Monotone Operators, Springer-Verlag, 2008. 24.

[8] S. Simons, From Hahn-Banach to Monotonicity, Springer-Verlag, 2008.

[9] C. Z ˇalinescu , Convex Analysis in General Vector Spaces, world Scientific Publishing, 2002. https://doi.org/10.1142/9789812777096.

[10] H.K.Xu," a nother control condition in an iterative method for nonexpansive mappings, bull, austral. Math. soc, 65 (2002), 109-113. https://doi.org/10.1017/S0004972700020116.

[11] H.K.Xu," Iterative Algorithm for nonlinear operators",J. london Math.soc .66(2002) 240-256 https://doi.org/10.1112/S0024610702003332.

[12] A.Moudafi, "Viscosity approximation method for fixed point problems", Journal of Mathematical Analysis and Applications, 241(2000) 46-55.

[13] H.K.Xu, "Viscosity approximation methods for non-expansive mapping ", J. Math .Anal.Appl. 298 (2004) 279-291. https://doi.org/10.1016/j.jmaa.2004.04.059.

[14] S. Kamimura, W. Takahashi, "Approximating solutions of maximal monotone operators in Hilbert spaces", J. Approx. Theory 106 (2000) 226–240. https://doi.org/10.1006/jath.2000.3493.

[15] Z. H. Mabeed, "Strongly Convergence Theorems of Ishikawa Iteration Process with Errors in Banach Space" Journal of Qadisiyah Computer Science and Mathematics, 3(2011)1-8.

[16] Z. H. Mabeed, Some Convergence Theorems for The Fixed Point in Banach Spaces, Journal of university of Anbar for pure science, 2(2013)7.2.

[17] Z. H. Maibed, Strong Convergence of Iteration Processes for Infinite Family of General Extended Mappings, IOP Conf. Series: Journal of Physics: Conf. Series 1003 (2018) 012042. https://doi.org/10.1088/1742-6596/1003/1/012042.

[18] Zeidler ,"Nonlinear Functional Analysis and Application" New York, (1986). https://doi.org/10.1007/978-1-4612-4838-5.

[19] H.H. Bauschke and P.L. Combettes, Convex Analysis and Monotone Operator Theory in Hilbert Spaces, Springer-Verlag, 2011. https://doi.org/10.1007/978-1-4419-9467-7.

[20] B.E. Rhoades, Comments on two fixed point iteration methods, J. Math. Anal. Appl. 56 (1976) 741–750. https://doi.org/10.1016/0022-247X(76)90038-X.

[21] W.G. Dotson Jr., On the Mann iterative process, Trans. Amer. Math. Soc. 149 (1970)65–73. https://doi.org/10.1090/S0002-9947-1970-0257828-6.

[22] R.L. Franks, R.P. Marzec, A theorem on mean value iterations, Proc. Amer. Math. Soc. 30 (1971) 324–326. https://doi.org/10.1090/S0002-9939-1971-0280656-9.

[23] C.W. Groetsch, A note on segmenting Mann iterates, J. Math. Anal. Appl. 40 (1972) 369–372. https://doi.org/10.1016/0022-247X(72)90056-X.

[24] S. Ishikawa, Fixed points and iteration of a nonexpansive mapping in a Banach space, Proc. Amer. Math. Soc. 59 (1976) 65–71. https://doi.org/10.1090/S0002-9939-1976-0412909-X.

[25] A.K. Kalinde, B.E. Rhoades, Fixed point Ishikawa iterations, J. Math. Anal. Appl. 170 (1992) 600–606. https://doi.org/10.1016/0022-247X(92)90040-K.
[26] K.K. Tan, H.K. Xu, Approximating fixed points of nonexpansive mappings by the Ishikawa iteration process, J. Math. Anal. Appl. 178 (1993) 301–308. https://doi.org/10.1006/jmaa.1993.1309.
[27] M.K. Ghosh, L. Debnath, Convergence of Ishikawa iterates of quasi-nonexpansive mappings, J. Math. Anal. Appl. 207 (1997) 96–103. https://doi.org/10.1006/jmaa.1997.5268.
[28] C.E. Chidume, S.A. Mutangadura, An example of the Mann iteration method for Lipschitz pseudocontractions, Proc. Amer. Math. Soc. 129 (2001) 2359–2363. https://doi.org/10.1090/S0002-9939-01-06009-9.
[29] Z. H. Maibed,generalized tupled common fixed point theorems for weakly compatibile mappings in fuzze metric space,(IJCIET)10(2019)255-273.