International Journal of Advanced Chemistry, 2 (1) (2014) 39-43

SPC

International Journal of Advanced Chemistry

Journal home page: www.sciencepubco.com/index.php/IJAC
doi: 10.14419/ijac.v2i1.1734
Research Paper



Kinetics and mechanism of oxidation of GSH by cis-(diaqua)-bis-(ethylenediamine) Cobalt(III) ion

B. Mohanty [1*], J. Behera [1], S. Acharya [2], P. Mohanty [1], A.K. Patnaik [3]

[1] *Department of Chemistry, Utkal University, Bhubaneswar, 751004, Odisha, India*
[2] *IMMT, Bhubaneswar, Odisha, India*
[3] *Ravenshaw University, Cuttack, 753003, Odisha, India*
**Corresponding author E-mail: omm.gayatri@gmail.com*

Abstract

The kinetics of GSH oxidation to GSSG by cis-(diaqua)-bis-(ethylenediamine) cobalt(III) perchlorate was studied spectophotometrically under pseudo-first order condition using 10^3[Co(III)] = 5 mol dm^{-3}, $2.5 \leq 10^2$[GSH] ≤ 10.00 mol dm^{-3}, $3.5 \leq$ pH ≤ 5.0, $318K \leq T \leq 333K$ at a fixed ionic strength I = 0.3 mol dm^{-3} ($NaClO_4$). The disappearance of [Co (III)] at 500 nm with time showed first-order kinetic trend. The first order dependence on [GSH] and pH was observed. Temperature dependence of the second order rate constant $k'_2 = k_{obs}$/[GSH] were analysed for the Co(III)-OH_2^{3+} (k_1) and Co(III)-OH^{2+} (k_2) reactivities. An outer-sphere complex formation between the Co(III) and GSH followed by one-electron transfer from GSH to Co(III) resulting the formation of Co(II) and GS^x radical. The GS^x radical undergoes fast dimerisation to GSSG. Activation parameters were calculated. These values favor the electron transfer reaction.

Keywords: *Co (III) complex, GSH, GSSG, Spectrophotometer, Redox reaction.*

1. Introduction

Glutathione (GSH) is a tripeptide containing three amino acids L-cysteine, L-glutamic acid and L-glycine with unusual peptide linkage between amine group of cysteine and carboxylic group of glutamate. The thiol group (SH) of cysteine serves as proton donor and is responsible for the biological activity of glutathione. It plays a fundamental role in numerous metabolic and biochemical reactions such as DNA synthesis and repair, protein synthesis, amino acid transport and enzyme activation.

Because of the importance of GSH, we are interested to study the kinetics of oxidation of GSH by biological active cobalt complex. Cobalt is an essential trace element for all organisms as the active center of co-enzyme called cobalamines. This includes vitamin B_{12} which is essential for mammals. A deficiency of cobalt leads to pernicious anaemia. Studies on the chemistry of electron transfer reaction of cobalt(III) complexes have received a sustained high level of attention from the scientific community for decades due to their relevance in various redox processes in biological system and act as promising agent for antitumor (Osinsky *et al*. 2003, Osinsky *et al*. 2004), anthelmintic (Behm *et al*. 1993), antiparasitic (Behm *et al*. 1995, Karaman *et al*. 1995), antibiotics (Ghirlanda *et al*. 1998), and antimicrobial activities (Srinivasan *et al*. 2005). Numerous studies have been performed, addressing the dependence of electron transfer on different environments including metalloproteins (Bernauer *et al*. 1999), vitamin B_{12} (Wolak *et al*. 2003) , liquids (Burel *et al*. 1999, Saik *et al*. 2004), micelles (Weidemaier *et al*. 1997, Travernier *et al*. 1998), vesicles (Gerasimov et al. 1988), and DNA (Srinivasan *et al*. 2005).

We report here redox reactions of GSH with Co (III) complex as oxidant and GSH as reductant. The product was isolated, identified and mechanism was proposed.

2. Experimental

2.1. Material and methods

Cis -(diaqua) - bis - (ethylenediamine) cobalt (III) perchlorate was prepared following the reported method (Basolo & Pearson 1964, Krisnamurty 1972). It was recrystallised and dried. The solid complex was characterized by elemental analysis and spectroscopic methods. All other chemicals used were of AnalaR grade. Doubly distilled water was used to prepare all the solutions. The acid strength of the medium was maintained by addition of $HClO_4$. The UV-Vis spectra of the complex showed the characteristic peaks at 380 nm (ε = 65 dm^{-3} mol^{-1} cm^{-1}) and 500 nm (ε = 80 dm^{-3} mol^{-1} cm^{-1}).

2.2. Kinetic study

Kinetics of oxidation of GSH by Co (III) complex in perchloric acid medium was studied from 318 to 333K, I = 0.3 mol dm^{-3} ($NaClO_4$). The kinetic measurements were carried out with a systronic 2202 UV-Vis spectrophotometer equipped with a thermostatic water bath for temperature control (accuracy = 0.1^oC). The progress of the reaction was monitored by following the decrease in absorbance at 500 nm "Fig. 1". The detail experimental procedure of kinetic study was reported in our published paper (Mohanty *et al.* 2011). The kinetics were studied under pseudo-first order conditions, where the substrate[GSH] varied from 2.5 x 10^{-2} mol dm^{-3} to 10.0 x 10^{-2} mol dm^{-3} and pH was varied from 3.5 to 5.0 and the [Co(III)L] complex is 5 x 10^{-3} mol dm^{-3}. The rate constants (k_{obs}) were calculated from the slope of ln (A_t - A_∞) *versus* t (s) plots from the relationship using computer program.

Copyright © 2014 B. Mohanty et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

$\ln (A_t - A_\infty) = \ln (A_0 - A_\infty) - k_{obs}.t$

Where A_0, A_t, A_∞ denote optical density of the reaction mixture at zero time, time *t* and at infinite time respectively. A_∞ was measured after completion of the reaction. The correlation coefficients of plots used to determine k_{obs} were found to be 0.99 in most of the cases.




Fig. 1: UV-Vis spectral scan of GSH with Co (III) at 303K, pH = 4.0, Co (III) = 0.005 mol dm^{-3}, I = 0.3 mol dm^{-3}($NaClO_4$), (1) Immediate after mixing, curves (2, 3) Δt = 5 mins, curves (4, 5) Δt = 10 mins, (6) after 24hs

3. Stoichiometry and characterization of the product

The reaction mixture containing [substrate] and [Co (III) L] in ratio 1:10, I = 0.3 mol dm^{-3} ($NaClO_4$) were allowed to react till completion of the reaction. From the above stoichiometry study it was revealed that the reaction exhibited as 1:1 stoichiometry for GSH oxidation.

$2\ Co^{3+} + 2\ e = 2\ Co^{2+}$

$2\ GSH = GSSG + 2\ H^+ + 2\ e$

$2\ Co\ (III) + 2\ GSH + 2\ OH^- = 2\ Co\ (II) + GSSG + 2\ H_2O$

To prepare the reaction product [Co (III) L] = 0.05 mol dm^{-3} was mixed with hot solution of [$HClO_4$] = 0.005 mol dm^{-3}, then pH of the solution was adjusted to 4.5 (Sol-I). In a separate beaker [GSH] = 0.05 mol dm^{-3} was mixed with $NaClO_4$ = 0.3 mol dm^{-3}. Its pH was adjusted to 4.5 (Sol-II). Both the solutions (I and II) were mixed and evaporated in a thermostat at 60°C for 3h till a pasty solution was formed. Then it was slowly dried in a desiccator. The yield of the product was nearly 70%. The FTIR spectra of the dried product were recorded "Fig. 3" in a Perkin Elmer (UK) FTIR spectrophotometer using KBr pellet technique and it was compared with FTIR spectra of the substrate GSH "Fig. 2". Co (II) in the product was identified by Kitson method (Kitson 1950).




Fig. 2: FTIR spectra of pure GSH






Fig. 3: FTIR spectra of oxidation product GSSG

"Fig. 3" showed a broad peak at 3396.54 cm^{-1} in the product may be assigned to υ_{N-H} (NH_3^+) as compared to 3252 cm^{-1} and 3026 cm^{-1} in GSH. The shifting to higher frequency was probably due to an association of water molecules with the product. The NH_3^+ bending bands and a strong absorption peak of carboxylate ion are mixed up and a broad band is observed at 1650.93 cm^{-1} in the product compared to 1600 cm^{-1}, 1538 cm^{-1} and 1395 cm^{-1} peak in GSH. The strong peaks at 1541.93 cm^{-1}, 1403.78 cm^{-1}, 1227.83 cm^{-1} and 537.06 cm^{-1} in the product corresponds to 1599.66 cm^{-1}, 1383.54 cm^{-1}, 1278.57 cm^{-1} and 549.61 cm^{-1} in GSH is due to the

free NH_2 twisting and rocking (Nakamoto 5th Edn). The weak band at 2526 cm^{-1} in GSH due to S-H stretching is absent in the product suggesting the dimerisation of GSH to GSSG having S-S linkage. The product was isolated as GSSG. [(2S)-2-amino-5-[[(2R)-3-[(2R)-2-[[(4S)-4-amino-5-hydroxy-5-oxopentanoyl]amino]-3-[carboxymethylamino)-3-oxopropyl]disulfanyl-1-(carboxymethylamino)-1-oxopropan-2-yl]-5-oxopentanoic acid]. The structure of substrate (GSH) and product (GSSG) was shown as



Structure of substrate GSH



Structure of product (GSSG)

4. Results

All kinetic runs were performed under pseudo-first order conditions.

4.1. Effect of [GSH] on reaction rate

With the varying concentration of [GSH] = 2.5 x 10^{-2} to 10.0 x 10^{-2}mol dm^{-3},10^4 k_{obs}(s^{-1}) (323K) increased from 6.8 to 10.42 mol dm^{-3} when pH = 4.0, I = 0.3mol dm^{-3} and [Co (III) L] = 5.0 x 10^{-3} mol dm^{-3} "Table 1". The plot of k_{obs} *versus* [GSH] "Fig. 4" was linear at different temperatures indicating first order dependence of rate on [GSH]. Furthermore, the second order rate constant k_2' ($mol^{-1}dm^3s^{-1}$) = k_{obs}/ $[GSH]_T$ almost remains constant. As the order of the reaction is one with respect to $[GSH]_T$, the order of the reaction is one with respect to [Co(III)L]. The rate law is therefore expressed as

Rate = k_{obs} [Co^{III}-OH]




Fig. 4: Plot of k_{obs} vs [GSH] at I = 0.3 mol dm^{-3}, temp=318 K (1), 323K (2), 328K (3) and 333K (4)

4.2. Effect of pH on reaction rate

The effect of pH on reaction rate was studied varying pH = 3.5 to 5.0, [Co (III) L] = 5 x 10^{-3} mol dm^{-3}, [GSH] = 2.5 x 10^{-2} mol dm^{-3} and I = 0.3 mol dm^{-3}. The rate10^4 k_{obs} (318K) increased from 2.48 to 6.8 mol dm^{-3} as pH was increased from 3.5 to 5.0. This behavior was repeated for the entire $[GSH]_T$ range 0.025 to 0.1 mol dm^{-3}. Plots of k_{obs}/[GSH] *versus* [H^+] "Fig 5" was a straight line indicating first order dependence of the rate on [H^+]. The rate constants k_1 and k_2 were obtained from plots of k_{obs}/ [GSH] *versus* [H^+].




Fig. 5: Variation of $k'_2 = k_{obs}$ / [GSH] vs [H^+] at 328K, [GSH] = 0.05 mol dm^{-3}, I = 0.3 mol dm^{-3}

4.3. Effect of temperature on reaction rate

The rate of the reaction was studied varying the temperature 318K to 333K, [Co (III) L] = 5 x 10^{-3} mol dm^{-3}, $[GSH]_T$ = 2.5 x 10^{-3} mol dm^{-3}, I = 0.3 mol dm^{-3}, pH = 3.5. Pseudo-first order rate constant k_{obs}(s^{-1}) were found to increase from 2.48 x 10^{-4} to 10.71 x 10^{-4} mol dm^{-3}as temperature increased from 318K to 333K, "Table 1".

Table 1: Pseudo-first order reaction rate constants of oxidation of GSH at different temperatures

[GSH] (mol dm^{-3})	pH	10^4[H^+] (mol dm^{-3})	10^4 k_{obs} (s^{-1})			
			318K	323K	328K	333K
0.025	3.5	3.160	2.48	5.20	7.55	10.71
0.025	4.0	1.000	3.65	6.80	12.22	12.23
0.025	4.5	0.316	6.53	8.67	11.08	13.95
0.025	5.0	0.100	8.35	10.65	13.11	15.86
0.05	3.5	3.160	3.97	6.58	8.74	12.04
0.05	4.0	1.000	5.23	7.87	11.51	14.15
0.05	4.5	0.316	8.10	10.30	13.45	15.81
0.05	5.0	0.100	9.90	12.58	15.24	17.48
0.075	3.5	3.160	5.58	8.06	10.54	13.55
0.075	4.0	1.000	7.03	9.57	13.29	16.04
0.075	4.5	0.316	9.43	12.17	14.66	17.61
0.075	5.0	0.100	11.23	14.37	16.89	19.56
0.10	3.5	3.160	6.80	9.32	12.33	14.82
0.10	4.0	1.000	8.32	10.42	15.05	17.93
0.10	4.5	0.316	11.15	13.58	16.00	19.44
0.10	5.0	0.100	12.60	15.90	18.21	21.10

[Co^{III}L] = 5 x 10^{-3} mol dm^{-3}, I = 0.3 mol dm^{-3}

5. Discussion


The oxidation reaction was occurred through two paths with the rate constants k_1 and k_2. Here k_2 path was almost remains constant. So the activation parameters were determined from the electron transfer rate constants k_1. The values of $\Delta H_1^{\#}$ and $\Delta S_1^{\#}$ were found to be 66.15 ± 9.5 kJ mol^{-1} and -77.77 ± 29.2 JK^{-1} mol^{-1} respectively. The moderate values of activation parameters favor the electron transfer reaction. The negative values of activation parameters suggested the formation of the ordered transition state.

Since pK_1, pK_2, pK_3 and pK_4 of GSH are 2.05, 3.47, 8.63 and 9.52(Scheme 1), at higher concentration of the acid 0.03 mol dm^{-3}, the undissociated form of GSH will participate in the electron

transfer reaction. The reaction sequence delineated below "Scheme 2" was consistent with the experimental data.

Mechanism

SCHEME – 1



Deprotonation equilibria of GSH

The above observation suggested the following mechanism as shown in "scheme 2". From the above reaction the rate law was derived as

SCHEME-2

Proposed mechanism of electron transfer reaction

$$Co^{III}\text{-}OH_2 \overset{K_h}{\rightleftharpoons} Co^{III}\text{-}OH + H^+$$

$$GSH \xrightarrow{k_1} GS^{\cdot} + Co(II) \qquad GHS \xrightarrow{k_2} GS^{\cdot} + Co(II)$$

$$2\,GS^{\cdot} = GSSG \text{ (fast)}$$

From the proposed mechanism, the rate law was derived as

$$K_h = \frac{Co^{III}\text{-}OH_2}{Co^{III}\text{-}OH_2} \qquad (1)$$

$$\text{Rate} = k_1 [GSH] [Co^{III}\text{-}OH_2]_e + k_2 [GSH] [Co^{III}\text{-}OH]_e$$

$$= k_1 [GSH] [Co^{III}\text{-}OH_2]_e + k_2 K_h [Co^{III}\text{-}OH_2]_e [GSH] / [H^+]$$

$$= [GSH] [Co^{III}\text{-}OH_2]_e \{k_1 [H^+] + k_2 K_h\} / [H^+] \qquad (2)$$

$$[Co^{III}\text{-}OH_2]_T = [Co^{III}\text{-}OH_2]_e + [Co^{III}\text{-}OH]_e$$

$$= [Co^{III}\text{-}OH_2]_e + \{[Co^{III}\text{-}OH_2] K_h / [H^+]\}$$

$$= [Co^{III}\text{-}OH_2]_e \{[H^+]+K_h\}/[H^+] \qquad (3)$$

$$[Co^{III}\text{-}OH_2]_e = \{[Co^{III}\text{-}OH_2]_T [H^+]\}/\{[H^+]+K_h\} \qquad (4)$$

$$\text{Rate} = [GSH]_T [Co^{III}\text{-}OH]_T \{k_1[H^+] + k_2 K_h\}/\{[H^+] + K_h\} \qquad (5)$$

$$\text{Rate} = k_{obs} [Co^{III}\text{-}OH]_T \qquad (6)$$

Comparing (5) & (6)

$$k_{obs} = [GSH]_T \{k_1[H^+]+k_2 K_h\}/\{[H^+]+K_h\} \qquad (7)$$

$$\frac{k_{obs}}{[GSH]_T} = k_2' = \frac{\{k_1[H^+]+k_2 K_h\}}{k_2'\{[H^+] + K_h\}}$$

$$k_2'\{[H^+] + k_h\} = \{k_1[H^+]+k_2 K_h\} \qquad (8)$$

Where $k_2' = k_{obs} / [GSH]_T$

There exists two species of Co (III) as $Co^{III} - OH_2$ and $Co^{III} - OH$ in acid solution which was in equilibrium with each other. Both the species reacted with GSH through two paths with rates k_1 and k_2 respectively producing radicals GS^x and Co (II). The GS^x radicals dimerised rapidly forming GSSG.

From equation (8) the left hand side expression was plotted against $[H^+]$. It showed a straight line with a positive slope. From the slope and intercept k_1 and k_2 were calculated. These values at four different temperatures 318 – 333K were calculated and tabulated in "Table 2". It shows that k_2 rate was much faster than k_1.

The values of k_1 (rate of oxidation of GSH by Co^{III}-OH_2 species) varies with temperature whereas k_2 (rate of oxidation of GSH by Co^{III}-OH species) remained almost constant and the activation parameters for the path k_1 were calculated and tabulated "Table 2". The activation parameters were determined from the electron transfer rate constants k_1.

Table 2: Calculation of electron transfer rate constants and activation parameter

Temp (K)	pK_h	k_1 $mol^{-1}dm^3s^{-1}$	k_2 $mol^{-1}dm^3s^{-1}$
318	5.65	0.0073	0.127
323	5.53	0.0128	0.112
328	5.41	0.0176	0.105
333	5.30	0.0242	0.084
	$\Delta H^{\#}$ =	(66.15 ± 9.15) $kJmol^{-1}$	
	$\Delta S^{\#}$ =	(-77.77 ± 29.2)$JK^{-1}mol^{-1}$	

6. Conclusion

The oxidation reaction of GSH by Co (III) complex proceeded through two steps (k_1 and k_2) producing free radical as GS^x and Co (II) respectively. The free radical GS^x dimerised to GSSG. Reduction at Co (III) center has been achieved due to generation of a radical at the bound ligand by the one equivalent oxidant. k_2 path is much faster than k_1 path. The activation parameters corresponding to k_1 were evaluated, such as activation enthalpy ($\Delta H^{\#}$ = 66.15 ± 9.5 kJ mol^{-1}) and activation entropy ($\Delta S^{\#}$ = -77.77 ± 29.2 JK-1 mol-1). Moderate values of activation parameters favor the electron transfer process. Negative value of activation entropy corresponds to ordered transition state.

Since there was no experimental evidence of bridging of ligand between the oxidant and reductant, inner sphere mechanism was ruled out. The electron transfer mechanism between GSH and Co (III) complexes was expected to be outer sphere mechanism.

Kinetics of oxidation of tripeptide (GSH) was compared with the kinetics of oxidation of one of its amino acid L – cysteine (Asemave *et al*. 2012) by Co (III) complex under similar conditions. It was concluded that the rate of electron transfer reaction of the tripeptide GSH is 40 times slower than amino acid.

Acknowledgement

The authors are thankful to H.O.D Chemistry, Utkal University, Vani Vihar, Bhubaneswar, Odisha, India for providing research facilities.

References

[1] Asemave K, Yiase SG, Adejo SO (2012) Kinetics and Mechanism of Substitution Reaction of Trans-Dichloro-bis-(Ethylenediammine) Cobalt (III) Chloride with Cysteine, Aspartic acid and Phenylalanine, International Journal of Science and Technology, 2(5):242-247.

[2] Basolo F and Pearson RG (1964) Mechanism of Inorganic Reaction, Wiley.

[3] Behm CA, Boreham PFL, Creaser II, Korybutdaszkiewicz B, Maddalena DJ, Sargeson AM & Snowdown GM (1995) Novel Cationic Surfactants Derived From Metal Ion Cage Complexes: Potential Antiparasitic Agents, Australian Journal of Chemistry, 48(5):1009-1030

[4] Behm CA, Creaser I, Daszkiewicz B, Geue RJ, Sargeson AM & Walker GW (1993) Novel cationic surfactants derived from metal ion cage complexes: potential anthelmintic agents, Journal of the Chemical Society, Chemical Communications, 24:1844-1846.

[5] Bernauer K, Ghizdavu S & Verardo L (1999) Chiral metal complexes as probes in electron-transfer reactions involving metalloproteins, Coordination Chemistry Reviews, 190-192:357–369.

[6] Burel L, Mostafavi M, Murata S, Tachiya M (1999) Transient Effect in Fluorescence Quenching by Electron Transfer. 4. Long-Range Electron Transfer in a Nonpolar Solvent, The Journal of Physical Chemistry A, 103(30):5882-5888.

[7] Gerasimov OV, Lymar SV, Tsevetkov TM, Parmon VN(1988) Cobalt water oxidation catalyst immobilized on membranes of lipid vesicles, Reaction Kinetics and Catalysis Letters, 36(1):145-149

[8] Ghirlanda G, Scrimin P, Tecilla P & Toffoletti A (1998) Amphiphilic Copper(II) Complexes Modeled after the Metal-Complexation Subunit of Bleomycin Antibiotics†, Langmuir, 14 (7):1646-1655.

[9] Karaman ME, Pashley RM & Bolonkin NK (1995) Study of the Surface and Biological Activity of a Trivalent Cage Surfactant, *Langmuir*, 11 (8):2872–2880.

[10] Kitson RE (1950) Simultaneous Spectrophotometric Determination of Cobalt, Copper, and Iron, Analytical Chemistry 22:664-667.

[11] Krisnamurty MJ (1972)Modified synthesis of trans-dichloro-bis-ethylenediamine cobalt(III) chloride, Journal of Inorganic and Nuclear Chemistry, 34:3915–3916

[12] Mohanty B, Behera J, Mohanty P, Acharya S, Patnaik AK & Das SP(2011) Kinetics and mechanism of oxidation of L-cysteine and DL-methionine be morpholinium chlorochromate in aqueous acidic medium, Journal of Indian Chemical Society, 88(10):1561-1566.

[13] Nakamoto K (1997) Infrared and Raman Spectra of Inorg. And Coord. Compexes, 5th Edn, John Willey and Sons Inc. Publication.

[14] Osinsky S, Levitin I, Bubnovskaya L, Sigan A, Ganusevich I, Kovelskaya A, Valkovaskaya N, Campanella L & Wardman P (2004) Selectivity of effects of redox-active cobalt(III) complexes on tumor tissue, Experimental Oncology, 26:140-144.

[15] Osinsky S, Levitin I, Sigan A, Bubnovskaya L, , Ganusevich I, Campanella L & Wardman P (2003) Redox-active cobalt complexes as promising antitumor agents, Russian Chemical Bulletin, 52:2636-2645.

[16] Saik VO, Goun AA, Nanda J, Shirota K, Tavernier HL & Fayer MD (2004) Photoinduced Intermolecular Electron Transfer in Liquid Solutions, The Journal of Physical Chemistry A, 108:6696-6703.

[17] Srinivasan S, Annaraj J & Athappan PR (2005) Spectral and redox studies on mixed ligand complexes of cobalt(III) phenanthroline/bipyridyl and benzoylhydrazones, their DNA binding and antimicrobial activity, Journal of Inorganic Biochemistry, 99:876-882.

[18] Tavernier HL, Barzykin AV, Tachiya M & Fayer MD, (1998)Solvent Reorganization Energy and Free Energy Change for Donor/Acceptor Electron Transfer at Micelle Surfaces: Theory and Experiment, The Journal of Physical Chemistry B, *102* (31):6078–6088

[19] Weidemaier K, Tavernier HL & Fayer MD (1997) Photoinduced Electron Transfer on the Surfaces of Micelles, The Journal of Physical Chemistry B, 101:9352-9361.

[20] Wolak M, Stochel G, Eldik Rv (2003) Mechanistic Studies on the Interaction of Reduced Cobalamin (Vitamin B12r) with Nitroprusside, Journal of the American Chemical Society, 125:1334-1351.