



Pharmacologically potentials of hydrazone containing compounds: a promising scaffold

Mohammad Asif

Department of Pharmacy, GRD (PG) IMT, Dehradun, (Uttarakhand), 248009, India

E-mail: aasif321@gmail.com

Abstract




Hydrazone considered as an important class of compounds for new drug development. Hydrazones are present in many of the bioactive compounds having wide interest because of their diverse biochemical applications. This created interest in researchers to developed variety of hydrazone compounds with effective biological activities. Therefore many compounds have been developed as target structures for their biological activities. This review is focus on the diverse biological activities of hydrazones.

Keywords: Hydrazones, Biological Activities, Azomethine, Pharmacological Potential.

1. Introduction

Hydrazones have wide interest because of their diverse biological applications. Hydrazones possess various biological activities like anticonvulsant, antidepressant, analgesic, anti-inflammatory, antiplatelet, antimalarial, antimicrobial, antitubercular, anticancer, vasodilator, antiviral, antischistosomiasis, anti-HIV, anthelmintic, antidiabetic, and trypanocidal activities (Ali et al., 2012; Kumar et al., 2010). Hydrazone-hydrazones are not only intermediates but they are also effective organic compounds. Some effective compounds like iproniazide and isocarboxazide. Iproniazid, like isoniazid is used in the treatment of tuberculosis (TB) and also showed antidepressant effect. Another effective


hydrazidehydrazones is nifuroxazide, which is used as an intestinal antiseptic (Negi et al., 2012; Rollas and Kucukguzel. 2007). The 2-Chloroquinoliny hydrazone derivative are anticonvulsants and ribavirin hydrazone derivatives are anticancer (Liu et al., 2009), hydrazones of indane-1, 3-dione are anticoagulant and antimicrobial activity (Jubie et al., 2010), 4-arylhydrazono-2-pyrazoline-5-one derivatives are anti-TB activity (Guniz et al., 2007). Some hydrazide hydrazones were active against Mtb H37Rv between the concs of 0.78-6.25µg/ml (Kaymakçioğlu, et al., 2006). A series of hydrazide-hydrazones reported antidepressant, sedative and analgesic activities (Mohareb et al., 2010).




2. Antimicrobial activity

The antibacterial activity of hydrazine derivatives [1] against various pathogenic bacterial strains (Sharma et al., 2011) and other hydrazine derivatives [2], [3] are also showed antibacterial activity (Jubie et al., 2010). Benzimidazole derivatives with hydrazone moiety [4] (Ozkay et al., 2010), chloropyrrole derivatives of aroylhydrazone [5] (Rane and Telvekar. 2010) and some hydrazoneyl substituted pyrimidinones [6], [7] showed antibacterial activity (Edrees et al., 2010). Vanillin based hydrazines [8], [9] showed antibacterial activity against *S. aureus* and *P. aeruginosa* (Govindasami et al. 2011). 2-quinoxalinone-3-hydrazine derivative [10] exhibited antibacterial activities against different bacterial strains (Ajani et al., 2010). Various hydrazones [11], [12] were act as selective inhibitors of *S. aureus* β -ketoacyl carrier proteinsynthase-3 (Lee et al., 2012). Aryloxyacetic acid


hydrazide [13] having promising antibacterial activity against some bacterial strains (Abdel-Wahab et al., 2012). Cholic acid based hydrazones [14] showed antibacterial activity against *Escherichia faecalis* and *E. coli* (Rasras et al., 2010). Anthraquinone based hydrazones [15] showed promising bacteriostatic activity against *P. aeruginosa* (Gouda et al., 2010). Various benzylidinehydrazides [16] showed bactericidal activity against *S. aureus* (Kumar et al., 2010). Some hydrazone with imidazoles [17] were exhibited antibacterial activity against various bacteria (Abdel-Wahab et al., 2011). The 2, 4-disubstituted thiazoles [18] have antibacterial activity against different strains (Vijesh et al., 2010). The 4-fluorobenzoic acid (5-nitro-2-furyl)-methylene-hydrazide, is effective against *S. aureus* (Gülerman et al., 2000).



1




2




3


4




5




6




7




8




9




10




11




12




13




14















15



16



17







Few hydrazone derivatives [Raja et al., 2010], 4-OH-8-trifluoromethylquinoline derivatives [19] (Thomas et al., 2011), fluorine-containing hydrazones [20] (Vavrikova et al., 2011) and some hydrazines [21], [22] and [23] showed promising antibacterial activity (Pavan et al., 2010). Ferrocenyl hydrazones [24], [25] (Magueue et al., 2011) and ferrocene-based hydrazone derivatives [26] showed significant antibacterial potential (Mahajan et al., 2011). Hydrazones [27] (Eswaran et al., 2010) and hydrazone derivatives [28] showed anti-TB activity (Jordao et al., 2011). Hydrazone derivatives [29] against different *Candida* spp, have promising antifungal potential (Ozdemir et al., 2010). Hydrazone derivatives [30], [31] have antiviral activity against viral HIV1-CA (Jin et al., 2010).

Various 2-quinoxalinone-3-hydrazone derivatives showed antimicrobial activities against the various pathogenic microbes, compounds 3-{2-(1-(6-chloro-2-oxo-2H-chromen-3-yl) ethylidene)

hydrazinyl}quinoxalin-2(1H)-one [32], 3-(2-(propan-2-ylidene)hydrazinyl) quinoxalin-2(1H)-one [33] showed antibacterial and antifungal activity (Olayinka et al., 2010). A aryl-hydrazone, N-(2-pyridinecarbaldehyde(-N-(4-(4-chlorophenyl)sulfonyl) benzoyl) hydrazone and its Cu(II), Co(II) and Ni(II) complexes showed antibacterial activity against Gram positive bacteria: *S. aureus*, *B. subtilis* and Gram-negative bacteria: *P. aeruginosa*, *E. coli*. The activity of N-(2-pyridinecarbaldehyde)-N-(4-(4-chloro-phenyl sulfonyl) benzoyl)-hydrazone becomes more pronounced when coordinated to the metal ions (Angelusiu et al., 2010). Sulfonyl hydrazone derivatives and their nickel complexes were exhibited antibacterial activity against gram-positive bacteria (*S. aureus*, *B. subtilis*, *B. magaterium*) and gram-negative bacteria (*Salmonella enteritidis*, *E. coli*) and possess broad spectrum of activity at MIC values between 145 and 683 µg/ml and presence of electron rich atoms (O, S, N, etc.) in ligands increase the antimicrobial activity

(Ozdemir and Gulcin. 2008). Biphenyl-4-carboxylic acid hydrazide hydrazone exhibited in-vitro antimicrobial activity against two gram negative strain *E. coli*, *P. aeruginosa* and two Gram positive strain, *B. subtilis*, *S. aureus* and fungal strain, *C.*




albicans, *A. niger* and presence of electron withdrawing groups on aromatic ring improved the antimicrobial activity (Deep et al., 2010).



A series of 2-arylquinoline-4-carboxylic acid hydrazidhydrazones were exhibited in vitro antimicrobial activity against *S. aureus*, *E. coli* and *C. albicans*. Out of these compounds 6-chloro-2-(4-methoxyphenyl) quinoline-4-carboxylic acid (4-nitrobenzylidene)-hydrazide [34] was found to be most potent (Metwally et al., 2006). Some p-substituted benzoic acid ((5-nitro-thiophen-2-yl)-methylene)-hydrazides were showed antimicrobial activity against standard and MDR strains of *S. aureus*. The 4-Acetyl-benzoic acid ((5-nitro-thiophen-2-yl)-methylene)-hydrazide [35] was the most active compound (Masunari, and Tavares. 2007). Derivatives of acylhydrazine such as substituted-2-mercapto-1, 3, 4-oxadiazoles, their corresponding ester, amide and benzene diasulfonamides were showed antimicrobial activity, the acyl hydrazine benzene diasulfonamide derivative was having antimicrobial activity (Zareef et al., 2008).


Anti-TB Activity: Various diclofenac acid hydrazones and amides were exhibited anti-TB activities against *Mtb*. The 1-cyclopropyl-6-fluoro-8-methoxy-7-((N4-(2-(2-(2,6-dichloro phenyl amino)phenyl)acetyl)-3-methyl)-N1-piperazinyl)-4-oxo-1,4-dihydro-

3-quinoline carboxylic acid [37] was found to be the most active compound in-vitro than anti-TB drug Isoniazid (INH). In the in vivo model, these compounds decreased the bacterial load in lung and spleen tissues (Sriram et al., 2006). Two series of 2-substituted quinolines 4-(adamantan-1-yl) group were exhibited in-vitro anti-TB activities against drug-sensitive *Mtb* H37Rv strain. Compound 4-adamantan-1-yl-quinoline-2-carboxylic acid (2-chlorobenzylidene) hydrazide [37] inhibited drug sensitive *Mtb* H37Rv and equipotent to INH (Nayyar et al., 2007). During the coumarin-4-acetic acid benzylidene hydrazides as anti-TB agents against *Mtb* H37Rv strain, (7-Hydroxy-2-oxo-2H-chromen-4-yl)-acetic acid (3-nitro-benzylidene) hydrazide [38] was found to be most potent (Manvar et al., 2008). Some 5, 6, 7, 8-tetrahydronaphthalen acetic acid benzylidene hydrazide derivatives showed anti-TB activity and compounds having -OH and -NO₂ in ortho and Meta position of aromatic ring having high anti-TB activity (Ozemir et al., 2010). Aryl acetic acid hydrazones showed moderate activity against *Mtb* H37 RV strain (Srivastava et al., 2010).




Some hydrazone compounds [39], [40] and N1-(4-methoxy benzamido) benzoyl)-N2-((5-nitro-2-furyl) methylene) hydrazine 41 inhibited the growth of several bacteria and fungi (Kuçukguzel et al., 2002). N2-Substituted alkylidene/arylidene-6-phenylimidazo [2], [1-b] thiazole-3-acetic acid hydrazides 42 were evaluated for their in vitro antimicrobial activity (Ulusoy et al.,

2000). The 5-bromoimidazo [1], [2-a] pyridine-2-carboxylic acid benzylidene hydrazide 43 and 5-bromoimidazo [1], [2-a] pyridine-2-carboxylic acid 4-methoxy benzylidene hydrazide 44 to possess antimicrobial activity against *E. fecalis* and *S. epidermis* (Turan-Zitouni et al., 2001).



A series of hydrazones derived from 1, 2-benzisothiazole hydrazides ($R_1=H$) 23-27 as well as the parent cyclic [45] and [46] and acyclic [47], [48] and [49] 1, 2-benzisothiazole hydrazides, were showed antibacterial and antifungal activity. The 2-amino-1,2-benzisothiazole-3(2H)-one derivatives, compound 45 and 46 showed antibacterial activity against Gram positive bacteria and most of them were also active against yeasts (Vicini et al., 2002). A series of hydrazide hydrazones and 1, 3, 4-oxadiazolines of 4-fluorobenzoic acid hydrazide as antibacterial and antifungal activities against *S. aureus*, *E. coli*, *P. aeruginosa* and *C. albicans*. The compound 4-fluorobenzoic acid ((5-nitro-2-furyl) methylene) hydrazide [50] showed equal activity as ceftriaxone against *S.*

aureus (Rollas et al., 2002). The 2', 4'-Difluoro-4-hydroxybiphenyl-3-carboxylic acid ((5-nitro-2-furyl) methylene) hydrazide has shown activity against *S. epidermis* HE-5 and *S. aureus* HE-9, respectively. 2', 4'-Difluoro-4-hydroxybiphenyl-3-carboxylic acid ((2, 4, 6-trimethyl phenyl) methylene) hydrazide [51] has exhibited activity against *Acinetobacter calcoaceticus* (Kucukguzel, et al., 2003). A series of hydrazones from cholesterol derivatives 52 were exhibited antimicrobial properties against pathogens. The tosylhydrazone cholesterol derivative exhibited activity against *C. albicans* (Loncle et al., 2004). 4-Substituted benzoic acid ((5-nitro-thiophene-2-yl) methylene) hydrazides 53 were exhibited bacteriostatic and some compounds showed bactericidal activity (Masunari, and Tavares. 2007).




Isoniazid (INH) hydrazide-hydrazones 1, the compounds have activity against various strains of Mtb, with lower toxicity than INH. The 1-methyl-1H-2-imidazo[4, [5-b]pyridinecarboxylic acid hydrazides, compound 54 showed anti-TB activity against Mtb H37 Rv, Mtb 192, Mtb 210, isolated from patients and resistant against INH, ethambutol, RIF at 31.2 $\mu\text{g/mL}$ (Bukowski and Janowiec. 1996). The 2-acetylimidazo[4, [5-b] pyridine with INH, this hydrazide hydrazones 55, exhibited anti-TB activity against Mtb H37 Rv, Mtb 192, Mtb 210, isolated from patients and resistant against INH, ethambutol, RIF (Bukowski et al., 1999). Various 2, 3, 4-pentanetrione-3-(4-(((5-nitro-2-furyl)pyridyl/substituted-phenyl)-methylene) hydrazine) carbonyl


phenyl) hydrazones were exhibited anti-TB activity against *M. fortuitum* ATCC 6841 and Mtb H37Rv. Some compounds were active against *M. fortuitum*. Compound, which exhibited >90% inhibition at 12.5 $\mu\text{g/mL}$ against Mtb H37Rv, was the most promising compound for anti-TB activity. The actual MIC and IC_{50} values of were 3.13 and 0.32 $\mu\text{g/mL}$. The compound 55 was also tested against *M. avium* (Kucukguzel et al., 2000). The pyridylmethyleneamino derivative, compound 56 was showed activity against Gram negative, Gram positive Mtb H37Rv strains and some isonicotinoylhydrazones also showed a moderate activity against clinically isolated Mtb (6.25-50 $\mu\text{g/mL}$) which was INH resistant (Cocco et al., 2005). The N2-Substituted

alkylidene/arylidene-6-phenylimidazo [2], [1-b] thiazole-3-acetic acid hydrazides 57 were showed in vitro anti-TB activity against Mtb H37Rv (Ulusoy et al., 2000). (5-(Pyridine-2-yl)-1, 3, 4-thiadiazole-2-yl) thio) acetic acid arylidene-hydrazide derivatives 58 were exhibited in-vitro anti-TB activity, some compounds showed activity at 20µg/mL against Mtb and at 40µg/mL against *M. avium* (Mamolo et al., 2001). N-Alkylidene/arylidene-5-(2-furyl)-4-ethyl-1,2,4-triazole-3-mercaptoacetic acid hydrazides 59 were showed in-vitro anti-TB activity against Mtb H37 Rv at 6.25µg/mL (Ulusoy et al., 2001). The 4-quinolyldiazones, compound 38 was active against Mtb H37Rv. The 4-quinolyldiazine and aryl- or heteroaryl carboxaldehydes, most of the derivatives had anti-TB properties (Savini et al., 2002). Benzoic acid ((5-nitro-


thiophene-2-yl) methylene) hydrazide series 60 were tested against Mtb H37Rv. 4-Methoxybenzoic acid ((5-nitrothiophene-2-yl) methylene) hydrazide [61] was the most active, with a MIC value of 2.0 µg/mL (Rando et al., 2002). Ethyl 2-((3, 5-dimethylpyrazole-4-yl) hydrazono)-3-oxobutyrates 62 and methyl 2-((3, 5-dimethylpyrazole-4-yl) hydrazono)-4-methoxy-3-oxobutyrates 63 showed 29 and 28% inhibition against Mtb H37Rv (Kaymakçioğlu and Rollas, 2002). Compound 64 was exhibited anti-TB activity against Mtb H37Rv and *M. avium*. Compound 64 was found potent with the MIC value of 6.25µg/mL against Mtb H37Rv (Kuşukgüzel and Rollas, 2002).




54




55




56




57




58




59




60




61




62




63




64




65



66










67



The (5-(Pyridine-2-yl)-1, 3, 4-thiazole-2-yl) acetic acid (3, 4-diaryl-3H-thiazole-2-ylidene) hydrazide derivatives 65 were exhibited in vitro anti-TB activity. Compound 65 was exhibited at 20 µg/mL against Mtb 190 (Mamolo et al., 2003). N'-{1-(2-hydroxy-3-(piperazine-1-yl-methyl) phenyl) ethylidene} isonicotinohydrazide 66 was found to be the most active compound with the MIC of 0.56 µM, and more potent than INH (Sriram et al., 2005). The INH-related isonicotinoyl hydrazones (ISNEs), 2'-monosubstituted isonicotino hydrazides and cyanoboranes 67 were exhibited in vitro anti-TB activity. Some compounds showed excellent (MICs 0.025 to 0.2 µg/mL) to moderate (6.25 to 12.5 µg/mL) MICs value against EMB and RIF resistant strains (Maccari et al., 2005). Fluoroquinolones 68 containing a hydrazones were exhibited in vivo against Mtb H37Rv (Shindikar and Viswanathan. 2005). N'-Arylidene-N-(2-oxo-2-(4-aryl-piperazin-1-yl)ethyl) hydrazide derivatives 69 containing INH hydrazide-hydrazones were showed anti-TB activity against Mtb H37Rv ATCC 27294 and Mtb clinical isolates (Sinha et al., 2005). Compound 70 containing INH hydrazidehydrazones, 1-(4-Fluorophenyl)-3-(4-{1-(pyridine-4-carbonyl) hydrazono} ethyl) phenyl) thiourea was found potent compound, with MIC of 0.49 µM against Mtb H37Rv and INH-resistant Mtb (Sriram et al., 2006). The compounds 71,72,73, N-(2-fluorophenyl)-N'-quinoline-2-yl-methylene hydrazine, N-(2-adamantan-1-yl)-N'-quinoline-4-ylmethylene)-N'-4-fluorophenyl) hydrazine and N-(2-

cyclohexyl)-N'-quinoline-4-yl-methylene)-(2-fluorophenyl) hydrazine exhibited 99% inhibition at the lowest tested conc. of 3.125 µg/mL against drug-sensitive Mtb H37 strain (Nayyar et al., 2006). Various diclofenac acid hydrazones 74 were exhibited anti-TB activities against Mtb in vitro at dose ranging from 0.0383 to 7.53 µM (Sriram et al., 2006). Hydrazide-hydrazones with 4-substituted benzoic acid 75 were showed anti-TB activity. 4-Fluorobenzoic ((5-nitrothiophene-2-yl) methylene) hydrazide 75 showed the inhibition (99%) at conc. (6.25 µg/mL) (Bijev. 2006). New hydrazones, compound 76, 77 containing a pyrrole ring was showed potential TB statics and some compounds showed 92-100% inhibition of Mtb H37Rv at 6.25 µg/mL (Kucukguzel et al., 2006). The N-(3-({2-((2E)-2-Benzylidenehydrazino)-2-oxoethyl)sulfanyl)-5-({2((acetyl)amino)-1,3-thiazol-4-yl)methyl)-4H-1,2,4-triazol-4-yl)-3-nitrobenzamide 78 and N-(3-({2-((2E)-2-benzylidene hydrazine)-2-oxoethyl)sulfanyl)-5-({2((benzoyl) amino)-1,3-thiazol-4-yl)methyl)-4H-1,2,4-triazol-4-yl)-3-nitrobenzamide 79 have most active, with MIC values ranging from 0.39 to 0.78 µM (Shiradkar et al., 2007). A series of hydrazones, compound 80 was synthesized from INH, PZA, p-aminosalicylic acid (PAS), EMB and ciprofloxacin. 2-Hydroxy-4-(((isonicotinoyl hydrazono) methyl) amino) benzoic acid 57d showed the highest inhibition (96%) of Mtb H37Rv at 0.39 µg/mL (Imramovsky et al., 2007).



5. Antiprotozoal activity

The inhibitory activity of hydrazones [86], [87] showed against cruzipena major cysteine protease of *T. cruzi* (Caputto et al., 2011). The in-vitro antiamebic activity of hydrazones [88], [89] against the HM1: IMSS strain of *Entamoeba histolytica*. The compounds are reported to have IC₅₀ value of 0.03 and 0.04 μM

respectively (Hayat et al., 2010). Hydrazone derivatives [90], [91] described to be of high utility in Chagas disease (Vaio et al., 2009). Some hydrazone derivatives [92], [93] were showed antiamebic activity (Siddiqui et al., 2012), some hydrazone derivatives [94], [95] act as cruzin inhibitors (Romeiro et al., 2009) and some hydrazone derivatives showed antitrypanosomal activity [96] (Aponte et al., 2010).





Figure 2 Chemical structure of hydrazone derivatives no.44-75


6. Trypanocidal activity

A non-peptidyl acyl hydrazide acted as proteinase inhibitor for inhibition of brucipain (major cystein proteinase). The compound showed IC₅₀ values <40 μM against *Trypanosoma brucei* cell culture. Some acyl hydrazides showed 50% or more inhibition of trypanosome replication at <1 μM. The trypanocidal activity of most effective compound 4-Nitro-benzoic acid (4-(4-nitrophenyl)-cyclopenta-1, 3-dienylmethylene)-hydrazide [97] was comparable to drugs sumarin and diminazeneaceturate (Caffrey et al., 2002). A series of thiosemicarbazone and amino acyl thiazolidones derivatives, 2-[N'-(2-(4-Chloro-phenylsulfanyl)-ethylidene)-hydrazino]-5-methyl-thiazol-4-one [98] exhibited significant in vitro activity against epimastigot *Trypanosoma cruzi* (Leite et al., 2006).





7. Leishmanicidal activity:

Two trans (Pt (Hpy1)₂(Cl)₂) and trans (Pt(Hpy2)₂ (Cl)₂) complexes by the reaction of K₂PtCl₄ with sterol hydrazone ligands 20-hydrazone-pyridin-2-yl-5a-pregnan-3b-ol (Hpy1) and 22-hydrazone-pyridin-2-yl-chol-5-ene-3b-ol (Hpy2). These compounds are specific inhibitors of sterol methyl transferase (SMT). Promastigotes of *Leishmania (L) Mexicana* were treated for 48 h with 10 μM of the sterol hydrazones Hpy1 or Hpy2 alone or coordinated to Pt. Hpy1 produced higher leishmanistatic activity than Hpy2 (39% growth inhibition vs. 16%), which significantly increased (71%) when the complex trans- (Pt (Hpy1)₂(Cl)₂) was used (Visbal et al., 2008).


















8. Antitumoral activity

Substituted pyrazole-5-carbohydrazide hydrazone derivatives were showed potent apoptosis inducer in A549 lung cancer cells. The (E)-1-(4-tert-butylbenzyl)-N'-(1-(5-chloro-2-hydroxyphenyl) ethylidene)-3 (4-chlorophenyl) -1H-pyrazole-5-carbohydrazide [99] showed the highest growth inhibitory effect and induced apoptosis of A549 cells (Zheng et al., 2009). Several benzo (d) isothiazolehydrazones showed potential antiretroviral activity. Among the compounds benzo (d) isothiazole-3-carboxylic acid (4-methoxy-benzylidene) -hydrazide [100] was found to be the most potent antiproliferative compound (Vicini et al., 2006).



Some of diphenolic hydrazones showed uterotrophic inhibition of 70%, whereas compound 101 exhibited cytotoxicity in the range of 50-70% against MCF-7 and ZR-75-1 human malignant breast cell lines (Pandey et al., 2002). N'-(1-{1-(4-nitrophenyl-3-phenyl-1H-pyrazole-4-yl} methylene)-2-chlorobenzohydrazide 102 was found to be the most active (Abadi et al., 2003). Some 2, 6-dimethyl-N'-substituted-phenylmethyleneimidazo [2], [1-b] [1], [3], [4] thiadiazole-5-carbohydrazides, (2, 6-Dimethyl-N'-(2-hydroxy phenyl methylidene) imidazo[2], [1-b] [1], [3], [4] thiadiazole-5-carbohydrazide 103 showed the cytotoxicity. In the in vitro screening of 60 human tumors cell lines, this compound showed marked effects on the ovarian cancer cell line (Terzioğlu and Gürsoy. 2003). 3-(((6-Chloro-3-phenyl-4(3H)-quinazolinone-2-yl) mercapto acetyl) hydrazono)-5-fluoro-1H-2 indolinone 104 showed the favourable cytotoxicity against the renal cancer cell line UO-31 (Gürsoy and Karali. 2003). Some active compound was the 3- and 5-methylthiophene-2-carboxaldehyde α -(N)-heterocyclichydrazones derivatives 105, which exhibited tumor growth inhibition activity against all cell lines at GI50 values between 1.63 and 26.5 μ M (Savini et al., 2004).






5-Chloro-3-methylindole-2-carboxylic acid (4-nitrobenzylidene) hydrazide 106 arrests T47D cells in G2/Mphase of the cell cycle and to induce apoptosis. A 20-fold increase of apoptotic activity was achieved from the 5-methyl-3-phenylindole-2-carboxylic acid (4-methylbenzylidene) hydrazide 107 and 5-chloro-3-phenylindole-2-carboxylic acid (4-nitrobenzylidene) hydrazide 108, with EC_{50} values of 0.1 μ M in the caspase activation assay in T47D breast cancer cells. Compound 108 was found to be highly active in a standard growth inhibition assay with a GI_{50} value of 0.9 μ M in T47D cells. Compound 106 and its analogs were found to inhibit tubulin polymerization (Zhang et al., 2004). Hydrazide-hydrazones containing 5-oxo-[1], [2], [4] triazole ring. Some compounds had inhibitory effect on mycelial growth whereas compounds 109 and 110 were found antitumor activity in breast cancer (Demirbas et al., 2004). Hydrazino pyrimidine derivatives 111 were showed in vitro antitumoral activity in nine different types of human cancers. Some compounds demonstrated inhibitory effects on the growth of a wide range of cancer cell lines (Cocco et al., 2005). Several benzo (d) isothiazole hydrazones

have been tested for antitumoral activity. Compound 112, bearing a hydroxy group at o-position of the benzylidene moiety, was the most potent, thus acting equally potent as 6-mercaptopurine against the haematological tumors (Vicini et al., 2006). N'-Substituted-benzylidene-3, 4, 5-trimethoxy benzohydrazide 113 were showed antitumoral activity against some cancer cells. Many hydrazone compounds showed good antitumor activity against PC3 cells and showed moderate activities against Bcap37 and BGC823 cells (Jin et al., 2006). 6-Amino-4-aryl-2-oxo-1-(1-pyrid-3-yl- or 4-yl-ethylidene-amino)-1, 2-dihydropyridine-3, 5-dicarbonyl nitrile series 114 exhibited a high percentage of tumor growth inhibition all cancer cell lines (Gürsoy and Güzeldemirci-Ulusoy, 2007). N'-(3, 5-Di-tert-butyl-4-hydroxybenzylidene)-6-nitro-1, 3-benzo dioxole-5-carbohydrazine as a novel anti-proliferative compound. They observed that was able to inhibit T-cell proliferation (66 % at 10 μ M) (Duarte et al., 2007). A series of arylidenehydrazides were evaluated against the full panel of 60 human tumour cell lines. Compound 115 demonstrated the most effect on prostate cancer cell line (El-Hawash et al., 2006).


Some derivatives act as PI3K p110 α inhibitors. PI3K are signaling proteins in different cell types responsible for phosphorylation of lipids in cell membranes (Kendall et al., 2012). Various bis (indolyl) based hydrazones active against multiple cancer cell lines (Kumar et al., 2012). Hydrazone derivative was showed potent activity against HL-60 leukaemia and 518A2 melanoma (Effenberger et al., 2010). Acylhydrazones have potent activity against the human promyelocytic leukemic cells (HL-60) [Cui et al., 2010]. Copper based hydrazone derivatives are act against integrin β_4 in H322 lung carcinoma cell lines (Fan et al., 2010). Palladium based hydrazones have active against human head and neck squamous carcinoma cell lines SQ20B and SCC-25 (Abu-Surrah et al., 2010). Various hydrazones derivatives, 2-phenylindole based hydrazone exhibited against breast carcinoma cell lines and reported to have an IC_{50} of 1.60nM (El-Nakkady et al., 2012; Al-Said et al., 2011; Hassan et al., 2011).

9. Analgesic, anti-inflammatory and antiplatelet activity


A series of N-Arylhydrazone derivatives mefenamic acid were showed analgesic and anti-inflammatory activity. The pyridine rings at the aryl moiety of the arylhydrazones have good analgesic activity in comparison to mefenamic acid. Compounds possessing the 4-tolyl or 4-fluorophenyl moiety are more active than 4-bromophenyl and 4-N, N-dimethylaminophenyl. The anti-inflammatory some potent compounds showed that replacement of carboxylic acid group of mefenamic acid with N-arylhydrazone moiety cannot produce any advantage in the anti-inflammatory property (Ali et al., 2005). Analgesic activity of some (4Z)-3-methyl-1-((2-oxo-2H-chromene-4-yl) carbonyl)-1H-pyrazole-4, 5-dione 4-((4-substitutedphenyl) hydrazine) some of them showed significant analgesic activity. The presence of 4-chloro, 4-bromo, 3, 4-dichloro, 3, 4-dibromo and 4-methyl group in the aromatic ring of 4-position of the pyrazole-hydrazono nucleus gave rise to an increased analgesic activities (Sivakumar et al., 2010). Hydrazone derivatives of quinoxalinone were evaluated for anti-inflammatory activity which showed that compounds having methoxy group at the para position showed comparatively good percentage of inhibition of edema (Khan et al., 2009). The 3-(1H-Indol-3-yl)-3-oxo-2-((5-phenyl-2H-pyrazol-3-yl)-hydrazono)-propionitrile [116] was found to possess appreciable analgesic and anti-inflammatory activity (Radhwan et al., 2007). The antiplatelet activity of tricyclic acylhydrazone derivatives was showed ability to inhibit platelet aggregation. Benzylidene-/4'-bromobenzylidene 3-OH-8-CH₃-6- phenylpyrazolo [3], [4-b] thieno-[2], [3-d] pyridine-2-carbohydrazide were exhibited inhibition of the PAF-induced platelet aggregation (Todeschini et al., 1998). The benzylidene 10H-phenothiazine-1-carbohydrazides were acts in the AA pathway probably by inhibition of platelet COX-1 enzyme (Fraga et al., 2000).




114




115




116




117




118




119




120



121








122






123


N-acylarylhydrazone class, ((4'-N, N-Dimethylaminobenzylidene-3-(3', 4'methylene dioxy phenyl) propionylhydrazine) 117 was more potent than dipyrone and indomethacine (Lima et al., 2000). The tricyclic acylhydrazone derivatives 118 were showed antiplatelet activity. Benzylidene-4'-bromobenzylidene 3-hydroxy-8-methyl-6-phenyl pyrazolo [3], [4-b] thieno-[2], [3-d] pyridine-2-carbohydrazone were evaluated as PAF-induced platelet aggregation inhibitor (Fraga et al., 2000). The benzylidene 10H-phenothiazine-1-carbohydrazone ($IC_{50}=2.3\mu M$) 119, inhibit COX-1 enzyme. The change in para-substituent group of acylhydrazone permitted to identify a hydrophilic carboxylate derivative and a hydrophobic bromo derivative as analgesic that are more potent than dipyrone, possessing selective peripheral or central mechanism of action (Silva et al., 2004). Hydrazones containing 5-methyl-2-benzoxazoline, the analgesic effects of 2-(2-(5-methyl-2-benzoxazoline-3-yl) acetyl)-4-chloro-4-methyl benzylidene hydrazine 120 and 121 were found to be higher than those of morphine and aspirin. In addition, 2-(2-(5-methyl-2-benzoxazoline-3-yl) acetyl)-4- methoxy benzylidene hydrazine 122 at 200 mg/kg dose possessed the most antiinflammatory activity (Salgın-Gökşen et al., 2007). N'-(3, 5-Di-tert-butyl-4-hydroxybenzylidene)-6-nitro-1, 3-benzodi oxole-5-carbohydrazone 123 as an antiinflammatory compound (Duarte et al., 2007). Compound furoxanyl-N-acylhydrazones [124], [125] were showed Analgesic and anti-inflammatory activity (Hernandez et al., 2012). The anti-inflammatory activity exhibited by some aryl hydrazones [126] (Rajitha et al. 2011). Various hydrazone derivatives [127] have promising in-vivo anti-inflammatory activity (Moldovan et al., 2011). Hydrazone derivatives [128], [129] with selective COX-2 inhibition with ED_{50} value of 0.2 mmol/Kg (El-Sayed et al., 2011). Benzo-thiophene derivatives [130] have inhibition of 50.2% inflammation (Isloor et al., 2010).




10. Central nervous system

Hydrazones [131] have inhibited PDE10A-a phosphodiesterase responsible for neurological and psychological disorder such as Parkinson's, Schizophrenia and Huntington's disease (Gage et al., 2011). The anticonvulsant potential of some hydrazone derivatives [91], [92] having long duration of action and a rapid onset of action (Kaushik et al., 2010). Antidepressant activity of hydrazones [93] has been reported (de Oliveira et al., 2011). The in-vitro β -amyloid aggregation inhibitory activity showed by hydrazone derivatives [96] with an IC_{50} of $23\mu M$ (Catto et al., 2010).










Antiplatelet activity: Antiplatelets prevent the formation of thrombus. Hydrazone derivatives [136], [137], and [138] exhibited in-vitro antiplatelet activity (Jordao et al., 2011).



11. Anticonvulsant Activity


The acetylhydrazones 139 provided good protection against convulsions while the oxamoylhydrazones 140 were significantly less active (Dimmock et al., 2000). Some hydrazones of (2-oxobenzoxazoline-3-yl) acetohydrazide 141 were tested for their antiepileptic activity against scPTZ test. The 4-fluoro derivative was found to be more active than the other compounds (Çakır et al., 2001). The GABA hydrazones 142 were showed anticonvulsant properties (Ragavendran et al., 2007). The 2-Chloroquinolinyl Hydrazone [143] derivatives have anticonvulsant activity (Kumar et al., 2010).






The N'-((5-chloro-3-methyl-1-phenyl-1H-pyrazol-4-yl) methylene) 2/4-substituted hydrazides showed anticonvulsant activity, aryl derivative exhibited activity against MES and scPTZ models. When the para to the aryl ring was substituted with electron withdrawing group the compounds were found to be more potent and active in both models as compared to electron donating group. The 3, 4-di (substituted) oxy-N2, N5-bis (substituted) thiophene-2, 5-dicarbohydrazide exhibited anticonvulsant activity against MES, and scPTZ induced seizure (Darpan et al., 2010). Some hydrazide-hydrazone derivatives showed antidepressant activity (Mohareb et al., 2010). A series of 2-(1H-indol-3-yl) acetyl-N-(substituted phenyl) hydrazine carbothioamides and their related heterocyclic derivatives exhibited anticonvulsant activity against MES method. The hydrazone-indole based heterocyclic derivatives were found to have encouraging anticonvulsant activity (Siddiqui et al., 2012). Anticonvulsants from a combined phthalimide-GABA-anilide and hydrazone pharmacophore, all of the compounds were ineffective in the MES test. Most of the compounds were found to be effective in the scSTY and ipPIC models and very few compounds showed protection in the sc PTZ model. The pharmacophoric combinations of phthalimide-GABA-anilide/hydrazones are as anticonvulsants. The phthalimide GABA-anilides were found to be more effective than the corresponding phthalimide-GABA-hydrazone derivatives (Darpan et al., 2010; Ragavendran et al., 2007).

12. Antioxidant activity


Hydrazone derivatives [144], [145] were exhibited free radical scavenging activity (Musad et al., 2011). Imidazoline based hydrazones [146], [147] reported as promising antioxidant activity (Abdel-Wahab et al., 2011).








13. Vasodilator activity

The octahydroquinazolinehydrazones [148], [149], and [150] were exhibited potential hypotensive agents (El-Sabbagh et al., 2010). A compound of the N-acylhydrazone, 3, 4-methylene dioxylbenzoyl-2-thienyl hydrazone, (LASSBio-294), showed inotropic and vasodilatory effects. Derivatives of LASSBio-294 were exhibited contractile responses of rat vascular smooth muscle. Phenylephrine-induced contractions of aorta was inhibited by the derivatives N-methyl-2-thienylidene-3, 4-methylenedioxy-benzoyl hydrazine (LASSBio-785) and N-allyl-2-thienylidene-3, 4-methylenedioxy-benzoyl hydrazine (LASSBio-788). LASSBio-785 was seven times more potent than the LASSBio-294 in endothelium-independent vasodilator effect (Silva et al., 2005).



14. Antiparasitic activity

The anti-parasitic activity of hydrazone derivatives [151], [152] showed against *Ctenocephalides felis* and *Rhipicephalus sanguineus* (Ali et al., 2010). Hydrazone derivatives [153], [154] as a urease inhibitors, urease catalyzes the hydrolysis of urea to ammonia and carbamate and it is beneficial for the pathogenesis of urolithiasis, pyelonephritis, ammonia and hepatic encephalopathy, hepatic coma and urinary catheter encrustation (Aslam et al., 2011).



15. Antidiabetic activity

Pyrimidylhydrazones as inhibitors of glycogen synthase kinase-3 (GSK-3) with the most active compound exhibited antidiabetic activity (Smalley et al., 2006).

16. Discussion

Hydrazones constitutes an important class of compounds for various biological activities. These observations promote us for the development of new hydrazones that possess varied biological activities and has been considerable interest in the development of novel compounds with diverse biological activities (Govindasami et al., 2011; Rajitha et al., 2011; Uppal et al., 2011; Belskaya et al., 2010; Scior and Garcés-Eisele, 2006; Nayyar and Jain, 2005). Vast works have been done on hydrazones across the world for the development of better biological molecules as well as with low toxicity profiles. Different hydrazones have been developed and found to be active pharmacologically.

17. Conclusion

The use of hydrazones as lead for development of newer agents and hydrazone moiety possesses an array of activity. With proper synthesis of potential compounds can be designed for different biological activity.

References

- [1] Abadi AH, Eissa AAH, Hassan GS. Synthesis of novel 1, 3, 4-trisubstituted pyrazole derivatives and their evaluation as antitumor and antiangiogenic agents. *Chem. Pharm. Bull.*, 2003, 51, 838-844.
- [2] Abdel-Aal MT, El-Sayed WA, El-Ashry EH. Synthesis and antitumor evaluation of some sugar arylglycinoylhydrazones and their oxadiazoline derivatives. *Arch. Pharm. Chem., Life Sci.* 2006, 339, 656-663.
- [3] Abdel-Wahab FB, Awad AEG, Badria AF. Synthesis, antimicrobial, antioxidant, anti-hemolytic and cytotoxic evaluation of new imidazole-based heterocycles. *Eur. J. Med. Chem.*, 2011, 46, 1505-1511.
- [4] Abdel-Wahab FB, Khidre ER, Awad AEG. Regioselective synthesis and antimicrobial activities of some novel aryloxyacetic acid derivatives. *Eur. J. Med. Chem.*, 2012, 50, 55-62.
- [5] Abu-Surrah AS, Safieh KAA, Ahmad IM, Abdalla MY, Ayoub MT, Qaroush AK, Abu-Mahtheieh AM. New palladium (II) complexes bearing pyrazole-based Schiff base ligands: Synthesis, characterization and cytotoxicity. *Eur. J. Med. Chem.*, 2010, 45, 471-475.
- [6] Ajani OO, Obafemi, C.A, Nwinyi, O.C, Akinpelu, D.A. Microwave assisted synthesis and antimicrobial activity of 2-quinoxalinone-3-hydrazone derivatives. *Bioorg. Med. Chem.*, 2010, 18, 214-221.
- [7] Ali, A, Fisara, P, Freemont, J.A, Kyi, S, Meyer, A.G, Andrew, G, Riches, A.G, Sargent, R.M, Sawutz, D.G, Turner, K.A, Winzenberg, KN, Yang, Q. Discovery of ectoparasiticide hydrazono-trifluoromethanesulfonamides, *Bioorg. Med. Chem. Lett.*, 2010, 20, 649-652.
- [8] Ali, A, Mohammad, T, Davood, Abbas, S. Synthesis and analgesic activity of N-aryl hydrazone derivatives of mefenamic acid. *J Pharm Pharmaceut Sci.*, 2005, 8(3):419-25.
- [9] Ali, M.R, Marella, A, Alam, M.T, Naz, R, Akhter, M, Shaquiquzzaman, M, Saha, R, Tanwar, O, Alam, M.M, Hooda, J. Review of Biological Activities of Hydrazones. *Indonesian J. Pharm.*, 2012, 23(4): 193-202.
- [10] Al-Macrosaur, L.Q, Dayam, R, Taheri, L, Witvrouw, M, Debyser, Z, Neamati, N. Discovery of novel non-cytotoxic salicylhydrazine containing HIV-1 integrase Inhibitors. *Bioorg. Med. Chem. Lett.* 2007, 17: 6472-75.
- [11] Al-Said, M.S, Bashandy, M.S, Al-qasoumi, S.I, Ghorab, M.M. Anti-breast cancer activity of some novel 1,2-dihydropyridine, thiophene and thiazole derivatives. *Eur. J. Med. Chem.*, 2011, 46, 137-141.
- [12] Angelusiua, M.V, Barbuceanu, S-F, Draghicic, C, Almajan, G.L. New Cu(II), Co(II), Ni(II) complexes with aroyl- hydrazone based ligand. Synthesis, spectroscopic characterization and in vitro antibacterial evaluation. *Eur. J. Med. Chem.*, 2010; 45:2055-62.
- [13] Aponte, J.C, Vaisberg, A.J, Castillo, D, Gonzalez, G, Estevez, Y, Arevalo, J, Quiliano, M, Zimic, M, Verastegui, M, Malaga, E, Juan, R.H.G, Bustamante, B, Tarleton, R.L, Wang, Y, Franzblau, S.G, Pauli, G.F, Sauvain, M, Hammond, G.B. Trypanoside, antituberculosis, leishmanicidal, and cytotoxic activities of tetrahydrobenzothienopyrimidines, *Bioorg. Med. Chem.*, 2010, 18, 2880-2886.
- [14] Aslam, M.A.S, Mahmood, S, Shahid, M, Saeed, A, Iqbal, J. Synthesis, biological assay in vitro and molecular docking studies of new Schiff base derivatives as potential urease inhibitors. *Eur. J. Med. Chem.*, 2011, 46, 5473-5479.
- [15] Belskaya, N.P, Dehaen, W, Bakulev, V.A. Synthesis and properties of hydrazones bearing amide, thioamide and amidine functions. *ARKIVOC*, 2010, 1, 275-332.
- [16] Bernardino, A, Gomes, A, Charret, K, Freitas, A, Machado, G, Canto-Cavalheiro, M, Leon, L, Amaral, V. Synthesis and leishmanicidal activities of 1-(4-X-phenyl)-N'-((4-Yphenyl) methylene)-1H-pyrazole-4-carbohydrazides *Eur. J. Med. Chem.* 2006, 41, 80-87.
- [17] Bijev A. New heterocyclic hydrazones in the search for antitubercular agents: Synthesis and in vitro evaluations. *Lett. Drug Des. Discov.* 2006, 3, 506-512.
- [18] Caffrey, C.R, Schanz, M, Nkemgu-Njinkeng, J, Brush, M, Hausell, E, Cohen, F.E, Flaherty, T.M, Mckerrrow, J.H, Steverding, D. Screening of acyl hydrazide proteinase inhibitors for antiparasitic activity against *Trypanosoma brucei*. *Int. J. Antimicrob. Agents.* 2002, 19, 227-251.
- [19] Çakır, B, Dağ, Ö, Yıldırım, E, Erol, K, Şahin, M.F. Synthesis and anticonvulsant activity of some hydrazones of 2-((3H)-oxobenzoxazolin-3-yl-aceto)hydrazide. *J. Fac. Pharm. Gazi.* 2001, 18, 99-106.
- [20] Caputto, M.E, Fabian, L.E, Benitez, D, Merlinoz, A, Rios, N, Cerecetto, H, Moltrasio, G.Y, Moglioni, A.G, Gonzalez, M, Finkielstein, L.M. Thiosemicarbazones derived from 1-indanones as new anti-*Trypanosoma cruzi* agents, *Bioorg. Med. Chem.*, 2011, 19, 6818-6826.
- [21] Catto, M, Aliano, R, Carotti, A, Cellamare, S, Palluotto, F, Purgatorio, R, Stradis, A. D, Campagna, F, Design, synthesis and biological evaluation of indane-2-arylhydrazinylmethylene-1,3-diones and indol arylidiazinylmethylene-3-ones as β -amyloid aggregation inhibitors, *Eur. J. Med. Chem.*, 2010, 45, 1359-136.
- [22] Cocco, M. T, Congiu, C, Lilliu, V, Onnis, V. Synthesis and in vitro antitumoral activity of new hydrazinopyrimidine-5-carbonitrile derivatives. *Bioorg. Med. Chem.* 2005, 14, 366-372.
- [23] Cui, Z, Li, Y, Ling, Y, Huang, J, Cui, J, Wang, R, Yang, X. New class of potent antitumor acylhydrazone derivatives containing furan, *Eur. J. Med. Chem.*, 2010, 45, 5576-5584.
- [24] Darpan, K, Khan, S.A, Gita, P, C, Verma, P, Kumar, S. N'-(5-chloro-3-methyl-1-phenyl-1H-pyrazol-4-yl) methylene) 2/4-substitutedhydrazides: Synthesis and anticonvulsant activity. *Eur. J. Med. Chem.*, 2010, 45, 3943-49.
- [25] de Oliveira, K.N, Costa, P, Santin, J. R, Mazzambani, L, Burger, C, Mora, C, Nunes, R. J, de-Souza, M. M. Synthesis and antidepressant-like activity evaluation of sulphonamides and sulphonyl-hydrazones, *Bioorg. Med. Chem.*, 2011, 19, 4295-4306.
- [26] Deep, A, Jain, S, Sharma, P, Kumar, M, Dora, C.P. Design and biological evaluation of biphenyl-4- carboxylic acid hydrazone-hydrazone for antimicrobial activity. *Acta Pol Pharm.*, 2010, 67(3), 255-9.
- [27] Demirbas, N, Karaoglu, S, Demirbas, A, Sancak, K. Synthesis and antimicrobial activities of some new 1-(5-phenylamino-(1,3,4)thiadiazol-2-yl)methyl-5-oxo-(1,2,4)triazole and 1-(4-phenyl-5-thioxo-(1,2,4) triazol-3-yl) methyl-5-oxo-(1,2,4)triazole derivatives. *Eur. J. Med. Chem.* 2004, 39, 793-804.
- [28] Dimmock, J.R, Vashishtha, S.C, Stables, J.P. Anticonvulsant properties of various acetylhydrazones, oxamoylhydrazones and semicarbazones derived from aromatic and unsaturated carbonyl compounds. *Eur. J. Med. Chem.* 2000, 35, 241-248.
- [29] Duarte, C.D, Tributino, J.L.M, Lacerda, D.I, Martins, M.V, Alexandre-Moreira, M.S, Dutra, F, Bechara, E.J.H, De-Paula, F.S, Goulart, M.O.F, Ferreira, J, Calixto, J.B, Nunes, M.P, Bertho, A.L, Miranda, A.L.P, Barreiro, E.J, Fraga, C.A.M. Synthesis, pharmacological evaluation and electrochemical studies of novel 6-nitro-3,4-methylenedioxyphenyl-Nacylhydrazone derivatives : Discovery of LASSBio-881, a new ligand of cannabinoid receptors. *Bioorg. Med. Chem.*, 2007, 15, 2421-2433.
- [30] Duarte, C.D, Tributino, J.L.M, Lacerda, D.I, Martins, M.V, Alexandre-Moreira, M.S, Dutra, F, Bechara, E.J.H, De-Paula, F.S, Goulart, M.O.F, Ferreira, J, Calixto, J.B, Nunes, M.P, Bertho, A.L, Miranda, A.L.P, Barreiro, E.J, Fraga, C.A.M. Synthesis, pharmacological evaluation and electrochemical studies of novel 6-nitro-3,4-methylenedioxyphenyl-Nacylhydrazone derivatives : Discovery of LASSBio-881, a new ligand of cannabinoid receptors. *Bioorg. Med. Chem.*, 2007, 15, 2421-2433.
- [31] Edrees, M.M, Farghaly, T.A, El-Hag, A.A.F, Abdalla, M.M. Antimicrobial, antitumor and 5 α -reductase inhibitor activities of some hydrazoneoyl substituted pyrimidinones, *Eur. J. Med. Chem.*, 2010, 45, 5702-5707.
- [32] Effenberger, K, Breyer, S, Schobert R. Modulation of doxorubicin activity in cancer cells by conjugation with fatty acyl and terpenyl hydrazones, *Eur. J. Med. Chem.*, 2010, 45, 1947-1954.
- [33] El-Hawash, S.A.M, Abdel Wahab, A.E, El-Dewellawy, M.A. Cyanoacetic acid hydrazones of 3-(and 4-) acetylpyridine and some derived ring systems as potential antitumor and anti-HCV agents. *Arch. Pharm. Chem. Life Sci.* 2006, 339, 14-23.
- [34] El-Nakkady, S.S, Hanna, M.M, Roaiah, H. M, Ghannam, I.A.Y. Synthesis, molecular docking study and antitumor activity of novel 2-phenylindole derivatives, *Eur. J. Med. Chem.*, 2012, 47, 387-398.
- [35] El-Sabbagh, O. I, Shabaan, M.A, Kadry, H. H, Al-Din, E.S. New octahydroquinazoline derivatives: Synthesis and hypotensive activity, *Eur. J. Med. Chem.*, 2010, 45, 5390-5396.
- [36] El-Sayed, M.A.A, Abdel-Aziz, N.I, Abdel-Aziz, N.A.M, El-Azab, A.S, Asiri, Y.A, Tahir, K.E.H.E. Design, synthesis, and biological evaluation of substituted hydrazone and pyrazole derivatives as selective COX-2 inhibitors, molecular docking study, *Bioorg. Med. Chem.*, 2011, 19, 3416-3424.
- [37] Eswaran, S, Adhikari, A.V, Chowdhury, I.H, Pal, N.K, Thomas, K.D. New quinoline derivatives: Synthesis and investigation of antibacterial and antituberculosis properties, *Eur. J. Med. Chem.*, 2010, 45, 3374-3383.
- [38] Fan, C.D, Su, H, Zhao, J, Zhao, B.X, Zhang, S.L, Miao, J.Y. A novel copper complex of salicylaldehyde pyrazole hydrazone induces apoptosis through up-regulating integrin β -4 in H322 lung carcinoma cells, *Eur. J. Med. Chem.*, 2010, 45, 1438-1446.
- [39] Fraga, A.G.M, Rodrigues, C.R, Miranda, A.L.P, Barreiro, E.J, Fraga, C.A.M. Synthesis and pharmacological evaluation of novel heterocy-

- clic acylhydrazone derivatives, designed as PAF antagonists. *Eur. J. Pharm. Sci.* 2000, 11, 285-290.
- [40] Gage, J.L., Onrust, R., Johnston, D., Osnowski, A., MacDonald, W., Mitchell, L., Urogdi, L., Rohde, A., Harbol, K., Gragerov, S., Dorman, G., Wheeler, T., Florio, V., Cutshall, N.S. N-Acylhydrazones as inhibitors of PDE10A, *Bioorg. Med. Chem. Lett.*, 2011, 21, 4155-4159.
- [41] Gemma, S., Kukreja, G., Fattorusso, C., Persico, M., Romano, M., Altarelli, M., Savini, L., Campiani, G., Fattorusso, E., Basilio, N. Synthesis of N1-arylidene-N2-quinolyl- and N2- acrydinyldhydrazones as potent antimalarial agents active against CQ-resistant *P. falciparum* strains. *Bioorg. Med. Chem. Lett.*, 2006, 16, 5384-5388.
- [42] Gouda, M.A., Berghot, M.A., Shoeib, A.I., Khalil, M.A. Synthesis and antimicrobial of new anthraquinone derivatives incorporating pyrazole moiety, *Eur. J. Med. Chem.*, 2010, 45, 1843-1848.
- [43] Govindasami, T., Pandey, A., Palanivelu, N., Pandey, A. Synthesis, characterization and antibacterial activity of biologically important vanillin related hydrazone derivatives, *Int. J. Org. Chem.*, 2011, 1, 71-77.
- [44] Gülerman, N.N., Oruc, E.E., Kartal, F., Rollas, S. In vivo metabolism of 4-fluorobenzoic acid ((5-nitro-2-furanyl) methylene) hydrazide in rats. *Eur. J. Drug Metab. Pharmacokinet.*, 2000, 25, 103-108.
- [45] Guniz, K.S., Rollasa, S., Erdeniz, H., Kiraz, M., Cevdet, E.A., Vidinc, A. Some 4-arylhydrazono-2-pyrazoline-5-one derivatives obtained from heterocyclic amines, tested against *Mycobacterium avium*. *Eur. J. Med. Chem.*, 2007, 35, 761-771.
- [46] Gürsoy, A., Karali, N. Synthesis and primary cytotoxicity evaluation of 3-(((3-phenyl-4(3H) - quinazolinone-2-yl)mercaptoacetyl)hydrazono)-1H-2-indolinones. *Eur. J. Med. Chem.* 2003, 38, 633-643.
- [47] Gürsoy, E., Güzeldemirci-Ulusoy, N. Synthesis and primary cytotoxicity evaluation of new imidazo(2,1-b)thiazole derivatives. *Eur. J. Med. Chem.*, 2007, 42, 320-326.
- [48] Hassan, G.S., Kadry, H.H., Abou-Seri, S.M., Ali, M.M., Mahmoud, A.E.E. Synthesis and in vitro cytotoxic activity of novel pyrazolo(3,4-d)pyrimidines and related pyrazole hydrazones toward breast adenocarcinoma MCF-7 cell line, *Bioorg. Med. Chem.*, 2011, 19, 6808-6817.
- [49] Hayat, F., Salahuddin, A., Zargan, J., Azam, A. Synthesis, characterization, antiamebic activity and cytotoxicity of novel 2-(quinolin-8-yloxy) acetohydrazones and their cyclized products (1,2,3-thiadiazole and 1,2,3-selenadiazole derivatives), *Eur. J. Med. Chem.*, 2010, 45, 6127- 6134.
- [50] Hernandez, P., Cabrera, M., Lavaggi, M.L., Celano, L., Tiscornia, I., da Costa, T.R., Thomson, L., Bollati-Fogolin, M., Miranda, A.L.P., Lima, L.M., Barreiro, E.J., Gonzalez, M., Cerecetto, H. Discovery of new orally effective analgesic and anti-inflammatory hybrid furoxanyl N-acylhydrazone derivatives, *Bioorg. Med. Chem.*, 2012, 20, 2158-2171.
- [51] Imramovsky, A., Polanc, S., Vinsova, J., Kocivar, M., Jampilek, J., Recková, Z., Kaustova, J. A new modification of anti-tubercular active molecules. *Bioorg. Med. Chem.* 2007, 15, 2551-2559.
- [52] Isloor, A.M., Kalluraya, B., Pai, K.S. Synthesis, characterization and biological activities of some new benzo(b) thiophene derivatives, *Eur. J. Med. Chem.*, 2010, 45, 825-830.
- [53] Jin, C.A.Y., Tan, Z., He, M., Tian, B., Tang, S., Hewlett, I., Yang, M. SAR and molecular mechanism study of novel acylhydrazone compounds targeting HIV-1, *Bioorg. Med. Chem.*, 2010, 18, 2135-2140.
- [54] Jin, L., Chen, J., Song, B., Chen, Z., Yang, S.; Li, Q., Hu, D., Xu R. Synthesis, structure, and bioactivity of N'-substitutedbenzylidene-3,4,5-trimethoxybenzohydrazide and 3-acetyl-2- substituted phenyl-5-(3,4,5-trimethoxyphenyl)-2,3-dihydro-1,3,4-oxadiazole derivatives. *Bioorg. Med. Chem. Lett.* 2006, 16, 5036-5040.
- [55] Jordao, A.K., Sathler, P.C., Ferreira, V.F., Campos, V.R., de Souza, M.C.B.V., Castro, H.C., Lannes, A., Lourenco, A., Rodrigues, C.R., Bel-lo, M.L., Lourenco, M.C.S., Carvalho, G.S.L., Almeida, M.C.B., Cunha, A.C. Synthesis, antitubercular activity, and SAR study of N-substitutedphenylamino-5-methyl-1H-1,2,3-triazole-4-carbohydrazides, *Bioorg. Med. Chem.*, 2011, 19, 5605-5611.
- [56] Jubie, S., Meena, S., Ramashesu, K. V., Jawahar, N., Vijaykumar, S. Synthesis and biological evaluation of some hydrazones and carbazones of indane 1,3-dione. *Ind. J. Chem.*, 2010, 49B, 1261-1263.
- [57] Kaushik, D., Khan, S.A., Chawla, G., Kumar, S. N'-(5-chloro-3-methyl-1- phenyl-1H-pyrazol-4-yl) methylene] 2/4-substituted hydrazides: Synthesis and anticonvulsant activity, *Eur. J. Med. Chem.*, 2010, 45, 3943-3949.
- [58] Kaymakcioglu, K. B., Oruc, E.E., Unsalan, S., Kandemirli, F., Shvets, N., Rollas, S., Anatholy, D. Synthesis and characterization of novel hydrazide-hydrazones and the study of their structureantituberculosis activity. *Eur. J. Med. Chem.*, 2006, 41, 1253-1261.
- [59] Kaymakcioglu, K.B., Rollas, S. Synthesis, characterization and evaluation of antituberculosis activity of some hydrazones. *Farmaco*, 2002, 57, 595-599.
- [60] Kendall, J.D., Giddens, A.C., Tsang, K.Y., Frederick, R., Marshall, E.S., Singh, R., Lill, C.L., Lee, W.J., Kolekar, S., Chao, M., Malik, A., Yu, S., Chaussade, C., Buchanan, C., Rewcastle, G.W., Baguley, B.C., Flanagan, J.U., Jamieson, S.M.F., Denny, W.A., Shepherd, P.R., Novel pyrazolo(1,5-a) pyridines as p110 α -selective PI3 kinase inhibitors: Exploring the benzenesulfonylhydrazide SAR. *Bioorg. Med. Chem.*, 2012, 20, 58-68.
- [61] Khan, S.A., Pooja, M., Shabir, P., Darpan, K. Hydrazone derivatives of quinoxalinone as antimicrobial and antiinflammatory agents. *Acta Pol Pharm.*, 2009, 66(2), 169-72.
- [62] Kucukguzel, S.G., Kocatepe A., De Clercq, E., Sahin F., Gulluce, M. Synthesis and biological activity of 4-thiazolidinones, thiosemicarbazides derived from diflunisal hydrazide. *Eur. J. Med. Chem.*, 2006, 41, 353-359.
- [63] Kuçukguzel, S.G., Kuçukguzel, I., Ulgen M. Metabolic and Chemical Studies on N-(4- chlorobenzyl)-N'-benzoylhydrazine, *Farmaco*, 2000, 55, 624-630.
- [64] Kucukguzel, S.G., Mazi, A., Sahin, F., Öztürk S., Stables, J. P. Synthesis and biological activities of diflunisal hydrazide-hydrazones. *Eur. J. Med. Chem.* 2003, 38, 1005-1009.
- [65] Kuçukguzel, S.G., Oruc E.E., Rollas S., Şahin, F., Ozbek, A. Synthesis, Characterization and biological activity of novel 4-thiazolidinones, 1,3,4-oxadiazoles and some related compounds. *Eur. J. Med. Chem.*, 2002, 37, 197-206.
- [66] Kuçukguzel, S.G., Rollas, S. Synthesis, Characterization of Novel Coupling Products and 4- Arylhydrazono-2-pyrazoline-5-ones as Potential Antimycobacterial Agents. *Farmaco*, 2002, 57, 583-587.
- [67] Kumar, D., Judge, V., Narang, R., Sangwan, S., Clercq, D.E., Balzarini, J., Narasimhan, B. Benzylidene/2-chlorobenzylidene hydrazides: Synthesis, antimicrobial activity, QSAR studies and antiviral evaluation, *Eur. J. Med. Chem.*, 2010, 45, 2806-2816.
- [68] Kumar, D., Kumar, N. M., Ghosh, S., Shah, K. Novel bis(indolyl)hydrazide-hydrazones as potent cytotoxic agents, *Bioorg. Med. Chem. Lett.*, 2012, 22, 212-215.
- [69] Kumar, S., Bawai, S., Drabu, S., Kumar, R., Machwal, L. Chloroquinolinylhydrazone derivative as anticonvulsant. *Acta Pol Pharm.*, 2010, 67, 567-3.
- [70] Lee, Y.J., Jeong, W.K., Shin, S., Lee, U.J., Kim, Y. Discovery of novel selective inhibitors of *Staphylococcus aureus* β -ketoacyl acyl carrier protein synthase III. *Eur. J. Med. Chem.*, 2012, 47, 261-269.
- [71] Leite, A.C.L., deLima, R.S., Moreira, D.R., Cardoso, M.V., de Brito, A.C.G., dos Santos, L.M.F., Hernandes, M.Z., Kipustok, A.C., de Lima, R.S., Soares, M.B.P. Synthesis, docking, and *in vitro* activity of thiosemiconbazone, aminoacylthiosemicarbazides and acylthiazolidones against *Trypanosomacruzi*. *Bioorg. Med. Chem.*, 2006, 14, 3749-57.
- [72] Lima, P.C., Lima, L. M., Silva, K.C., Leda, P.H., Miranda, A.L.P., Fraga, C.A.M.; Barreiro, E.J., Synthesis and analgesic activity of novel N-acylhydrazones and isosters, derived from natural saffrole. *Eur. J. Med. Chem.*, 2000, 35, 187-203.
- [73] Liu, W.-Y., Li, H.-Y., Zhao, B.-X., Shin, D.-S., Lian, S., Miaonovel, J.-Y. Ribavirin hydrazone derivatives and anti-proliferative activity against A549 lung cancer cells. *Carbohydrate Res.*, 2009, 344, 1270-5.
- [74] Loncle, C., Brunel, J., Vidal, N., Dherbomez, M., Letourneux, Y. Synthesis and antifungal activity of cholesterol-hydrazone derivatives. *Eur. J. Med. Chem.*, 2004, 39, 1067-1071.
- [75] Maccari, R., Ottana, R., Vigorita, M.G. In vitro advanced antimycobacterial screening of isoniazid-related hydrazones, hydrazides and cyanoboranes: Part 14. *Bioorg. Med. Chem. Lett.*, 2005, 15, 2509-2513.
- [76] Mague, G.M., Jakhlal, J., Ladyman, M., Vallin, A., Ralambomanana, D.A., Bousquet, T., Maugein, J., Lebibi, J., Pelinski, L. Synthesis and antimycobacterial activity of a series of ferrocenyl derivatives. *Eur. J. Med. Chem.*, 2011, 46, 31-38.
- [77] Mahajan, A., Kremer, L., Louw, S., Gueradel, Y., Chibale, K., Biot, C. Synthesis and in vitro antitubercular activity of ferrocene-based hydrazones. *Bioorg. Med. Chem. Lett.*, 2011, 21, 2866-2868.
- [78] Mamolo, M.G., Falagiani, V., Zampieri, D., Vio, L., Banfi, E., Scialino, G. Synthesis and antimycobacterial activity of (3,4-diaryl-3H-thiazole-2-ylidene)hydrazide derivatives. *Farmaco*, 2003, 58, 631-637.
- [79] Mamolo, M.G., Falagiani, V., Zampieri, D., Vio, L., Banfi, E. Synthesis and antimycobacterial activity of (5-(pyridin-2-yl)-1, 3, 4-thiadiazole-2-ylthio) acetic acid arylidene-hydrazide derivatives. *Farmaco*, 2001, 56, 587-592.
- [80] Manvar, A., Malde, A., Verma, J., Virsodia, V., Mishra, A., Upadhyay, K., Acharya, H., Countinhe, E., Shah, A. Synthesis, antitubercular activ-

- ity and 3D-QSAR study of coumarin-4-acetic acid benzylidene hydrazides. *Eur. J. Med. Chem.*, 2008, 3, 342-4.
- [81] Masunari, A, Tavares, L.C. A new class of nifuroxazide analogues: Synthesis of 5-nitrophenyl derivatives with antimicrobial activity against multidrug-resistant *Staphylococcus aureus*. *Bioorg. Med. Chem.*, 2007, 15, 4229-4236.
- [82] Metwally KA, Abdel-Aziz LM, Lashine EM, Husseiny MI, Badawy RH. Hydrazones of 2-aryl-quinoline-4-carboxylic acid hydrazides: synthesis and preliminary evaluation as antimicrobial agents. *Bioorg. Med. Chem.*, 2006, 14, 8675-82.
- [83] Mohareb RM, El-Sharkawy KA, Hussein MM, El-Sehrawi HM. Hydrazidehydrazones derivatives as antidepressant, sedative and analgesic agents. *J. Pharm. Sci. & Res.*, 2010, 2, 185-196.
- [84] Moldovan CM, Oniga O, Parvu A, Tiperciuc B, Verite P, Pirnau A, Bojit OCM, Pop R. Synthesis and antiinflammatory evaluation of some new acyl-hydrazones bearing 2-aryl-thiazole. *Eur. J. Med. Chem.*, 2011, 46, 526-534.
- [85] Musad EA, Mohamed R, Saeed BA, Vishwanath BS, Rai KML. Synthesis and evaluation of antioxidant and antibacterial activities of new substituted bis (1,3,4-oxadiazoles), 3,5-bis(substituted) pyrazoles and isoxazoles. *Bioorg. Med. Chem. Lett.*, 2011, 21, 3536-3540.
- [86] Nayyar A, Jain R. Recent advances in new structural classes of anti-tuberculosis agents. *Curr. Med. Chem.*, 2005, 12, 1873-1886.
- [87] Nayyar A, Malde A, Coutinho E, Jain R. Synthesis, anti-tuberculosis activity, and 3D-QSAR study of ring-substituted-2/4-quinolinecarbaldehyde derivatives. *Bioorg. Med. Chem.*, 2006, 14, 7302-7310.
- [88] Nayyar A, Monga V, Malde A, Coutinho E, Jain R. Synthesis, anti-tuberculosis activity and 3D-QSAR study of 4-(adamantan-1-yl)-2-substituted quinolines. *Bioorg. Med. Chem.* 2007, 15, 626-40.
- [89] Negi VJ, Sharma AK, Negi JS, Ram V. Biological activities of hydrazone derivatives in the new millennium. *Int. J. Pharm. Chem.*, 2012, 2 (4), 100-109.
- [90] Olayinka O, Craig A, Obafemi B, Obinna C, Nwinyi C, David A. Akinpelu. Microwave assisted synthesis and antimicrobial activity of 2-quinoxalinone-3-hydrazone derivatives. *Bioorg & Med Chem.*, 2010, 18, 214-21.
- [91] Ozdemir A, Turan-Zitouni G, Kaplanciklia AZ, Iscan G, Khan S, Demirci F. Synthesis and the selective antifungal activity of 5,6,7,8-tetrahydroimidazo (1,2-a)pyridine derivatives. *Eur. J. Med. Chem.*, 2010, 45, 2080-2084.
- [92] Ozdemir OU, Gulcin O. Sufonylhydrazone derivatives and their nickel complexes as a antibacterial agents. *Spectrochimica Acta.*, 2008, 70, 641-5.
- [93] Ozemir A, Kalancikli ZA, Zitouni GT, Revial G. Synthesis of some novel hydrazone derivatives and evaluation of their antituberculosis activity. *Marmara Pharm J.*, 2010, 14, 79-83.
- [94] Ozkay Y, Tunali Y, Karaca H, Isikdag I. Antimicrobial activity and a SAR study of some novel benzimidazole derivatives bearing hydrazone moiety. *Eur. J. Med. Chem.*, 2010, 45, 3293-3298.
- [95] Pandey J, Pal R, Dwivedi A, Hajela K. Synthesis of some new diaryl and triaryl hydrazone derivatives as possible estrogen receptor modulators. *Arzneimittelforschung.* 2002, 52, 39-44.
- [96] Pavan FR, Maia PIS, Leite SRA, Deflon VM, Batista AA, Sato DN, Franzblau SG, Leite CQF. Thiosemicarbazones, semicarbazones, dithiocarbazates and hydrazide/ hydrazones: Anti-*Mycobacterium tuberculosis* activity and cytotoxicity. *Eur. J. Med. Chem.*, 2010, 45, 1898-1905.
- [97] Radhwan MAA, Ragab EA, Sabry NM, El-Shenawy SM. Synthesis and biological evaluation of new 3-substituted indole derivatives as potential antiinflammatory and analgesic agents. *Bioorg. Med. Chem.*, 2007, 15: 3832-41.
- [98] Ragavendran J, Sriram D, Patel S, Reddy I, Bharathwajan N, Stables, J, Yogeeswari P. Design and synthesis of anticonvulsants from a combined phthalimide-GABA-anilide and hydrazone pharmacophore. *Eur. J. Med. Chem.*, 2007, 42, 146-151.
- [99] Raja AS, Agarwal AK, Mahajan N, Pandeya SN, Ananthan A. Antibacterial and antitubercular activity of some diphenyl hydrazone and semicarbazone. *Ind. J. Chem.*, 2010, 49B, 1384-1388.
- [100] Rajitha G, Saideepa N, Praneetha P. Synthesis and evaluation of N-(α - benzamidocinnamoyl)aryl hydrazones Activities of Hydrazones derivative for anti-inflammatory and antioxidant activities, *Ind. J. Chem.*, 2011, 50B, 729-733.
- [101] Rando D, Sato, D.N, Siqueira, L, Malvezzi, A, Leite, C.Q.F, Amaral, A.T, Ferreira, E.I, Tavares, L.C. Potential tuberculostatic agents. Topliss application on benzoic acid ((5-Nitrothiophene-2-yl) methylene) hydrazide series. *Bioorg. Med. Chem.*, 2002, 10, 557-560.
- [102] Rane RA, Telvekar VN. Synthesis and evaluation of novel chloropyrrole molecules designed by molecular hybridization of common pharmacophores as potential antimicrobial agents. *Bioorg. Med. Chem. Lett.*, 2010, 20, 5681-5685.
- [103] Rasras MJA, Al-Tel HT, Al-Aboudi FA, Al-Qawasmeh AR. Synthesis and antimicrobial activity of cholic acid hydrazone analogues. *Eur. J. Med. Chem.*, 2010, 45, 2307-2313.
- [104] Rollas S, Gulerman N, Erdeniz H. Synthesis and antimicrobial activity of some new hydrazones of 4-fluorobenzoic acid hydrazide and 3-acetyl-2,5-disubstituted-1,3,4-oxadiazolines. *Farmaco.* 2002, 57, 171-174.
- [105] Rollas S, Kucukguzel SG. Biological Activities of Hydrazone Derivatives. *Molecules.* 2007, 12, 1910-1939.
- [106] Romeiro NC, Aguirre G, Hernandez P, González M, Cerecetto H, Aldana I, Silanes SP, Monge A, Barreiro EJ, Lima LM. Synthesis, trypanocidal activity and docking studies of novel quinoxaline-N-acylhydrazones, designed as cruzain inhibitors candidates, *Bioorg. Med. Chem.*, 2009, 17, 641-652.
- [107] Salgın-Gökşen, U, Gökhan-Kelekci, N, Goktas, O, Koyas, Y, Kılıç, E, Isık, S, Aktay, G, Özalp, M. 1-Acylthiosemicarbazides, 1,2,4-triazole-5(4H)-thiones, 1,3,4-thiadiazoles and hydrazones containing 5-methyl-2-benzoxazolones: Synthesis, analgesic-anti-inflammatory and antimicrobial activities. *Bioorg. Med. Chem.*, 2007, 15, 5738-5751.
- [108] Savini L, Chiasserini, L, Gaeta, A, Pellerano, C. Synthesis and Anti-tubercular Evaluation of Quinolyl hydrazones. *Bioorg. Med. Chem.*, 2002, 10, 2193-2198.
- [109] Savini L, Chiasserini, L, Travagli, V, Pellerano, C, Novellino, E, Cosentino, S, Pisano, M.B. New α -heterocyclic hydrazones: evaluation of anticancer, anti-HIV and antimicrobial activity. *Eur. J. Med. Chem.*, 2004, 39, 113-122.
- [110] Scior T, Garcés-Eisele SJ. Isoniazid is not a lead compound for its pyridyl ring derivatives, isonicotinoyl amides, hydrazides and hydrazones: A critical review. *Curr. Med. Chem.*, 2006, 13, 2205-2219.
- [111] Sharma RN, Sharma KP, Dikshit SN. Synthesis, Characterization and biological activities of some new hypophosphorousadducts of acid hydrazones derived from 2-((N-benzoyl) 2,3-dichloroanilido) acetohydrazide. *Arch Appl Sci Res.*, 2011, 3, 415-424.
- [112] Shindikar AV, Viswanathan CL. Novel fluoroquinolones: design, synthesis, and in vivo activity in mice against *Mycobacterium tuberculosis* H37 Rv. *Bioorg. Med. Chem. Lett.*, 2005, 15, 1803-1806.
- [113] Shiradkar MR, Murahari KK, Gangadasu HR, Suresh T, Kalyan CC, Panchal D, Kaur R, Burange P, Ghogare J, Mokale V, Raut M. Synthesis of new S-derivatives of clubbed triazolyl thiazole as anti-*Mycobacterium tuberculosis* agents. *Bioorg. Med. Chem.*, 2007, 15, 3997-4008.
- [114] Siddiqui SM, Salahuddin A, Azam A. Synthesis, characterization and antiamoebic activity of some hydrazone and azole derivatives bearing pyridyl moiety as a promising heterocyclic scaffold. *Eur. J. Med. Chem.*, 2012, 49, 411-416.
- [115] Silva AG, Zapata-Suto G, Kummerle AE, Fraga CAM, Barreiro EJ, Sudo RT. Synthesis and vasodilatory activity of new N-acylhydrazone derivatives, designed as LASS Bio-alogues. *Bioorg. Med. Chem.*, 2005, 13, 3431-37.
- [116] Silva GA, Costa LMM, Brito FCF, Miranda ALP, Barreiro EJ, Fraga CAM. New class of potent antinociceptive and antiplatelet 10H-phenothiazine-1-acylhydrazone derivatives. *Bioorg. Med. Chem.* 2004, 12, 3149-3158.
- [117] Sinha N Jain S, Tilekar A, Upadhyaya RS, Kishore N, Jana GH, Arora SK. Synthesis of isonicotinic acid N'-Arylidene-N-(2-oxo-2-(4-aryl-piperazin-1-yl)ethyl)hydrazides as antituberculosis agents. *Bioorg. Med. Chem. Lett.*, 2005, 15, 1573-1576.
- [118] Sivakumar KK, Rajasekaran A, Ponnilaravan I, Somasundaram A, Sivasakthi R, Kamalaveni S. Antimicrobial and analgesic activity of some (4Z)-3-methyl-1-((2-oxo-2H-chromen-4-yl)carbonyl)-1H-pyrrole-4,5-dione 4-((4-substitutedphenyl) hydrazone). *Scholars Res Lib.*, 2010, 2(1), 211-19.
- [119] Smalley TL Jr, Peat AJ, Boucheron JA, Dickerson S, Garrido D, Preugschat F, Schweiker SL, Thomson SA, Wang TY. Synthesis and evaluation of novel heterocyclic Inhibitors of GSK-3. *Bioorg. Med. Chem. Lett.*, 2006, 16, 2091-94.
- [120] Sriram D, Yogeeswari P, Devakaram RV. Synthesis, in vitro and in vivo antimycobacterial activities of diclofenac acid hydrazones and amides. *Bioorg. Med. Chem.*, 2006, 14, 3113-3118.
- [121] Sriram D, Yogeeswari P, Madhu K. Synthesis and in vitro and in vivo antimycobacterial activity of isonicotinoyl hydrazones. *Bioorg. Med. Chem. Lett.*, 2005, 15, 4502-4505.
- [122] Sriram D, Yogeeswari P, Madhu K. Synthesis and in vitro antitubercular activity of some 1-(4-sub phenyl)-3-(4-{1-((pyridine-4-carbonyl)hydrazono)ethyl} phenyl)thiourea. *Bioorg. Med. Chem.*, 2006, 14, 876-878.

- [123] Srivastava R, Pandeya SN, Singh AK, Sam A. Synthesis of some new acid hydrazone and their activity against mycobacterium. *Inter J Chem Res*, 2010, 2(1), 18-19.
- [124] Terzioğlu N, Gursoy A. Synthesis and anticancer evaluation of some new hydrazone derivatives of 2,6-dimethylimidazo(2,1-b)-(1,3,4)thiadiazole-5-carbohydrazide. *Eur. J. Med. Chem.*, 2003, 38, 781-786.
- [125] Thomas KD, Adhikari AV, Telkar S, Chowdhury IH, Mahmood R, Pal KN, Rowd G, Sumesh E. Design, synthesis and docking studies of new quinoline-3-carbohydrazide derivatives as antitubercular agents. *Eur. J. Med. Chem.*, 2011, 46, 5283-5292.
- [126] Todeschini AR, Miranda AL, Silva CM, Parrini SC, Barreiro EJ. Synthesis and evaluation of analgesic, anti-inflammatory and antiplatelet properties of new 2- pyridylarylhydrazone derivatives. *Eur. J. Med. Chem.*, 1998, 33, 189-99.
- [127] Turan-Zitouni G, Blache Y, Guven K. Synthesis and antimicrobial activity of some imidazo-(1,2-a) pyridine-2-carboxylic acid arylidenehydrazide derivatives. *Boll. Chim. Farm.* 2001, 140, 397-400.
- [128] Ulusoy N, Çapan G, Ötük G, Kiraz M. Synthesis and antimicrobial activity of new 6- phenylimidazo(2,1-b)thiazole derivatives. *Boll. Chim. Farm*, 2000, 139, 167-172.
- [129] Ulusoy N, Gürsoy A, Otuk G, Kiraz M. Synthesis and antimicrobial activity of some 1,2,4-triazole-3-mercaptoacetic acid derivatives. *Farmaco*, 2001, 56, 947-952.
- [130] Uppal G, Bala S, Kamboj S, Saini M. Therapeutic Review Exploring Antimicrobial Potential of Hydrazones as Promising Lead. *Der Pharma Chemica*, 2011, 3, 250-268.
- [131] Vaio MAFVD, Freitas ACC, Castro HC, Albuquerque S.D, Cabral LM, Rodrigues CR, Albuquerque MG, Martins RCA, Henriques MGMO, Dias LRS. Synthesis, antichagasic in vitro evaluation, cytotoxicity assays, molecular modeling and SAR/QSAR studies of a 2-phenyl-3-(1-phenyl-1Hpyrazol-4-yl)-acrylic acid benzylidenecarbohydrazide series, *Bioorg. Med. Chem.*, 2009, 17, 295-302.
- [132] Vavrikova E, Polanc S, Kocovar M, Horvati K, Bosze S, Stolarikova J, Vavrova K, Vinsova J. New fluorinecontaining hydrazones active against MDR-tuberculosis, *Eur. J. Med. Chem.*, 2011, 46, 4937-4945.
- [133] Vicini P, Incerti M, Doytchinova I, La Colla P, Busonera B, Loddò R. Synthesis and antiproliferative activity of benzo(d)isothiazole hydrazones. *Eur. J. Med. Chem.*, 2006, 41, 624-632.
- [134] Vicini P, Zani F, Cozzini P, Doytchinova I. Hydrazones of 1,2-benzisothiazole hydrazides: synthesis, antimicrobial activity and QSAR investigations. *Eur. J. Med. Chem.*, 2002, 37, 553-564.
- [135] Vijesh AM, Isloor MA, Prabhu V, Ahmad S, Malladi S. Synthesis, characterization and anti-microbial studies of some novel 2,4-disubstituted thiazoles, *Eur. J. Med. Chem.*, 2010, 45, 5460-5464.
- [136] Visbal G, Marchan E, Maldonado, A, Simoni, Z, Navarro, M. Synthesis and characterization of platinum-sterol hydrazone complexes with biological activity against *Leishmania (L.) mexicana*. *J. Inorg. Biochem.*, 2008, 102, 547-54.
- [137] Walcourt A, Loyevsky M, Lovejoy DB, Gordeuk, V. R, Richardson, D.R. Novel aroylhydrazone and thiosemicarbazone iron chelators with anti-malarial activity against chloroquine-resistant and -sensitive parasites. *Int. J. Biochem. Cell Biol.*, 2004, 36, 401-407.
- [138] Zareef M, Iqbal R, Mirza B, Khan KM, Manan A, Asim F, Khan W. Synthesis and antimicrobial activity of some derivatives of acylhydrazine including novel benzenidiazasulfonamides. *Eur. J. Med. Chem.*, 2008, 141-52.
- [139] Zhang H, Drewe J, Tseng B, Kasibhatla S, Cai SX. Discovery and SAR of indole-2- carboxylic acid benzylidenehydrazides as a new series of potent apoptosis inducers using a cellbased HTS assay. *Bioorg. Med. Chem.*, 2004, 12, 3649-3655.
- [140] Zheng L-W, Wu L-L, Zhao B-X, Dong W-L, Miao J-Y. Synthesis of novel substituted pyrazole-5-carbohydrazide hydrazone derivatives and discovery of a potent apoptosis inducer in A549 lung cancer cells. *Bioorg. Med. Chem.*, 2009, 17, 1957-1962.