



General solution and generalized Ulam-Hyers stability of a generalized n - type additive quadratic functional equation in Banach space and Banach algebra: direct and fixed point methods

S. Murthy^{1*}, M. Arunkumar², V. Govindan³

¹ Department of Mathematics, Government Arts College (For Men), Krishnagiri -635 001, Tamilnadu, India

² Department of Mathematics, Government Arts College, Thiruvannamalai -606 603, Tamilnadu, India

³ Department of Mathematics, Sri Vidya Mandir Arts & Science College, Uthangarai -636 902, Tamilnadu, India

*Corresponding author E-mail: smurthy07@yahoo.co.in

Copyright © 2014 S. Murthy et al. This is an open access article distributed under the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

In this paper, the authors introduce and investigate the general solution and generalized Ulam-Hyers stability of a generalized n -type additive-quadratic functional equation.

$$\begin{aligned} & f(nx + n^2y + n^3z) + f(nx - n^2y + n^3z) + f(nx + n^2y - n^3z) + f(-nx + n^2y + n^3z) \\ &= n[f(x) - f(-x)] + n^2[f(y) - f(-y)] + n^3[f(z) - f(-z)] \\ &+ 2n^2[f(x) + f(-x)] + 2n^4[f(y) + f(-y)] + 2n^6[f(z) + f(-z)], \end{aligned}$$

Where n is a positive integer with $n \neq 0$, in Banach Space and Banach Algebras using direct and fixed point methods.

Keywords: Additive Functional Equations; Quadratic Functional Equations; Mixed Type Functional Equations; Banach Space; Banach Algebra; Generalized Ulam-Hyers Stability; Fixed Point.

1. Introduction and preliminaries

The stability of functional equation was raised by S.M. Ulam [52] for what metric group G is it true that a ε -automorphism of G is necessary near to a strict automorphism?

In 1941, D.H. Hyers [27] gave a positive answer to the question of Ulam for Banach spaces. In 1950, T. Aoki [2] was second author to treat this problem for additive mappings. Th.M. Rassias [48] succeeded in extending the result of Hyers Theorem by weakening the condition for the Cauchy difference controlled by $(\|x\|^p + \|y\|^p)$, $p \in [0, 1)$ to be unbounded. Considering a lot of influence of Ulam, Hyers and Rassias on the development of stability problems of functional equations, the stability phenomenon that was proved by Th.M. Rassias is called Hyers-Ulam-Rassias stability (see [19], [21], [28], [32], [36], and [50]).

The stability problems of several functional equations have been extensively investigated by a number of authors [16], [20], [24], [47], and [51]. The terminology generalized Ulam-Hyers stability originates from these historical backgrounds.

Over the last seven decades, the above problem was talked by numerous authors and its solutions via various forms of functional equations like, additive, quadratic, cubic, quartic mixed type functional equations were discussed. We refer the interested readers for more information on such problems to the monographs of [4], [6], [9], [11], and [13].

C. Park [41], [43], [44] applied Gavruta's result to Banach Modules over a C^* -Algebra. Many authors have studied the structure of C^* -Algebra for different types of functionals in various settings one can refer [42]. It seems that an approximate derivation was first investigated by K.W. Jun and D.W. Park [45]. Recently, the stability of derivations has been investigated in [42], [43], [52] and references therein. The stability of cubic derivations was first time introduced and investigated by M.E. Gordji et al, [25]. With the help of [43] the stability of quadratic derivations was discussed by M. Arunkumar ET. al., [3].

One of the most famous functional equations is the additive functional equation

$$f(x+y) = f(x) + f(y). \quad (1)$$

In 1821, it was first solved by A.L. Cauchy in the class of continuous real-valued functions. It is often called Cauchy additive functional equation in honor of A.L. Cauchy. The theory of additive functional equations is frequently applied to the development of theories of other functional equations. Moreover, the properties of additive functional equations are powerful tools in almost every field of natural and social sciences. Every solution of the additive functional equation (1) is called an additive function.

The quadratic function $f(x) = cx^2$ satisfies the functional equation

$$f(x+y) + f(x-y) = 2f(x) + 2f(y) \quad (2)$$

And therefore, the equation (2) is called quadratic functional equation. The Hyers - Ulam stability theorem for the quadratic functional equation (2) was proved by F.Skof for the functions $f: E_1 \rightarrow E_2$ where E_1 a normed space is and E_2 be a Banach space. The result of Skof is still true if the relevant domain E_1 is replaced by an Abelian group, and it was delt by P.W.Cholewa [19]. S.Czerwik [20], [21] proved the Hyers-Ulam-Rassias stability of the quadratic functional equation (2). This result was further generalized by Th.M.Rassais, C.Borelli and G.L.Forti [16].

The solution and stability of the following mixed type additive-quadratic functional equations

$$f(2x+y) + f(2x-y) = 2f(x+y) + 2f(x-y) + 2f(2x) - 4f(x) \quad (3)$$

$$f(x+2y) + f(x-2y) + 8f(y) = 2f(x) + 4f(2y) \quad (4)$$

$$f\left(\sum_{i=1}^n x_i\right) + (n-2) \sum_{i=1}^n f(x_i) = \sum_{1 \leq i < j \leq n} f(x_i + x_j) \quad (5)$$

$$f(x+ay) + af(x-y) = f(x-ay) + af(x+y) \quad (6)$$

$$f(x+y) + f(x-y) = 2f(x) + f(y) + f(-y) \quad (7)$$

$$f(2x \pm y \pm z) = 2f(-x \mp y \mp z) - 2f(\mp y \mp z) + f(\pm y \pm z) + 3f(x) - f(-x) \quad (8)$$

$$\begin{aligned} f(x-t) + f(y-t) + f(z-t) &= 3f\left(\frac{x+y+z}{3} - t\right) + 3f\left(\frac{2x-y-z}{3}\right) \\ &+ 3f\left(\frac{-x+2y-z}{3}\right) + 3f\left(\frac{-x-y+2z}{3}\right) \end{aligned} \quad (9)$$

$$\begin{aligned} &f(x+2y+3z) + f(x-2y+3z) + f(x+2y-3z) + f(x-2y-3z) \\ &= 4f(x) + 8[f(y) + f(-y)] + 18[f(z) + f(-z)] \end{aligned} \quad (10)$$

Were discussed by A.Najati and M.B.Moghim [40], G.Zamani Eskandani [53], K.W.Jun, H.M.Kim [30], [31], M.Arunkumar, J.M.Rassias [47], M.J.Rassias et al, [5], S. Murthy et.al, [39] and M.Arunkumar, P.Agilan [10].

Motivated by the above findings in this paper, we introduce and investigate that the general solution and generalized Ulam-Hyers stability of a generalized n-type Additive Quadratic functional equation of the form

$$\begin{aligned}
& f(nx + n^2y + n^3z) + f(nx - n^2y + n^3z) + f(nx + n^2y - n^3z) + f(-nx + n^2y + n^3z) \\
&= n[f(x) - f(-x)] + n^2[f(y) - f(-y)] + n^3[f(z) - f(-z)] \\
&+ 2n^2[f(x) + f(-x)] + 2n^4[f(y) + f(-y)] + 2n^6[f(z) + f(-z)],
\end{aligned} \tag{11}$$

Where n is a positive integer with $n \neq 0$, in Banach spaces and Banach Algebras using direct and fixed point methods?

2. General solution of the functional equation (11): when f is odd

In this section, the general solution of the functional equation (11), for the odd case is discussed. Throughout this section, let us consider X and Y to be real vector spaces.

Theorem 2.1: If an odd mapping $f : X \rightarrow Y$ satisfies the functional equation

$$f(x + y) = f(x) + f(y) \tag{12}$$

For all $x, y \in X$, if and only if $f : X \rightarrow Y$ satisfies the functional equation

$$\begin{aligned}
& f(nx + n^2y + n^3z) + f(nx - n^2y + n^3z) + f(nx + n^2y - n^3z) + f(-nx + n^2y + n^3z) \\
&= n[f(x) - f(-x)] + n^2[f(y) - f(-y)] + n^3[f(z) - f(-z)] \\
&+ 2n^2[f(x) + f(-x)] + 2n^4[f(y) + f(-y)] + 2n^6[f(z) + f(-z)]
\end{aligned} \tag{13}$$

For all $x, y, z \in X$.

Proof. Let $f : X \rightarrow Y$ satisfies the functional equation (12). Setting (x, y) by $(0, 0)$ in (12), we get $f(0) = 0$. Replacing (x, y) by (x, x) and $(x, 2x)$ respectively in (12), we obtain

$$f(2x) = 2f(x) \text{ And } f(3x) = 3f(x) \tag{14}$$

For all $x \in X$. In general for any positive integer a , we have

$$f(ax) = af(x) \tag{15}$$

For all $x \in X$. It is easy to verify from (15) that

$$f(a^2x) = a^2f(x) \text{ And } f(a^3x) = a^3f(x) \tag{16}$$

For all $x \in X$. Replacing (x, y) by $(nx + n^2y, n^3z)$ in (12) and using (12), (15) and (16), we get

$$f(nx + n^2y + n^3z) = nf(x) + n^2f(y) + n^3f(z) \tag{17}$$

For all $x, y, z \in X$. Again replacing z by $-z$ in equation (17) and using oddness of f , we obtain

$$f(nx + n^2y - n^3z) = nf(x) + n^2f(y) - n^3f(z) \tag{18}$$

For all $x, y, z \in X$. Also replacing y by $-y$ in (17) and using oddness of f , we get

$$f(nx - n^2y + n^3z) = nf(x) - n^2f(y) + n^3f(z) \tag{19}$$

For all $x, y, z \in X$. Finally replacing x by $-x$ in (17) and using oddness of f , we obtain

$$f(-nx + n^2y + n^3z) = -nf(x) + n^2f(y) + n^3f(z) \quad (20)$$

For all $x, y, z \in X$. Adding the equations (17), (18), (19) and (20), we have

$$\begin{aligned} & f(nx + n^2y + n^3z) + f(nx - n^2y + n^3z) + f(nx + n^2y - n^3z) + f(-nx + n^2y + n^3z) \\ &= 2nf(x) + 2n^2f(y) + 2n^3f(z) \end{aligned} \quad (21)$$

For all $x, y, z \in X$. Using oddness of f in (21) and remodifying, we arrive

$$\begin{aligned} & f(nx + n^2y + n^3z) + f(nx - n^2y + n^3z) + f(nx + n^2y - n^3z) + f(-nx + n^2y + n^3z) \\ &= n[f(x) - f(-x)] + n^2[f(y) - f(-y)] + n^3[f(z) - f(-z)] \end{aligned} \quad (22)$$

For all $x, y, z \in X$. Adding $2n^2f(x) + 2n^4f(y) + 2n^6f(z)$ on both sides, remodifying and using oddness of f , we reach (13) as desired.

Conversely, $f : X \rightarrow Y$ satisfies the functional equation (13) using oddness of f in (13), we arrive

$$\begin{aligned} & f(nx + n^2y + n^3z) + f(nx - n^2y + n^3z) + f(nx + n^2y - n^3z) + f(-nx + n^2y + n^3z) \\ &= 2nf(x) + 2n^2f(y) + 2n^3f(z) \end{aligned} \quad (23)$$

For all $x, y, z \in X$. Replacing (x, y, z) by $(x, 0, 0)$, $(0, x, 0)$ and $(0, 0, x)$, respectively in (23), we obtain

$$f(nx) = nf(x), \quad f(n^2x) = n^2f(x) \quad \text{and} \quad f(n^3x) = n^3f(x) \quad (24)$$

For all $x \in X$. One can easily verify from (24) that

$$f\left(\frac{x}{n^i}\right) = \frac{1}{n^i}f(x); \quad i = 1, 2, 3 \quad (25)$$

For all $x \in X$. Replacing (x, y, z) by $\left(\frac{x}{n}, \frac{y}{n^2}, 0\right)$ in equation (23) and using oddness of f and (25), we arrive our result.

3. General solution of the functional equation (11): when f is even

In this section, the general solution of the functional equation (11) for even case is given. Throughout this section, let us consider X and Y to be real vector spaces.

Theorem 3.1: If an even mapping $f : X \rightarrow Y$ satisfies the functional equation

$$f(x + y) + f(x - y) = 2f(x) + 2f(y) \quad (26)$$

For all $x, y \in X$ if and only if $f : X \rightarrow Y$ satisfies the functional equation

$$\begin{aligned} & f(nx + n^2y + n^3z) + f(nx - n^2y + n^3z) + f(nx + n^2y - n^3z) + f(-nx + n^2y + n^3z) \\ &= n[f(x) - f(-x)] + n^2[f(y) - f(-y)] + n^3[f(z) - f(-z)] \\ &+ 2n^2[f(x) + f(-x)] + 2n^4[f(y) + f(-y)] + 2n^6[f(z) + f(-z)] \end{aligned} \quad (27)$$

For all $x, y, z \in X$.

Proof: Let $f : X \rightarrow Y$ satisfies the functional equation (26). Setting (x, y) by $(0, 0)$ in (26), we get $f(0) = 0$. Replacing y by x and y by $2x$ in (26), we obtain

$$f(2x) = 4f(x) \text{ And } f(3x) = 9f(x) \quad (28)$$

For all $x \in X$. In general for any positive integer b , such that

$$f(bx) = b^2 f(x) \quad (29)$$

For all $x \in X$. It is easy to verify from (29) that

$$f(b^2x) = b^4 f(x) \text{ And } f(b^3x) = b^6 f(x) \quad (30)$$

For all $x \in X$. Replacing (x, y) by (nx, n^2y) in (26) and using (26), we get

$$f(nx + n^2y) + f(nx - n^2y) = 2n^2 f(x) + 2n^4 f(y) \quad (31)$$

For all $x, y \in X$. Setting (x, y) by (nx, n^3z) in (26) and using (26), we obtain

$$f(nx + n^3z) + f(nx - n^3z) = 2n^2 f(x) + 2n^6 f(z) \quad (32)$$

For all $x, z \in X$. Replacing (x, y) by (n^2y, n^3z) in (26) and using (26), we have

$$f(n^2y + n^3z) + f(n^2y - n^3z) = 2n^4 f(y) + 2n^6 f(z) \quad (33)$$

For all $y, z \in X$. Adding equations (31), (32) and (33), we arrive

$$\begin{aligned} & f(nx + n^2y) + f(nx - n^2y) + f(nx + n^3z) + f(nx - n^3z) + f(n^2y + n^3z) + f(n^2y - n^3z) \\ &= 4n^2 f(x) + 4n^4 f(y) + 4n^6 f(z) \end{aligned} \quad (34)$$

For all $x, y, z \in X$. Replacing (x, y) by $(nx + n^2y, n^3z)$ in (26), we get

$$f(nx + n^2y + n^3z) + f(nx + n^2y - n^3z) = 2f(nx + n^2y) + 2f(n^3z) \quad (35)$$

For all $x, y, z \in X$. Setting (x, y) by $(n^3z, nx - n^2y)$ in (26), we obtain

$$f(nx - n^2y + n^3z) + f(-nx + n^2y - n^3z) = 2f(n^3z) + 2f(nx - n^2y) \quad (36)$$

For all $x, y, z \in X$. Replacing (x, y) by $(n^2y, nx + n^3z)$ in (26), we get

$$f(nx + n^2y + n^3z) + f(-nx + n^2y - n^3z) = 2f(n^2y) + 2f(nx + n^3z) \quad (37)$$

For all $x, y, z \in X$. Setting (x, y) by $(n^2y, nx - n^3z)$ in (26), we obtain

$$f(nx + n^2y - n^3z) + f(-nx + n^2y + n^3z) = 2f(n^2y) + 2f(nx - n^3z) \quad (38)$$

For all $x, y, z \in X$. Replacing (x, y) by $(nx, n^2y + n^3z)$ in (26), we get

$$f(nx + n^2y + n^3z) + f(nx - n^2y - n^3z) = 2f(nx) + 2f(n^2y + n^3z) \quad (39)$$

For all $x, y, z \in X$. Setting (x, y) by $(nx, n^2y - n^3z)$ in (26), we have

$$f(nx + n^2y - n^3z) + f(nx - n^2y + n^3z) = 2f(nx) + 2f(n^2y - n^3z) \quad (40)$$

For all $x, y, z \in X$. Now multiply by 2 on both sides of (34), we obtain

$$\begin{aligned} 2f(nx + n^2y) + 2f(nx - n^2y) + 2f(nx + n^3z) + 2f(nx - n^3z) + 2f(n^2y + n^3z) + 2f(n^2y - n^3z) \\ = 8n^2f(x) + 8n^4f(y) + 8n^6f(z) \end{aligned} \quad (41)$$

For all $x, y, z \in X$. Adding $f(n^3z)$ on both sides of (41), we get

$$\begin{aligned} 2f(nx + n^2y) + 2f(n^3z) + 2f(nx - n^2y) + 2f(nx + n^3z) + 2f(nx - n^3z) + 2f(n^2y + n^3z) \\ + 2f(n^2y - n^3z) = 8n^2f(x) + 8n^4f(y) + 8n^6f(z) + 2f(n^3z) \end{aligned} \quad (42)$$

For all $x, y, z \in X$. Using (35), (40) in (42), we arrive

$$\begin{aligned} f(nx + n^2y + n^3z) + f(nx + n^2y - n^3z) + 2f(nx - n^2y) + 2f(nx + n^3z) + 2f(nx - n^3z) \\ + 2f(n^2y + n^3z) + 2f(n^2y - n^3z) = 8n^2f(x) + 8n^4f(y) + 8n^6f(z) + 2n^6f(z) \end{aligned} \quad (43)$$

For all $x, y, z \in X$. Adding $2f(n^3z)$ on both sides of (43), we get

$$\begin{aligned} f(nx + n^2y + n^3z) + f(nx + n^2y - n^3z) + 2f(n^3z) + 2f(nx - n^2y) + 2f(nx + n^3z) + 2f(nx - n^3z) \\ + 2f(n^2y + n^3z) + 2f(n^2y - n^3z) = 8n^2f(x) + 8n^4f(y) + 10n^6f(z) + 2f(n^3z) \end{aligned} \quad (44)$$

For all $x, y, z \in X$. Using (36) in (44), we arrive

$$\begin{aligned} f(nx + n^2y + n^3z) + f(nx + n^2y - n^3z) + f(n^3z + nx - n^2y) + f(n^3z - nx + n^2y) \\ + 2f(nx + n^3z) + 2f(nx - n^3z) + 2f(n^2y + n^3z) + 2f(n^2y - n^3z) \\ = 8n^2f(x) + 8n^4f(y) + 12n^6f(z) \end{aligned} \quad (45)$$

For all $x, y, z \in X$. Adding $2f(n^2y)$ on both sides of (45), we get

$$\begin{aligned} f(nx + n^2y + n^3z) + f(nx + n^2y - n^3z) + f(nx - n^2y + n^3z) + f(-nx + n^2y + n^3z) \\ + 2f(n^2y) + 2f(nx + n^3z) + 2f(nx - n^3z) + 2f(n^2y + n^3z) + 2f(n^2y - n^3z) \\ = 8n^2f(x) + 8n^4f(y) + 2f(n^2y) + 12n^6f(z) \end{aligned} \quad (46)$$

For all $x, y, z \in X$. Using (37) in (46), we arrive

$$f(nx + n^2y + n^3z) + f(nx + n^2y - n^3z) + f(nx - n^2y + n^3z) + f(-nx + n^2y + n^3z) +$$

$$\begin{aligned}
& f(nx + n^2y + n^3z) + f(-nx + n^2y - n^3z) + 2f(nx - n^3z) + 2f(n^2y + n^3z) + 2f(n^2y - n^3z) \\
& = 8n^2f(x) + 10n^4f(y) + 12n^6f(z)
\end{aligned} \tag{47}$$

For all $x, y, z \in X$. Adding $2f(n^2y)$ on both sides of (47), we obtain

$$\begin{aligned}
& f(nx + n^2y + n^3z) + f(nx + n^2y - n^3z) + f(nx - n^2y + n^3z) + f(-nx + n^2y + n^3z) \\
& + f(nx + n^2y + n^3z) + f(-nx + n^2y - n^3z) + 2f(nx - n^3z) + 2f(n^2y) \\
& + 2f(n^2y + n^3z) + 2f(n^2y - n^3z) = 8n^2f(x) + 10n^4f(y) + 2f(n^2y) + 12n^6f(z)
\end{aligned} \tag{48}$$

For all $x, y, z \in X$. Using (38) in (48), we arrive

$$\begin{aligned}
& f(nx + n^2y + n^3z) + f(nx + n^2y - n^3z) + f(nx - n^2y + n^3z) + f(-nx + n^2y + n^3z) \\
& + f(nx + n^2y + n^3z) + f(-nx + n^2y - n^3z) + f(nx + n^2y - n^3z) \\
& + f(-nx + n^2y + n^3z) + 2f(n^2y + n^3z) + 2f(n^2y - n^3z) \\
& = 8n^2f(x) + 12n^4f(y) + 12n^6f(z)
\end{aligned} \tag{49}$$

For all $x, y, z \in X$. Adding $2f(nx)$ on both sides of (49), we get

$$\begin{aligned}
& f(nx + n^2y + n^3z) + f(nx + n^2y - n^3z) + f(nx - n^2y + n^3z) + f(-nx + n^2y + n^3z) \\
& + f(nx + n^2y + n^3z) + f(-nx + n^2y - n^3z) + f(nx + n^2y - n^3z) \\
& + f(-nx + n^2y + n^3z) + 2f(nx) + 2f(n^2y + n^3z) + 2f(n^2y - n^3z) \\
& = 8n^2f(x) + 12n^4f(y) + 12n^6f(z) + 2f(nx)
\end{aligned} \tag{50}$$

For all $x, y, z \in X$. Using (39) in (50), we arrive

$$\begin{aligned}
& f(nx + n^2y + n^3z) + f(nx + n^2y - n^3z) + f(nx - n^2y + n^3z) + f(-nx + n^2y + n^3z) \\
& + f(nx + n^2y + n^3z) + f(-nx + n^2y - n^3z) + f(nx + n^2y - n^3z) + f(-nx + n^2y + n^3z) \\
& + f(nx + n^2y + n^3z) + f(nx - n^2y - n^3z) + 2f(n^2y - n^3z) \\
& = 10n^2f(x) + 12n^4f(y) + 12n^6f(z) + 2f(nx)
\end{aligned} \tag{51}$$

For all $x, y, z \in X$. Adding $2f(nx)$ on both sides of (51), we have

$$\begin{aligned}
& f(nx + n^2y + n^3z) + f(nx + n^2y - n^3z) + f(nx - n^2y + n^3z) + f(-nx + n^2y + n^3z) \\
& + f(nx + n^2y + n^3z) + f(-nx + n^2y - n^3z) + f(nx + n^2y - n^3z) + f(-nx + n^2y + n^3z)
\end{aligned}$$

$$\begin{aligned}
& +f(nx+n^2y+n^3z)+f(nx-n^2y-n^3z)+2f(nx)+2f(n^2y-n^3z) \\
& =10n^2f(x)+12n^4f(y)+12n^6f(z)+2f(nx)
\end{aligned} \tag{52}$$

For all $x, y, z \in X$. Using (40) in (52), we arrive

$$\begin{aligned}
& f(nx+n^2y+n^3z)+f(nx+n^2y-n^3z)+f(nx-n^2y+n^3z)+f(-nx+n^2y+n^3z) \\
& +f(nx+n^2y+n^3z)+f(-(nx-n^2y+n^3z))+f(nx+n^2y-n^3z)+f(-nx+n^2y+n^3z) \\
& +f(nx+n^2y+n^3z)+f(-(-nx+n^2y+n^3z))+f(nx+n^2y-n^3z)+f(nx-n^2y+n^3z) \\
& =12n^2f(x)+12n^4f(y)+12n^6f(z)
\end{aligned} \tag{53}$$

For all $x, y, z \in X$. Using evenness of f in (53), we have

$$\begin{aligned}
& f(nx+n^2y+n^3z)+f(nx-n^2y+n^3z)+f(nx+n^2y-n^3z)+f(-nx+n^2y+n^3z) \\
& =4n^2f(x)+4n^4f(y)+4n^6f(z)
\end{aligned} \tag{54}$$

For all $x, y, z \in X$. Using evenness of f in (54) one can get,

$$\begin{aligned}
& f(nx+n^2y+n^3z)+f(nx-n^2y+n^3z)+f(nx+n^2y-n^3z)+f(-nx+n^2y+n^3z) \\
& =2n^2[f(x)+f(-x)]+2n^4[f(y)+f(-y)]+2n^6[f(z)+f(-z)]
\end{aligned} \tag{55}$$

For all $x, y, z \in X$. Adding $nf(x)+n^2f(y)+n^3f(z)$ on both sides of (55) and using evenness of f , we desired our result.

Conversely, $f : X \rightarrow Y$ satisfies the functional equation (27). Using evenness of f in (27), we have

$$\begin{aligned}
& f(nx+n^2y+n^3z)+f(nx-n^2y+n^3z)+f(nx+n^2y-n^3z)+f(-nx+n^2y+n^3z) \\
& =4n^2f(x)+4n^4f(y)+4n^6f(z)
\end{aligned} \tag{56}$$

Setting (x, y, z) by $(x, 0, 0)$, $(0, x, 0)$ and $(0, 0, x)$ in (56), we obtain

$$f(nx)=n^2f(x); \quad f(n^2x)=n^4f(x) \quad \text{and} \quad f(n^3x)=n^6f(x) \tag{57}$$

For all $x \in X$. It is easy to verify from (57), that

$$f\left(\frac{x}{n^i}\right)=\frac{1}{n^i}f(x), \quad i=1, 2, 3 \tag{58}$$

For all $x \in X$. Replacing (x, y, z) by $\left(\frac{x}{n}, \frac{y}{n^2}, 0\right)$ in (56) and using evenness of f and (58), we desired our result.

For sections 4, 5 and 6, let us consider X and Y to a normed space and a Banach space. Define a mapping $Df : X \rightarrow Y$ by

$$\begin{aligned}
Df(x, y, z) &= f(nx+n^2y+n^3z)+f(nx-n^2y+n^3z)+f(nx+n^2y-n^3z) \\
&+f(-nx+n^2y+n^3z)-n[f(x)-f(-x)]-n^2[f(y)-f(-y)]-n^3[f(z)-f(-z)]
\end{aligned}$$

$$-2n^2[f(x) + f(-x)] - 2n^4[f(y) + f(-y)] - 2n^6[f(z) + f(-z)]$$

For all $x, y, z \in X$.

Banach Space Stability Results of (11)

4. Stability results for (11): odd case – direct method

In this section, we present the generalized Ulam-Hyers stability of the functional equation (11) for odd case.

Theorem 4.1: Let $j \in \{-1, 1\}$ and $\alpha: X^3 \rightarrow [0, \infty)$ be a function such that

$$\sum_{k=0}^{\infty} \frac{\alpha(n^{kj}x, n^{kj}y, n^{kj}z)}{n^{kj}} \text{ Converges in } \mathbb{R} \text{ and } \lim_{k \rightarrow \infty} \frac{\alpha(n^{kj}x, n^{kj}y, n^{kj}z)}{n^{kj}} = 0 \quad (59)$$

For all $x, y, z \in X$. Let $f_a: X \rightarrow Y$ be an odd function satisfying the inequality

$$\|Df_a(x, y, z)\| \leq \alpha(x, y, z) \quad (60)$$

For all $x, y, z \in X$. There exists a unique additive mapping $A: X \rightarrow Y$ which satisfies the functional equation (11) and

$$\|f_a(x) - A(x)\| \leq \frac{1}{2n} \sum_{k=\frac{1-j}{2}}^{\infty} \frac{\alpha(n^{kj}x, 0, 0)}{n^{kj}} \quad (61)$$

For all $x \in X$. The mapping $A(x)$ is defined by

$$A(x) = \lim_{k \rightarrow \infty} \frac{f_a(n^{kj}x)}{n^{kj}} \quad (62)$$

For all $x \in X$.

Proof. Assume that $j = 1$. Replacing (x, y, z) and $(x, 0, 0)$ in (60) and using oddness of f_a , we get

$$\|2f_a(nx) - 2nf_a(x)\| \leq \alpha(x, 0, 0) \quad (63)$$

For all $x \in X$. It follows from (63) that

$$\left\| \frac{f_a(nx)}{n} - f_a(x) \right\| \leq \frac{\alpha}{2n}(x, 0, 0) \quad (64)$$

For all $x \in X$. Replacing x by nx in (64) and dividing by n , we obtain

$$\left\| \frac{f_a(n^2x)}{n^2} - \frac{f_a(nx)}{n} \right\| \leq \frac{\alpha}{2n^2}(nx, 0, 0) \quad (65)$$

For all $x \in X$. It follows from (64) and (65) that

$$\left\| \frac{f_a(n^2x)}{n^2} - f_a(x) \right\| \leq \frac{1}{2n} \left[\alpha(x, 0, 0) + \frac{\alpha}{n}(nx, 0, 0) \right] \quad (66)$$

For all $x \in X$. Generalizing, we have

$$\left\| f_a(x) - \frac{f_a(n^k x)}{n^k} \right\| \leq \frac{1}{2n} \sum_{k=0}^{n-1} \frac{\alpha(n^k x, 0, 0)}{n^k} \leq \frac{1}{2n} \sum_{k=0}^{\infty} \frac{\alpha(n^k x, 0, 0)}{n^k} \quad (67)$$

For all $x \in X$. In order to prove convergence of the sequence $\left\{ \frac{f_a(n^k x)}{n^k} \right\}$, replace x by $n^l x$ and dividing n^l (67), for any $k, l > 0$, to deduce

$$\begin{aligned} \left\| \frac{f_a(n^l x)}{n^l} - \frac{f_a(n^{k+l} x)}{n^{k+l}} \right\| &= \frac{1}{2n} \left\| f_a(n^l x) - \frac{f_a(n^{k+l} x)}{n^k} \right\| \leq \frac{1}{2n} \sum_{k=0}^{n-1} \frac{\alpha(n^{k+l} x, 0, 0)}{n^{k+l}} \\ &\leq \frac{1}{2n} \sum_{k=0}^{\infty} \frac{\alpha(n^{k+l} x, 0, 0)}{n^{k+l}} \rightarrow 0 \text{ as } l \rightarrow \infty \end{aligned} \quad (68)$$

For all $x \in X$. Hence the sequence $\left\{ \frac{f_a(n^k x)}{n^k} \right\}$ is a Cauchy sequence. Since Y is complete, there exists a mapping

$A : X \rightarrow Y$ such that $A(x) = \lim_{k \rightarrow \infty} \frac{f_a(n^k x)}{n^k}$, $\forall x \in X$. Letting $k \rightarrow \infty$ in (67), we see that (61) holds for all

$x \in X$. To prove that A satisfies (11), replacing (x, y, z) by $(n^k x, n^k y, n^k z)$ and dividing n^k in (60), we obtain $\frac{1}{n^k} \|Df_a(n^k x, n^k y, n^k z)\| \leq \frac{1}{n^k} \alpha(n^k x, n^k y, n^k z)$ for all $x, y, z \in X$. Letting $k \rightarrow \infty$ in the above inequality and using the definition of $A(x)$, we see that $DA(x, y, z) = 0$. Hence A satisfies (11) for all $x, y, z \in X$. To show that A is unique, let $B(x)$ be another additive mapping satisfying (11) and (61), then

$$\|A(x) - B(x)\| \leq \frac{1}{n^l} \|A(n^l x) - f_a(n^l x)\| + \|f_a(n^l x) - B(n^l x)\| \leq \frac{1}{2n} \sum_{k=0}^{\infty} \frac{\alpha(n^{k+l} x, 0, 0)}{n^{k+l}} \rightarrow 0 \text{ as } l \rightarrow \infty$$

For all $x \in X$. Hence A is unique. Now, replacing x by $\frac{x}{n}$ in (63), we get

$$\left\| 2f_a(x) - 2nf_a\left(\frac{x}{n}\right) \right\| \leq \alpha\left(\frac{x}{n}, 0, 0\right) \quad (69)$$

For all $x \in X$. It follows from (69) that

$$\left\| f_a(x) - nf_a\left(\frac{x}{n}\right) \right\| \leq \frac{1}{2} \alpha\left(\frac{x}{n}, 0, 0\right) \quad (70)$$

For all $x \in X$. The rest of the proof is similar to that of $j = 1$. Hence for $j = -1$ also the theorem is true. This completes the proof of the theorem.

The following corollary is an immediate consequence of Theorem 4.1 concerning the stability of (11).

Corollary 4.2. Let λ and s be a nonnegative real numbers. Let an odd function $f_a : X \rightarrow Y$ satisfying the inequality

$$\|Df_a(x, y, z)\| \leq \begin{cases} \lambda & s \neq 1; \\ \lambda \left(\|x\|^s + \|y\|^s + \|z\|^s \right), & s \neq \frac{1}{3}; \\ \lambda \left(\|x\|^s \|y\|^s \|z\|^s + \left\{ \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \right\} \right), & s \neq \frac{1}{3}; \end{cases} \quad (71)$$

For all $x, y, z \in X$. Then there exists a unique additive function $A : X \rightarrow Y$ such that

$$\|f_a(x) - A(x)\| \leq \begin{cases} \frac{\lambda}{2|n-1|}, \\ \frac{\lambda \|x\|^s}{2|n-n^s|}, \\ \frac{\lambda \|x\|^{3s}}{2|n-n^{3s}|}, \end{cases} \text{ For all } x \in X. \quad (72)$$

Proof: If we replace

$$\alpha(x, y, z) = \begin{cases} \lambda; \\ \lambda \left(\|x\|^s + \|y\|^s + \|z\|^s \right); \\ \lambda \left(\|x\|^s \|y\|^s \|z\|^s + \left\{ \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \right\} \right); \end{cases} \quad (73)$$

In Theorem 4.1, we arrive (72).

5. Stability result for (11): even case – direct method

In this section, we investigate the generalized Ulam-Hyers stability of the functional equation (11) for even case.

Theorem 5.1: Let $j \in \{-1, 1\}$ and $\alpha : X^3 \rightarrow [0, \infty)$ be a function such that,

$$\sum_{k=0}^{\infty} \frac{\alpha(n^{kj}x, n^{kj}y, n^{kj}z)}{n^{2kj}} \text{ Converges to R and } \lim_{k \rightarrow \infty} \frac{\alpha(n^{kj}x, n^{kj}y, n^{kj}z)}{n^{2kj}} = 0 \quad (74)$$

For all $x, y, z \in X$. Let $f_Q : X \rightarrow Y$ be an even function satisfying the inequality,

$$\|Df_Q(x, y, z)\| \leq \alpha(x, y, z) \quad (75)$$

For all $x, y, z \in X$. Then there exists unique quadratic mapping $Q : X \rightarrow Y$ which satisfies the functional equation (11) and

$$\|f_Q(x) - Q(x)\| \leq \frac{1}{4n^2} \sum_{k=\frac{1-j}{2}}^{\infty} \frac{\alpha(n^{kj}x, 0, 0)}{n^{2kj}} \quad (76)$$

For all $x \in X$. The mapping $Q(x)$ is defined by

$$Q(x) = \lim_{k \rightarrow \infty} \frac{f_Q(n^{kj}x)}{n^{2kj}} \quad (77)$$

For all $x \in X$.

Proof. Assume that $j = 1$, replacing (x, y, z) by $(x, 0, 0)$ in (75) and using of evenness of f_Q , we get

$$\|4f_Q(nx) - 4n^2f_Q(x)\| \leq \alpha(x, 0, 0) \quad (78)$$

For all $x \in X$. The rest of the proof is similar to that Theorem 4.1.

The following corollary is an immediate consequence of Theorem 5.1 concerning the stability of (11).

Corollary 5.2. Let λ and s be a nonnegative real numbers. Let an even function $f_Q : X \rightarrow Y$ satisfying the inequality

$$\|Df_Q(x, y, z)\| \leq \begin{cases} \lambda; \\ \lambda(\|x\|^s + \|y\|^s + \|z\|^s); & s \neq 2; \\ \lambda(\|x\|^s \|y\|^s \|z\|^s + \{\|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s}\}); & s \neq \frac{2}{3}; \end{cases} \quad (79)$$

For all $x, y, z \in X$. Then there exists a unique quadratic function $Q : X \rightarrow Y$ such that,

$$\|f_Q(x) - Q(x)\| \leq \begin{cases} \frac{\lambda}{4|n^2 - 1|}, \\ \frac{\lambda\|x\|^s}{4|n^2 - n^s|}, \\ \frac{\lambda\|x\|^{3s}}{4|n^2 - n^{3s}|}, \end{cases} \quad (80)$$

For all $x \in X$.

6. Stability results for (11): mixed case-direct method

In this section, we establish the generalized Ulam-Hyers stability of the functional equation (11) for mixed case.

Theorem 6.1: Let $j \in \{-1, 1\}$ and $\alpha : X^3 \rightarrow [0, \infty)$ be a function satisfying (59) and (74) for all $x, y, z \in X$. Let $f : X \rightarrow Y$ be a function satisfying the inequality

$$\|Df(x, y, z)\| \leq \alpha(x, y, z) \quad (81)$$

For all $x, y, z \in X$. Then there exists a unique additive mapping and a unique quadratic mapping $Q : X \rightarrow Y$ which satisfies the functional equation (11) and

$$\|f(x) - A(x) - Q(x)\| \leq \frac{1}{2} \left[\frac{1}{2n} \sum_{k=\frac{1-j}{2}}^{\infty} \left(\frac{\alpha(n^{kj}x, 0, 0)}{n^{kj}} + \frac{\alpha(-n^{kj}x, 0, 0)}{n^{kj}} \right) \right]$$

$$+\frac{1}{4n^2} \sum_{k=\frac{1-j}{2}}^{\infty} \left(\frac{\alpha(n^{kj}x, 0, 0)}{n^{2kj}} + \frac{\alpha(-n^{kj}x, 0, 0)}{n^{2kj}} \right) \Bigg] \quad (82)$$

For all $x \in X$. The mapping $A(x)$ and $Q(x)$ is defined in (62) and (77) respectively for all $x \in X$.

Proof. Let $f_o(x) = \frac{f_a(x) - f_a(-x)}{2}$ for all $x \in X$. Then $f_o(0) = 0$ and $f_o(-x) = -f_o(x)$ for all $x \in X$. Hence,

$$\|Df_o(x, y, z)\| \leq \frac{1}{2} (\|Df_o(x, y, z)\| + \|Df_o(-x, -y, -z)\|) \leq \frac{\alpha(x, y, z)}{2} + \frac{\alpha(-x, -y, -z)}{2} \quad (83)$$

For all $x, y, z \in X$. By Theorem 4.1, we have

$$\|f_o(x) - A(x)\| \leq \frac{1}{4n} \sum_{k=\frac{1-j}{2}}^{\infty} \left(\frac{\alpha(n^{kj}x, 0, 0)}{n^{kj}} + \frac{\alpha(-n^{kj}x, 0, 0)}{n^{kj}} \right) \quad (84)$$

For all $x \in X$. Also let, $f_e(x) = \frac{f_q(x) + f_q(-x)}{2}$ for all $x \in X$. Then $f_e(0) = 0$ and $f_e(-x) = f_e(x)$ for all $x \in X$. Hence,

$$\|Df_e(x, y, z)\| \leq \frac{1}{2} (\|Df_q(x, y, z)\| + \|Df_q(-x, -y, -z)\|) \leq \frac{\alpha(x, y, z)}{2} + \frac{\alpha(-x, -y, -z)}{2} \quad (85)$$

For all $x, y, z \in X$. By Theorem 5.1, we have

$$\|f_e(x) - Q(x)\| \leq \frac{1}{8n^2} \sum_{k=\frac{1-j}{2}}^{\infty} \left(\frac{\alpha(n^{kj}x, 0, 0)}{n^{2kj}} + \frac{\alpha(-n^{kj}x, 0, 0)}{n^{2kj}} \right) \quad (86)$$

For all $x \in X$. Define

$$f(x) = f_e(x) + f_o(-x) \quad (87)$$

For all $x \in X$. It follows from (84), (86) and (87), we arrive

$$\begin{aligned} \|f(x) - A(x) - Q(x)\| &= \|f_e(x) + f_o(-x) - A(x) - Q(x)\| \\ &\leq \|f_o(-x) - A(x)\| + \|f_e(x) - Q(x)\| \\ &\leq \frac{1}{4n} \sum_{k=\frac{1-j}{2}}^{\infty} \left(\frac{\alpha(n^{kj}x, 0, 0)}{n^{kj}} + \frac{\alpha(-n^{kj}x, 0, 0)}{n^{kj}} \right) \\ &\quad + \frac{1}{8n^2} \sum_{k=\frac{1-j}{2}}^{\infty} \left(\frac{\alpha(n^{kj}x, 0, 0)}{n^{2kj}} + \frac{\alpha(-n^{kj}x, 0, 0)}{n^{2kj}} \right) \end{aligned} \quad (88)$$

For all $x \in X$. Hence the theorem is proved.

Using Corollaries 4.2 and 5.2, we have the following corollary concerning the stability of (11).

Corollary 6.2 Let λ and s be a nonnegative real numbers. Let a function $f : X \rightarrow Y$ satisfying the inequality

$$\|Df(x, y, z)\| \leq \begin{cases} \lambda; & s \neq 1, 2; \\ \lambda(\|x\|^s + \|y\|^s + \|z\|^s); & s \neq \frac{1}{3}, \frac{2}{3}; \\ \lambda(\|x\|^s \|y\|^s \|z\|^s + \{ \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \}); & s \neq \frac{1}{3}, \frac{2}{3}; \end{cases} \quad (89)$$

For all $x, y, z \in X$. Then there exists a unique additive mapping $A: X \rightarrow Y$ and a unique quadratic mapping $Q: X \rightarrow Y$ such that

$$\|f(x) - A(x) - Q(x)\| \leq \begin{cases} \frac{\lambda}{2} \left[\frac{1}{|n-1|} + \frac{1}{2|n^2-1|} \right], \\ \frac{\lambda \|x\|^s}{2} \left[\frac{1}{|n-n^s|} + \frac{1}{2|n^2-n^s|} \right], \\ \frac{\lambda \|x\|^{3s}}{2} \left[\frac{1}{|n-n^{3s}|} + \frac{1}{2|n^2-n^{3s}|} \right], \end{cases} \quad (90)$$

For all $x \in X$.

For Sections 7, 8 and 9, let us consider W and B to a normed space and a Banach space, respectively. Define a mapping

$Df: W \rightarrow B$ by

$$Df(x, y, z) = f(nx + n^2y + n^3z) + f(nx - n^2y + n^3z) + f(nx + n^2y - n^3z)$$

$$+ f(-nx + n^2y + n^3z) - n[f(x) - f(-x)] - n^2[f(y) - f(-y)] - n^3[f(z) - f(-z)]$$

$$- 2n^2[f(x) + f(-x)] - 2n^4[f(y) + f(-y)] - 2n^6[f(z) + f(-z)]$$

For all $x, y, z \in W$.

Fixed Point Stability Results of (11)

The following theorems are useful to prove our fixed point stability results.

Theorem A: (Banach contraction principle) [37] Let (X, d) be a complete metric spaces and consider a mapping $T: X \rightarrow X$ which is strictly contractive mapping, that is,

$$(A_1) d(Tx, Ty) \leq d(x, y)$$

For some (Lipschitz constant) $L < 1$, then,

i) The mapping T has one and only fixed point $x^* = T(x^*)$.

ii) The fixed point for each given element x^* is globally contractive that is

$$(A_2) \lim_{n \rightarrow \infty} T^n x = x^*$$

For any starting point $x \in X$.

iii) One has the following estimation inequalities,

$$(A_3) d(T^n x, x^*) \leq \frac{1}{1-L} d(T^n x, T^{n+1} x), \quad \forall n \geq 0, \forall x \in X.$$

$$(A_4) d(x, x^*) = \frac{1}{1-L} d(x, T x), \quad \forall x \in X.$$

Theorem B: (The Alternative Fixed Point)[37] Suppose that for a complete generalized metric space (X, d) and a strictly contractive mapping $T : X \rightarrow X$ with Lipschitz constant L , then for each given element $x \in X$ either,

$$(B_1) d(T^n x, T^{n+1} x) = \infty, \quad \forall n \geq 0.$$

(B_2) There exists a natural number n_0 such that,

$$i) \quad d(T^n x, T^{n+1} x) < \infty \text{ for all } \forall n \geq 0.$$

ii) The sequence $\{T^n x\}$ is convergent to a fixed point y^* of T ,

iii) y^* is the unique fixed point of T in the set $Y = \{y \in Y; d(T^{n_0} x, y) < \infty\}$.

$$iv) \quad d(y^*, y) \leq \frac{1}{1-L} d(y, Ty) \text{ for all } y \in Y.$$

7. Fixed point stability of (11): odd case-fixed point method

In this section, we present the generalized Ulam-Hyers stability of the functional equation (11) for odd case using fixed point method.

Theorem 7.1: Let $f_a : W \rightarrow B$ be an odd mapping for which there exists a function $\alpha : W^3 \rightarrow [0, \infty)$ with the condition

$$\lim_{k \rightarrow \infty} \frac{\alpha(\eta_i^k x, \eta_i^k y, \eta_i^k z)}{\eta_i^k} = 0 \quad (91)$$

Where

$$\eta_i = \begin{cases} n, & i = 0; \\ \frac{1}{n}, & i = 1; \end{cases}$$

Such that the functional inequality

$$\|Df_a(x, y, z)\| \leq \alpha(x, y, z) \quad (92)$$

For all $x, y, z \in W$. If there exist $L = L(i)$ such that the function

$$x \rightarrow \beta(x) = \frac{1}{2} \alpha\left(\frac{x}{n}, 0, 0\right)$$

Has the property,

$$\frac{1}{\eta_i} \beta(\eta_i x) = L \beta(x) \quad (93)$$

For all $x \in W$. Then there exists a unique additive function $A : W \rightarrow B$ satisfying the functional equation (11) and

$$\|f_a(x) - A(x)\| \leq \frac{L^{1-i}}{1-L} \beta(x) \quad (94)$$

Holds for all $x \in W$.

Proof: Consider the set $X = \{P/P:W \rightarrow B, P(0)=0\}$ and introduce the generalized metric on X . $d(p,q) = \inf \{K \in (0,\infty): \|p(x)-q(x)\| \leq K\beta(x), x \in W\}$. It is easy to see that (X, d) is complete. Define $T: X \rightarrow X$ by $Tp(x) = \frac{1}{\eta_i} p(\eta_i x)$ for all $x \in W$. Now $p, q \in X$,

$$\begin{aligned} d(p,q) \leq K &\Rightarrow \|p(x)-q(x)\| \leq K\beta(x), & x \in W; \\ &\Rightarrow \left\| \frac{1}{\eta_i} p(\eta_i x) - \frac{1}{\eta_i} q(\eta_i x) \right\| \leq \frac{1}{\eta_i} K\beta(\eta_i x), & x \in W; \\ &\Rightarrow \|Tp(x)-Tq(x)\| \leq LK\beta(x), & x \in W; \\ &\Rightarrow d(Tp,Tq) \leq LK. \end{aligned}$$

This implies $d(Tp,Tq) \leq Ld(p,q)$ for all $p,q \in X$. (i.e.,) T is strictly contractive mapping on X with Lipschitz constant L . It follows from (63) that

$$\|2f_a(nx) - 2nf_a(x)\| \leq \alpha(x, 0, 0) \quad (95)$$

For all $x \in W$. It follows from (95) that

$$\left\| f_a(x) - \frac{f_a(nx)}{n} \right\| \leq \frac{1}{2n} \alpha(x, 0, 0) \quad (96)$$

For all $x \in W$. Using (93), for the case $i = 0$, it reduces to

$$\left\| f_a(x) - \frac{1}{n} f_a(nx) \right\| \leq \frac{1}{n} \beta(x) \quad (97)$$

For all $x \in W$.

$$(i.e.) \ d(f_a, Tf_a) \leq \frac{1}{n} \Rightarrow d(f_a, Tf_a) \leq \frac{1}{n} = L = L^1 < \infty.$$

Again replacing $x = \frac{x}{n}$ in (95), we get

$$\left\| f_a(x) - nf_a\left(\frac{x}{n}\right) \right\| \leq \frac{1}{2} \alpha\left(\frac{x}{n}, 0, 0\right) \quad (98)$$

For all $x \in W$. Using (93) for the case $i = 1$, it reduces to,

$$\left\| f_a(x) - nf_a\left(\frac{x}{n}\right) \right\| \leq \beta(x) \quad (99)$$

For all $x \in W$.

$$(i.e.,) \ d(f_a, Tf_a) \leq 1 \Rightarrow d(f_a, Tf_a) \leq 1 = L^0 < \infty.$$

In above case, we arrive $d(f_a, Tf_a) \leq L^{1-i}$

Therefore $(B_2(i))$ holds. By $(B_2(ii))$, it follows that there exists a fixed point A of T in X , such that

$$A(x) = \lim_{k \rightarrow \infty} \frac{f_a(\eta_i^k x)}{\eta_i^k}, \quad \forall x \in W. \quad (100)$$

In order to prove $A:W \rightarrow B$ is additive. Replacing (x, y, z) by $(\eta_i^k x, \eta_i^k y, \eta_i^k z)$ in (92) and dividing by η_i^k , it follows from (91) and (100), we see that A satisfies (11) for all $x, y, z \in W$. Hence A satisfies the functional equation (11).

By $(B_2(iii))$, A is the unique fixed point of T in the set, $Y = \{f_a \in X : d(Tf_a, A) < \infty\}$. Using the fixed-point alternative result, A is the unique function such that, $\|f_a(x) - A(x)\| \leq K\beta(x)$ for all $x \in W$, and $k > 0$. Finally by

$$(B_2(iv)), \text{ we obtain } d(f_a, A) \leq \frac{1}{1-L} d(f_a, Tf_a)$$

$$(i.e.) d(f_a, A) \leq \frac{L^{1-i}}{1-L}.$$

Hence, we conclude that

$$\|f_a(x) - A(x)\| \leq \frac{L^{1-i}}{1-L} \beta(x)$$

For all $x \in W$. This completes the proof of the theorem.

Corollary 7.2. Let $f_a:W \rightarrow B$ be an odd mapping and there exists a real numbers λ and s such that,

$$\|Df_a(x, y, z)\| \leq \begin{cases} \lambda, \\ \lambda \left\{ \|x\|^s + \|y\|^s + \|z\|^s \right\}; \\ \lambda \left\{ \|x\|^s \|y\|^s \|z\|^s + \left\{ \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \right\} \right\}; \end{cases} \quad (101)$$

For all $x, y, z \in W$. There exists a unique additive mapping $A:W \rightarrow B$ such that

$$\|f_a(x) - A(x)\| \leq \begin{cases} \frac{\lambda}{2^{\lfloor n-1 \rfloor}}, \\ \frac{\lambda \|x\|^s}{2^{\lfloor n-n^s \rfloor}}, & s \neq 1 \\ \frac{\lambda \|x\|^{3s}}{2^{\lfloor n-n^{3s} \rfloor}}, & s \neq \frac{1}{3} \end{cases} \quad (102)$$

For all $x \in W$.

Proof: Setting

$$\alpha(x, y, z) = \begin{cases} \lambda, \\ \lambda \left\{ \|x\|^s + \|y\|^s + \|z\|^s \right\}; \\ \lambda \left\{ \|x\|^s \|y\|^s \|z\|^s + \left\{ \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \right\} \right\}; \end{cases} \quad \text{for all } x, y, z \in W. \text{ Now}$$

$$\frac{\alpha(\eta_i^k x, \eta_i^k y, \eta_i^k z)}{\eta_i^k}$$

$$= \begin{cases} \frac{\lambda}{\eta_i^k}, \\ \frac{\lambda}{\eta_i^k} \left\{ \left\| \eta_i^k x \right\|^s + \left\| \eta_i^k y \right\|^s + \left\| \eta_i^k z \right\|^s \right\}; \\ \frac{\lambda}{\eta_i^k} \left\{ \left\| \eta_i^k x \right\|^s \left\| \eta_i^k y \right\|^s \left\| \eta_i^k z \right\|^s + \left\{ \left\| \eta_i^k x \right\|^{3s} + \left\| \eta_i^k y \right\|^{3s} + \left\| \eta_i^k z \right\|^{3s} \right\} \right\}; \end{cases} = \begin{cases} \rightarrow 0 \text{ as } k \rightarrow \infty, \\ \rightarrow 0 \text{ as } k \rightarrow \infty, \\ \rightarrow 0 \text{ as } k \rightarrow \infty, \end{cases} \quad (103)$$

i.e., (91) is holds. But we have $\beta(x) = \frac{1}{2}\alpha\left(\frac{x}{n}, 0, 0\right)$. Hence, $\beta(x) = \frac{1}{2}\alpha\left(\frac{x}{n}, 0, 0\right) = \begin{cases} \frac{\lambda}{2} \\ \frac{\lambda}{2n^s} \|x\|^s \\ \frac{\lambda}{2n^{3s}} \|x\|^{3s} \end{cases}$

Also,

$$\frac{1}{\eta_i} \beta(\eta_i x) = \begin{cases} \frac{\lambda}{2\eta_i} \\ \frac{\lambda}{2\eta_i} \left\| \eta_i x \right\|^s \\ \frac{\lambda}{2\eta_i} \left\| \eta_i x \right\|^{3s} \end{cases} = \begin{cases} \eta_i^{-1} \beta(x) \\ \eta_i^{s-1} \beta(x) \\ \eta_i^{3s-1} \beta(x) \end{cases} \quad (104)$$

Hence the inequality (93) holds

Either $L = n^{-1}$ for $s = 0$ if $i = 0$ and $L = \frac{1}{n-1}$ for $s = 0$ if $i = 1$.

Either $L = n^{s-1}$ for $s < 1$ if $i = 0$ and $L = \frac{1}{n^{s-1}}$ for $s > 1$ if $i = 1$.

Either $L = n^{3s-1}$ for $s < 1$ if $i = 0$ and $L = \frac{1}{n^{3s-1}}$ for $s > 1$ if $i = 1$.

Now from (94), we prove the following cases:

Case: 1 $L = n^{-1}$, $i = 0$

$$\|f_a(x) - A(x)\| \leq \frac{L^{1-i}}{1-L} \beta(x) = \frac{(n^{-1})^{1-0}}{1-n^{-1}} \frac{\lambda}{2} = \frac{\lambda}{2(n-1)}. \quad (105)$$

Case: 2 $L = \left(\frac{1}{n}\right)^{-1}$, $i = 1$

$$\|f_a(x) - A(x)\| \leq \frac{L^{1-i}}{1-L} \beta(x) = \frac{(n)^{1-1}}{1-n} \frac{\lambda}{2} = \frac{\lambda}{2(1-n)}. \quad (106)$$

Case: 3 $L = n^{s-1}$, $s < 1$, $i = 0$

$$\|f_a(x) - A(x)\| \leq \frac{L^{1-i}}{1-L} \beta(x) = \frac{(n^{s-1})^{1-0}}{1-n^{s-1}} \frac{\lambda}{2n^s} \|x\|^s = \frac{\lambda \|x\|^s}{2(n-n^s)}. \quad (107)$$

Case: 4 $L = \left(\frac{1}{n}\right)^{s-1}$, $s > 1$, $i = 1$

$$\|f_a(x) - A(x)\| \leq \frac{L^{1-i}}{1-L} \beta(x) = \frac{(n^{1-s})^{1-1}}{1-n^{1-s}} \frac{\lambda}{2n^s} \|x\|^s = \frac{\lambda \|x\|^s}{2(n^s - n)}. \quad (108)$$

Case: 5 $L = n^{3s-1}$, $s < \frac{1}{3}$, $i = 0$

$$\|f_a(x) - A(x)\| \leq \frac{L^{1-i}}{1-L} \beta(x) = \frac{(n^{3s-1})^{1-0}}{1-n^{3s-1}} \frac{\lambda}{2n^{3s}} \|x\|^s = \frac{\lambda \|x\|^s}{2(n - n^{3s})}. \quad (109)$$

Case: 6 $L = \left(\frac{1}{n}\right)^{3s-1}$, $s > \frac{1}{3}$, $i = 1$

$$\|f_a(x) - A(x)\| \leq \frac{L^{1-i}}{1-L} \beta(x) = \frac{(n^{1-3s})^{1-1}}{1-n^{1-3s}} \frac{\lambda}{2n^{3s}} \|x\|^s = \frac{\lambda \|x\|^s}{2(n^{3s} - n)}. \quad (110)$$

Hence the proof of the corollary.

8. Fixed point stability of (11): even case-fixed point method

In this section, we give the generalized Ulam-Hyers stability of the functional equation (11), for even case.

Theorem 8.1: Let $f_Q : W \rightarrow B$ be an even mapping for which there exists a function $\alpha : W^3 \rightarrow [0, \infty)$ with the condition

$$\lim_{k \rightarrow \infty} \frac{\alpha(\eta_i^k x, \eta_i^k y, \eta_i^k z)}{\eta_i^{2k}} = 0 \quad (111)$$

Where

$$\eta_i = \begin{cases} n, & i = 0; \\ \frac{1}{n}, & i = 1; \end{cases}$$

Such that the functional inequality with

$$\|Df_Q(x, y, z)\| \leq \alpha(x, y, z) \quad (112)$$

For all $x, y, z \in W$. If there exist $L = L(i) < 1$ such that the function

$$x \rightarrow \beta(x) = \frac{1}{2} \alpha\left(\frac{x}{n}, 0, 0\right)$$

Has the property,

$$\frac{1}{\eta_i^2} \beta(\eta_i x) = L \beta(x) \quad (113)$$

For all $x \in W$. Then there exists a unique quadratic function $Q : W \rightarrow B$ satisfying the functional equation (11) and

$$\|f_q(x) - Q(x)\| \leq \frac{L^{1-i}}{1-L} \beta(x) \quad (114)$$

Holds for all $x \in W$.

Proof: Consider the set $X = \{P/P : W \rightarrow B, P(0) = 0\}$ and introduce the generalized metric on X .

$$d(p, q) = \inf \{K \in (0, \infty) : \|p(x) - q(x)\| \leq K \beta(x), x \in W\}$$

It is easy to see that (X, d) is complete. Define $T : X \rightarrow X$ by

$$Tp(x) = \frac{1}{\eta_i^2} p(\eta_i x) \text{ for all } x \in W. \text{ Now } p, q \in X,$$

$$d(p, q) \leq K \Rightarrow \|p(x) - q(x)\| \leq K \beta(x); \quad x \in W$$

$$\Rightarrow \left\| \frac{1}{\eta_i^2} p(\eta_i x) - \frac{1}{\eta_i^2} q(\eta_i x) \right\| \leq \frac{1}{\eta_i^2} K \beta(\eta_i x); x \in W$$

$$\Rightarrow \left\| \frac{1}{\eta_i^2} p(\eta_i x) - \frac{1}{\eta_i^2} q(\eta_i x) \right\| \leq LK \beta(x); x \in W$$

$$\Rightarrow \|Tp(x) - Tq(x)\| \leq LK \beta(x); x \in W$$

$$\Rightarrow d(Tp, Tq) \leq LK.$$

This implies $d(Tp, Tq) \leq Ld(p, q)$ for all $p, q \in X$.

(i.e.) T is strictly contractive mapping on X with Lipschitz constant L . Replacing (x, y, z) by $(x, 0, 0)$ in (112) and using evenness of f , we get

$$\|4f(nx) - 4n^2 f(x)\| \leq \alpha(x, 0, 0) \quad (115)$$

$$\left\| f(x) - \frac{f(nx)}{n^2} \right\| \leq \frac{1}{4n^2} \alpha(x, 0, 0) \quad (116)$$

For all $x \in W$, using (113) for the case $i = 0$, it reduces to

$$\left\| f_q(x) - \frac{1}{n^2} f_q(nx) \right\| \leq \frac{1}{2n^2} \beta(x) \quad (117)$$

For all $x \in W$,

$$(i.e.) d(f_q, Tf_q) \leq \frac{1}{n^2} \Rightarrow d(f_q, Tf_q) \leq \frac{1}{2n^2} = L = L^1 < \infty.$$

Again replacing $x = \frac{x}{n}$ in (115), we get

$$\left\| f_q(x) - n^2 f_q\left(\frac{x}{n}\right) \right\| \leq \frac{1}{4} \alpha\left(\frac{x}{n}, 0, 0\right) \quad (118)$$

For all $x \in W$, using (113) for the case $i = 1$, it reduces to

$$\left\| f_q(x) - n^2 f_q\left(\frac{x}{n}\right) \right\| \leq \frac{1}{2} \beta(x); \quad (119)$$

For all $x \in W$,

$$(\text{i.e.}) \ d(f_q, Tf_q) \leq \frac{1}{2} < 1 \Rightarrow d(f_q, Tf_q) \leq 1 \Rightarrow L^0 < \infty.$$

In above cases, we arrive

$$d(f_q, Tf_q) \leq L^{1-i}.$$

The rest of the proof is similar to that of Theorem 7.1 This completes the proof of the theorem.

Corollary 8.2: Let $f_q : W \rightarrow B$ be an even mapping and there exists a real numbers λ and s such that,

$$\|Df_q(x, y, z)\| \leq \begin{cases} \lambda, & s \neq 2; \\ \lambda \left\{ \|x\|^s + \|y\|^s + \|z\|^s \right\}, & s \neq \frac{1}{3}; \\ \lambda \left\{ \|x\|^s \|y\|^s \|z\|^s + \left(\|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \right) \right\}, & s \neq \frac{1}{3}; \end{cases} \quad (120)$$

For all $x, y, z \in W$. There exists a unique quadratic mapping $Q : W \rightarrow B$ such that

$$\|f_q(x) - Q(x)\| \leq \begin{cases} \frac{\lambda}{4|n^2 - 1|}, \\ \frac{\lambda \|x\|^s}{4|n^2 - n^s|}, \\ \frac{\lambda \|x\|^{3s}}{4|n^2 - n^{3s}|}, \end{cases} \quad (121)$$

For all $x \in W$.

9. Fixed point stability of (11): mixed case-fixed point method

In this section, we discuss the generalized Ulam-Hyers stability of the functional equation (11) for the mixed case.

Theorem 9.1: Let $f : W \rightarrow B$ be a mapping for which there exists a function $\alpha : W^3 \rightarrow [0, \infty)$ with the condition (91) and (111) where

$$\eta_i = \begin{cases} n, & i = 0; \\ \frac{1}{n}, & i = 1; \end{cases}$$

Such that the functional inequality with

$$\|Df(x, y, z)\| \leq \alpha(x, y, z) \quad (122)$$

For all $x, y, z \in W$. If there exist $L = L(i)$ such that the function for all $x \in W$, such that the function

$$x \rightarrow \beta(x) = \frac{1}{2} \alpha\left(\frac{x}{n}, 0, 0\right),$$

Has the properties (93) and (113) for all $x \in W$. Then there exists a unique additive function $A: W \rightarrow B$ and a unique quadratic function $Q: W \rightarrow B$ satisfying the functional equation (11) and

$$\|f(x) - A(x) - Q(x)\| \leq \frac{L^{1-i}}{1-L} [\beta(x) + \beta(-x)] \quad (123)$$

Holds for all $x \in W$.

Proof: It follows from (83) and Theorem 7.1, that

$$\|f_o(x) - A(x)\| \leq \frac{1}{2} \frac{L^{1-i}}{1-L} [\beta(x) + \beta(-x)] \quad (124)$$

Similarly, it is follows from (85) and Theorem 8.1, that

$$\|f_e(x) - Q(x)\| \leq \frac{1}{2} \frac{L^{1-i}}{1-L} [\beta(x) + \beta(-x)] \quad (125)$$

For all $x \in W$. Define

$$f(x) = f_o(x) + f_e(x) \quad (126)$$

For all $x \in W$. From (124), (125) and (126), we have

$$\begin{aligned} \|f(x) - A(x) - Q(x)\| &\leq \|f_e(x) + f_o(x) - A(x) - Q(x)\| \\ &\leq \|f_o(x) - A(x)\| + \|f_e(x) - Q(x)\| = \frac{L^{1-i}}{1-L} [\beta(x) + \beta(-x)] \end{aligned}$$

For all $x \in W$. Hence the theorem is proved.

Using Corollaries 7.2 and 8.2 we have the following corollary concerning the stability of (11).

Corollary 9.2: Let $f: W \rightarrow B$ be a mapping and there exists a real numbers λ and s such that,

$$\|Df(x, y, z)\| \leq \begin{cases} \lambda, & s \neq 1, 2 \\ \lambda \left\{ \|x\|^s + \|y\|^s + \|z\|^s \right\}, & s \neq \frac{1}{3}, \frac{2}{3} \\ \lambda \left\{ \|x\|^s \|y\|^s \|z\|^s + \left(\|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \right) \right\}, & s \neq \frac{1}{3}, \frac{2}{3} \end{cases} \quad (127)$$

For all $x, y, z \in W$. There exists a unique additive mapping $A: W \rightarrow B$ and a unique quadratic mapping $Q: W \rightarrow B$ such that,

$$\|f(x) - A(x) - Q(x)\| \leq \begin{cases} \frac{\lambda}{2|n-1|} + \frac{\lambda}{4|n^2-1|}, \\ \frac{\lambda\|x\|^s}{2|n-n^s|} + \frac{\lambda\|x\|^s}{4|n^2-n^s|}, \\ \frac{\lambda\|x\|^{3s}}{2|n-n^{3s}|} + \frac{\lambda\|x\|^{3s}}{4|n^2-n^{3s}|}, \end{cases} \quad \text{for all } x \in W. \quad (128)$$

Banach Algebra Stability Results For (11)

For sections 10, 11 and 12, let us consider X and Y to a normed Algebra and a Banach Algebra, respectively. Define a mapping $Df : X \rightarrow Y$ by,

$$\begin{aligned} Df(x, y, z) &= f(nx + n^2y + n^3z) + f(nx - n^2y + n^3z) + f(nx + n^2y - n^3z) \\ &+ f(-nx + n^2y + n^3z) - n[f(x) - f(-x)] - n^2[f(y) - f(-y)] - n^3[f(z) - f(-z)] \\ &- 2n^2[f(x) + f(-x)] - 2n^4[f(y) + f(-y)] - 2n^6[f(z) + f(-z)] \end{aligned}$$

For all $x, y, z \in X$.

10. Stability results for (11): odd case-direct method

In this section, we present the generalized Ulam-Hyers stability of the functional equation (11), when f is odd.

Definition 10.1: Let X be Banach Algebra. A mapping $A : X \rightarrow X$ is said to be Additive derivation if the Additive function A satisfies,

$$A(ab) = A(a)b + aA(b) \quad (129)$$

For all $a, b \in X$. Also the additive derivation for three variables satisfies

$$A(abc) = A(a)bc + aA(b)c + abA(c) \quad (130)$$

For all $a, b, c \in X$.

Theorem 10.2: Let $j = \pm 1$. Let $f_a : X \rightarrow Y$ be a odd mapping for which there exists function $\alpha, \beta : X^3 \rightarrow [0, \infty)$ with the condition

$$\sum_{k=0}^{\infty} \frac{\alpha(n^{kj}x, n^{kj}y, n^{kj}z)}{n^{kj}} \text{ Converges in } \mathbb{R} \text{ and } \lim_{k \rightarrow \infty} \frac{\alpha(n^{kj}x, n^{kj}y, n^{kj}z)}{n^{kj}} = 0 \quad (131)$$

$$\sum_{k=0}^{\infty} \frac{\beta(n^{kj}x, n^{kj}y, n^{kj}z)}{n^{3kj}} \text{ Converges in } \square \text{ and } \lim_{k \rightarrow \infty} \frac{\beta(n^{kj}x, n^{kj}y, n^{kj}z)}{n^{3kj}} = 0 \quad (132)$$

Such that the functional inequalities

$$\|Df_a(x, y, z)\| \leq \alpha(x, y, z) \quad (133)$$

And

$$\|f_a(xyz) - f_a(x)yz - xf_a(y)z - xyf_a(z)\| \leq \beta(x, y, z) \quad (134)$$

For all $x, y, z \in X$. Then there exists a unique Additive Derivation mapping $A: X \rightarrow Y$ satisfying the functional equation (11) and

$$\|f_a(x) - A(x)\| \leq \frac{1}{2n} \sum_{k=\frac{1-j}{2}}^{\infty} \frac{\alpha(n^{kj}x, 0, 0)}{n^{kj}} \quad (135)$$

For all $x \in X$. The mapping $A(x)$ is defined by

$$A(x) = \lim_{n \rightarrow \infty} \frac{f_a(n^{kj}x)}{n^{kj}} \quad (136)$$

For all $x \in X$.

Proof: It follows from Theorem 4.1 that A is a unique additive mapping and satisfies (11) for all $x, y, z \in X$. It follows from (134) that

$$\begin{aligned} & \|A(xyz) - A(x)yz - xA(y)z - xyA(z)\| \\ & \leq \frac{1}{n^{3k}} \|f_a(n^k(xyz)) - f_a(n^kx)(n^ky n^kz) - n^kx f_a(n^ky) n^kz - n^kx n^ky f_a(n^kz)\| \\ & \leq \frac{1}{n^{3k}} \beta(n^kx, n^ky, n^kz) \rightarrow 0 \text{ As } k \rightarrow \infty \end{aligned}$$

For all $x, y, z \in X$. Hence, the mapping $A: X \rightarrow Y$ is a unique Additive Derivation satisfying (135).

This following corollary is a immediate consequence of Theorem 10.2 concerning the stability of (11).

Corollary 10.3: Let $f_a: X \rightarrow Y$ be a odd mapping and there exists a real numbers λ and s such that

$$\|Df_a(x, y, z)\| \leq \begin{cases} \lambda; & s \neq 1 \\ \lambda(\|x\|^s + \|y\|^s + \|z\|^s); & s \neq \frac{1}{3} \\ \lambda(\|x\|^s \|y\|^s \|z\|^s + \{ \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \}); & s \neq \frac{1}{3} \end{cases} \quad (137)$$

$$\|f_a(xyz) - f_a(x)yz - xf_a(y)z - xyf_a(z)\|$$

$$\leq \begin{cases} \lambda; \\ \lambda(\|x\|^s + \|y\|^s + \|z\|^s); \\ \lambda(\|x\|^s \|y\|^s \|z\|^s + \{ \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \}); \end{cases} \quad (138)$$

For all $x, y, z \in X$. Then there exists a unique additive derivation $A: X \rightarrow Y$ such that

$$\|f_a(x) - A(x)\| \leq \begin{cases} \frac{\lambda}{2|n-1|}, \\ \frac{\lambda \|x\|^s}{2|n-n^s|}, \\ \frac{\lambda \|x\|^{3s}}{2|n-n^{3s}|}, \end{cases} \quad (139)$$

For all $x \in X$.

11. Stability results for (11): even case-direct method

In this section, we discuss the generalized Ulam-Hyers stability of the functional equation (11), when f is even.

Definition 11.1: Let X be Banach Algebra. A mapping $Q: X \rightarrow X$ is said to be quadratic derivation if the quadratic function Q satisfies

$$Q(ab) = Q(a)b^2 + a^2Q(b) \quad (140)$$

For all $a, b \in X$. Also the quadratic derivation for three variables satisfies

$$Q(abc) = Q(a)b^2c^2 + a^2Q(b)c^2 + a^2b^2Q(c) \quad (141)$$

For all $a, b, c \in X$.

Theorem 11.2: Let $j = \pm 1$. Let $f_j: X \rightarrow Y$ be an even mapping for which there exists functions $\alpha, \beta: X^3 \rightarrow [0, \infty)$ with the conditions

$$\sum_{k=0}^{\infty} \frac{\alpha(n^{kj}x, n^{kj}y, n^{kj}z)}{n^{2kj}} \text{ Converges in } \mathbb{R} \text{ and } \lim_{k \rightarrow \infty} \frac{\alpha(n^{kj}x, n^{kj}y, n^{kj}z)}{n^{2kj}} = 0 \quad (142)$$

$$\sum_{k=0}^{\infty} \frac{\beta(n^{kj}x, n^{kj}y, n^{kj}z)}{n^{6kj}} \text{ Converges in } \mathbb{R} \text{ and } \lim_{k \rightarrow \infty} \frac{\beta(n^{kj}x, n^{kj}y, n^{kj}z)}{n^{6kj}} = 0 \quad (143)$$

Such that the functional inequalities

$$\|Df_Q(x, y, z)\| \leq \alpha(x, y, z) \quad (144)$$

And

$$\|f_Q(xyz) - y^2z^2f_Q(x) - x^2f_Q(y)z^2 - x^2y^2f_Q(z)\| \leq \beta(x, y, z) \quad (145)$$

For all $x, y, z \in X$. There exists a unique quadratic derivation $Q: X \rightarrow Y$ satisfying the functional equation (11) and

$$\|f_Q(x) - Q(x)\| \leq \frac{1}{4n^2} \sum_{k=0}^{\infty} \frac{\alpha(n^{kj}x, 0, 0)}{n^{2kj}} \quad (146)$$

For all $x \in X$. The mapping $Q(x)$ is defined by

$$Q(x) = \lim_{k \rightarrow \infty} \frac{f_q(n^{kj}x)}{n^{2kj}} \quad (147)$$

For all $x \in X$.

Proof: It follows from Theorem 5.1 that Q is a unique quadratic mapping and satisfies (11) for all $x, y, z \in X$. It follows from (145) that

$$\begin{aligned} & \|Q(xyz) - Q(x)y^2z^2 - x^2Q(y)z^2 - x^2y^2Q(z)\| \\ & \leq \frac{1}{n^{6k}} \|f_q(n^k(xyz)) - f_q(n^kx)n^{2k}y^2n^{2k}z^2 - n^{2k}xf_q(n^ky)n^{2k}z^2 - n^{2k}x^2n^{2k}yf_q(n^kz)\| \\ & \leq \frac{1}{n^{6k}} \beta(n^kx, n^ky, n^kz) \rightarrow 0 \text{ As } k \rightarrow \infty \end{aligned}$$

For all $x, y, z \in X$. Hence, the mapping $Q : X \rightarrow Y$ is a unique Quadratic Derivation satisfying (146).

Corollary 11.3: Let $f_q : X \rightarrow Y$ be a even mapping and there exists a real numbers λ and s such that,

$$\|Df_q(x, y, z)\| \leq \begin{cases} \lambda; & s \neq 2; \\ \lambda(\|x\|^s + \|y\|^s + \|z\|^s); & s \neq \frac{2}{3}; \end{cases} \quad (148)$$

$$\begin{aligned} & \|f_q(xyz) - y^2z^2f_q(x) - x^2f_q(y)z^2 - x^2y^2f_q(z)\| \\ & \leq \begin{cases} \lambda; \\ \lambda(\|x\|^s + \|y\|^s + \|z\|^s); \\ \lambda(\|x\|^s\|y\|^s\|z\|^s + \{ \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \}); \end{cases} \end{aligned} \quad (149)$$

For all $x \in X$. Then there exists a unique quadratic derivation $Q : X \rightarrow Y$ such that

$$\|f_q(x) - Q(x)\| \leq \begin{cases} \frac{\lambda}{4|n^2 - 1|}; \\ \frac{\lambda\|x\|^s}{4|n^2 - n^s|}; \\ \frac{\lambda\|x\|^s}{4|n^2 - n^{3s}|}; \end{cases} \quad (150)$$

For all $x \in X$.

12. Stability results for (11): mixed case-direct method

In this section, we investigate the generalized Ulam-Hyers stability of the functional equation (11), when f is mixed case.

Theorem 12.1: Let $j = \pm 1$. Let $f : X \rightarrow Y$ be a mapping for which there exists functions $\alpha, \beta : X^3 \rightarrow [0, \infty)$ with the conditions (131), (132) and (142), (143) such that the functional inequalities,

$$\|Df(x, y, z)\| \leq \alpha(x, y, z) \quad (151)$$

(134) and (145) for all $x, y, z \in X$. There exists a unique additive derivation mapping.

$A : X \rightarrow Y$ and a unique quadratic derivation mapping $Q : X \rightarrow Y$ satisfying the functional equation (11) and

$$\begin{aligned} \|f(x) - A(x) - Q(x)\| &\leq \frac{1}{2} \left[\frac{1}{2n} \sum_{k=\frac{1-j}{2}}^{\infty} \left(\frac{\alpha(n^{kj}x, 0, 0)}{n^{kj}} + \frac{\alpha(-n^{kj}x, 0, 0)}{n^{kj}} \right) \right] \\ &+ \frac{1}{4n^2} \left[\sum_{k=\frac{1-j}{2}}^{\infty} \left(\frac{\alpha(n^{kj}x, 0, 0)}{n^{2kj}} + \frac{\alpha(-n^{kj}x, 0, 0)}{n^{2kj}} \right) \right] \end{aligned} \quad (152)$$

For all $x \in X$. The mapping $A(x)$ and $Q(x)$ are defined in (135) and (146) respectively for all $x \in X$.

Proof: The proof follows by Theorems 6.1, using Theorems 10.2 and 11.2.

Using Corollaries 10.3 and 11.3 we have the following corollary concerning the stability of (11).

Corollary 12.2: Let $f : X \rightarrow Y$ be a mapping and there exists a real numbers λ and s such that

$$\|Df(x, y, z)\| \leq \begin{cases} \lambda; \\ \lambda \{ \|x\|^s + \|y\|^s + \|z\|^s \}; & s \neq 1, 2; \\ \lambda \{ \|x\|^s \|y\|^s \|z\|^s + \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \}; & s \neq \frac{1}{3}, \frac{2}{3}; \end{cases} \quad (153)$$

And (138), (149) for all $x, y, z \in X$. There exists a unique additive derivation $A : X \rightarrow Y$ and a unique quadratic derivation $Q : X \rightarrow Y$ such that

$$\|f(x) - A(x) - Q(x)\| \leq \begin{cases} \frac{\lambda}{2|n-1|} + \frac{\lambda}{4|n^2-1|}; \\ \frac{\lambda \|x\|^s}{2|n-n^s|} + \frac{\lambda \|x\|^s}{4|n^2-n^s|}; \\ \frac{\lambda \|x\|^{3s}}{2|n-n^{3s}|} + \frac{\lambda \|x\|^{3s}}{4|n^2-n^{3s}|}; \end{cases} \quad (154)$$

For all $x \in X$.

13. Stability results for (11): odd case-fixed point method

In this section, we give the generalized Ulam-Hyers stability of the functional equation (11), when f is odd case.

Theorem 13.1: Let $j = \pm 1$. Let $f_a : X \rightarrow Y$ be an odd mapping for which there exists functions $\alpha, \beta : X^3 \rightarrow [0, \infty)$ with the conditions

$$\sum_{k=0}^{\infty} \frac{\alpha(n^{kj}x, n^{kj}y, n^{kj}z)}{n^{kj}} \text{ Converges in } \mathbb{R} \text{ and } \lim_{k \rightarrow \infty} \frac{\alpha(n^{kj}x, n^{kj}y, n^{kj}z)}{n^{kj}} = 0 \quad (155)$$

$$\sum_{k=0}^{\infty} \frac{\beta(n^{kj}x, n^{kj}y, n^{kj}z)}{n^{3kj}} \text{ Converges in } \mathbb{R} \text{ and } \lim_{k \rightarrow \infty} \frac{\beta(n^{kj}x, n^{kj}y, n^{kj}z)}{n^{3kj}} = 0 \quad (156)$$

Where η is defined in (91) satisfying the functional inequalities

$$\|Df_a(x, y, z)\| \leq \alpha(x, y, z) \quad (157)$$

And

$$\|f_a(xyz) - f_a(x)yz - xf_a(y)z - xyf_a(z)\| \leq \beta(x, y, z) \quad (158)$$

For all $x, y, z \in X$. Then there exists $L = L(i) < 1$ such that the function

$$x \rightarrow \beta(x) = \frac{1}{2} \alpha\left(\frac{x}{n}, 0, 0\right)$$

As the property

$$\frac{1}{\eta_i} \beta(\eta_i x) = L \beta(x) \quad (159)$$

For all $x \in X$. Then there exists a unique Additive Derivation mapping $A : X \rightarrow Y$ satisfying the functional equation (11) and

$$\|f_a(x) - A(x)\| \leq \frac{L^{1-i}}{1-L} \beta(x) \quad (160)$$

For all $x \in X$.

Proof: It follows from Theorem 7.1 that A is a unique additive mapping and satisfies (11) for all $x \in X$. It follows from (155), (156) and (158) that

$$\begin{aligned} & \|A(xyz) - A(x)yz - xA(y)z - xyA(z)\| \\ & \leq \frac{1}{n^{3k}} \|f_a(n^k(xyz)) - f_a(n^kx)(n^ky n^kz) - n^kx f_a(n^ky) n^kz - n^kx n^ky f_a(n^kz)\| \\ & \leq \frac{1}{n^{3k}} \beta(n^kx, n^ky, n^kz) \rightarrow 0 \text{ as } k \rightarrow \infty. \end{aligned}$$

Thus the mapping $A : X \rightarrow Y$ is unique additive derivation mapping satisfying (11).

Corollary 13.2: Let $f_a : X \rightarrow Y$ be a odd mapping and there exists a real numbers λ and s such that,

$$\|Df_a(x, y, z)\| \leq \begin{cases} \lambda; & s \neq 1; \\ \lambda \left(\|x\|^s + \|y\|^s + \|z\|^s \right); & s \neq \frac{1}{3}; \\ \lambda \left(\|x\|^s \|y\|^s \|z\|^s + \left\{ \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \right\} \right); & s \neq \frac{1}{3}; \end{cases} \quad (161)$$

$$\|f_a(xyz) - f_a(x)yz - xf_a(y)z - xyf_a(z)\| \leq \begin{cases} \lambda; & s \neq 1; \\ \lambda(\|x\|^s + \|y\|^s + \|z\|^s); & \\ \lambda(\|x\|^s \|y\|^s \|z\|^s + \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s}); & s \neq \frac{1}{3}; \end{cases} \quad (162)$$

For all $x, y, z \in X$. Then there exists a unique additive derivation $A : X \rightarrow Y$ such that

$$\|f_a(x) - A(x)\| \leq \begin{cases} \frac{\lambda}{2|n-1|}, \\ \frac{\lambda\|x\|^s}{2|n-n^s|}, \\ \frac{\lambda\|x\|^{3s}}{2|n-n^{3s}|}, \end{cases} \quad (163)$$

For all $x \in X$.

14. Stability results for (11): even case-fixed point method

In this section, we discuss the generalized Ulam-Hyers stability of the functional equation (11), when f is even case.

Theorem 14.1: Let $j = \pm 1$. Let $f_a : X \rightarrow Y$ be a even mapping for which there exists functions $\alpha, \beta : X^3 \rightarrow [0, \infty)$ with the conditions

$$\sum_{k=0}^{\infty} \frac{\alpha(n^{kj}x, n^{kj}y, n^{kj}z)}{n^{2kj}} \text{ Converges in } \mathbb{R} \text{ and } \lim_{k \rightarrow \infty} \frac{\alpha(n^{kj}x, n^{kj}y, n^{kj}z)}{n^{2kj}} = 0 \quad (164)$$

$$\sum_{k=0}^{\infty} \frac{\beta(n^{kj}x, n^{kj}y, n^{kj}z)}{n^{6kj}} \text{ Converges in } \mathbb{R} \text{ and } \lim_{k \rightarrow \infty} \frac{\beta(n^{kj}x, n^{kj}y, n^{kj}z)}{n^{6kj}} = 0 \quad (165)$$

Where η_i is defined in (91) satisfying the functional inequalities

$$\|Df_Q(x, y, z)\| \leq \alpha(x, y, z) \quad (166)$$

And

$$\|f_Q(xyz) - f_Q(x)y^2z^2 - x^2f_Q(y)z^2 - x^2y^2f_Q(z)\| \leq \beta(x, y, z) \quad (167)$$

For all $x, y, z \in X$. Then there exists $L = L(i) < 1$ such that the function

$$x \rightarrow \beta(x) = \frac{1}{2} \alpha\left(\frac{x}{n}, 0, 0\right)$$

As the property

$$\frac{1}{\eta_i^2} \beta(\eta_i x) = L \beta(x) \quad (168)$$

For all $x \in X$. Then there exists a unique Quadratic Derivation mapping $Q : X \rightarrow Y$ satisfying the functional equation (11) and

$$\|f_Q(x) - Q(x)\| \leq \frac{L^{1-i}}{1-L} \beta(x) \quad (169)$$

For all $x \in X$.

Proof: It follows from Theorem 8.1 that Q is a unique quadratic mapping and satisfies (11) for all $x \in X$. It follows from (164), (165) and (167) that

$$\begin{aligned} & \|Q(xyz) - Q(x)y^2z^2 - x^2Q(y)z^2 - x^2y^2Q(z)\| \\ & \leq \frac{1}{n^{2k}} \|f_Q(n^k(xyz)) - f_Q(n^kx)n^{2k}y n^{2k}z - n^{2k}x f_Q(n^ky)n^{2k}z - n^{2k}x n^{2k}y f_Q(n^kz)\| \\ & \leq \frac{1}{n^{2k}} \beta(n^kx, n^ky, n^kz) \rightarrow 0 \text{ as } k \rightarrow \infty. \end{aligned}$$

Thus the mapping $Q : X \rightarrow Y$ is unique quadratic derivation mapping satisfying (11).

Corollary 14.2: Let $f_Q : X \rightarrow Y$ be a even mapping and there exists a real numbers λ and s such that,

$$\|Df_Q(x, y, z)\| \leq \begin{cases} \lambda; & s \neq 1; \\ \lambda(\|x\|^s + \|y\|^s + \|z\|^s); & s \neq \frac{1}{3}; \\ \lambda(\|x\|^s \|y\|^s \|z\|^s + \{ \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \}); & s \neq \frac{1}{3}; \end{cases} \quad (170)$$

$$\begin{aligned} & \|f_Q(xyz) - f_Q(x)y^2z^2 - x^2f_Q(y)z^2 - x^2y^2f_Q(z)\| \\ & \leq \begin{cases} \lambda; & s \neq 1; \\ \lambda(\|x\|^s + \|y\|^s + \|z\|^s); & s \neq \frac{1}{3}; \\ \lambda(\|x\|^s \|y\|^s \|z\|^s + \{ \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \}); & s \neq \frac{1}{3}; \end{cases} \quad (171) \end{aligned}$$

For all $x, y, z \in X$. Then there exists a unique quadratic derivation $Q : X \rightarrow Y$ such that

$$\|f_Q(x) - Q(x)\| \leq \begin{cases} \frac{\lambda}{4|n^2 - 1|}; \\ \frac{\lambda\|x\|^s}{4|n^2 - n^s|}; \\ \frac{\lambda\|x\|^{3s}}{4|n^2 - n^{3s}|}; \end{cases} \quad (172)$$

For all $x \in X$.

15. Stability results for (11): mixed case-fixed point method

In this section, we investigate the generalized Ulam-Hyers stability of the functional equation (11), when f is mixed case.

Theorem 15.1: Let $f : X \rightarrow Y$ be a mapping for which there exist a function $\alpha, \beta : X^3 \rightarrow [0, \infty)$ with the condition, (155), (156) and (164), (165) where η is defined in (91) such that the functional inequality

$$\|Df(x, y, z)\| \leq \alpha(x, y, z) \quad (173)$$

And (158), (167) for all $x, y, z \in X$. If there exists $L = L(i) < 1$ such that the function

$$x \rightarrow \beta(x) = \frac{1}{2} \alpha\left(\frac{x}{n}, 0, 0\right),$$

Has the properties (159) and (168) for all $x \in X$. Then there exists a unique additive derivation mapping $A : X \rightarrow Y$ and a unique quadratic derivation mapping $Q : X \rightarrow Y$ satisfying the functional equation (11) and

$$\|f(x) - A(x) - Q(x)\| \leq \frac{L^{1-i}}{1-L} [\beta(x) + \beta(-x)] \quad (174)$$

Holds for all $x \in X$.

Corollary 15.2: Let $f : X \rightarrow Y$ be a mapping and there exists a real numbers λ and s such that

$$\|Df(x, y, z)\| \leq \begin{cases} \lambda; \\ \lambda \{\|x\|^s + \|y\|^s + \|z\|^s\}; & s \neq 1, 2; \\ \lambda \{\|x\|^s \|y\|^s \|z\|^s + \{\|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s}\}\}; & s \neq \frac{1}{3}, \frac{2}{3}; \end{cases} \quad (175)$$

And (158), (167) for all $x, y, z \in X$. There exists a unique additive derivation mapping $A : X \rightarrow Y$ and a unique quadratic derivation mapping $Q : X \rightarrow Y$ such that

$$\|f(x) - A(x) - Q(x)\| \leq \begin{cases} \frac{\lambda}{2|n-1|} + \frac{\lambda}{4|n^2-1|} \\ \frac{\lambda\|x\|^s}{2|n-n^s|} + \frac{\lambda\|x\|^s}{4|n^2-n^s|}; \\ \frac{\lambda\|x\|^{3s}}{2|n-n^{3s}|} + \frac{\lambda\|x\|^{3s}}{4|n^2-n^{3s}|}; \end{cases} \quad (176)$$

For all $x \in X$.

16. Stability results for (11): another substitutions

In this section, the generalized Ulam-Hyers stability of generalized n-type additive quadratic functional equation (11) is investigated by using various substitutions. Hence the details of the proof are omitted.

Theorem 16.1: Let $j \in \{-1, 1\}$ and $\phi : X^3 \rightarrow [0, \infty)$ be a function satisfying the conditions

$$\sum_{k=0}^{\infty} \frac{\phi(n^{2kj} x, n^{2kj} y, n^{2kj} z)}{n^{2kj}} \text{ Converges to } \lim_{k \rightarrow \infty} \frac{\phi(n^{2kj} x, n^{2kj} y, n^{2kj} z)}{n^{2kj}} = 0 \quad (177)$$

$$\sum_{k=0}^{\infty} \frac{\phi(n^{2kj} x, n^{2kj} y, n^{2kj} z)}{n^{4kj}} \text{ Converges to } \lim_{k \rightarrow \infty} \frac{\phi(n^{2kj} x, n^{2kj} y, n^{2kj} z)}{n^{4kj}} = 0 \quad (178)$$

For all $x, y, z \in X$. Let $f : X \rightarrow Y$ be a function satisfying the inequality

$$\|Df(x, y, z)\| \leq \phi(x, y, z) \quad (179)$$

For all $x, y, z \in X$. Then there exists a unique additive mapping $A : X \rightarrow Y$ and a unique quadratic mapping $Q : X \rightarrow Y$ which satisfies the functional equation (11) and

$$\begin{aligned} \|f(y) - A(y) - Q(y)\| &\leq \frac{1}{2} \left[\frac{1}{2n^2} \sum_{k=\frac{1-j}{2}}^{\infty} \left(\frac{\phi(0, n^{2kj} y, 0)}{n^{2kj}} + \frac{\phi(0, n^{2kj} y, 0)}{n^{2kj}} \right) \right. \\ &\quad \left. + \frac{1}{4n^4} \sum_{k=\frac{1-j}{2}}^{\infty} \left(\frac{\phi(0, n^{2kj} y, 0)}{n^{4kj}} + \frac{\phi(0, n^{2kj} y, 0)}{n^{4kj}} \right) \right] \end{aligned} \quad (180)$$

For all $y \in X$. The mapping $A(y)$ and $Q(y)$ are defined by

$$A(y) = \lim_{k \rightarrow \infty} \frac{f_a(n^{2kj} y)}{n^{2kj}}, \quad Q(y) = \lim_{k \rightarrow \infty} \frac{f_a(n^{2kj} y)}{n^{4kj}} \quad (181)$$

Respectively for all $y \in X$.

Corollary 16.2: Let λ and s be a nonnegative real numbers. Let a function $f : X \rightarrow Y$ satisfying the inequality

$$\|Df(x, y, z)\| \leq \begin{cases} \lambda; & s \neq 1, 2; \\ \lambda \left(\|x\|^s + \|y\|^s + \|z\|^s \right); & s \neq \frac{1}{3}, \frac{2}{3}; \\ \lambda \left(\|x\|^s \|y\|^s \|z\|^s + \left(\|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \right) \right); & s \neq \frac{1}{3}, \frac{2}{3}; \end{cases} \quad (182)$$

For all $x, y, z \in X$. Then there exists a unique additive mapping $A : X \rightarrow Y$ and a unique quadratic mapping $Q : X \rightarrow Y$ such that

$$\|f(y) - A(y) - Q(y)\| \leq \begin{cases} \frac{\lambda}{2} \left[\frac{1}{|n^2-1|} + \frac{1}{|n^4-1|} \right], \\ \frac{\lambda \|y\|^s}{2} \left[\frac{1}{|n^2-n^{2s}|} + \frac{1}{2n|n^4-n^{2s}|} \right], \\ \frac{\lambda \|y\|^{3s}}{2} \left[\frac{1}{|n^2-n^{6s}|} + \frac{1}{2n|n^4-n^{6s}|} \right], \end{cases} \quad (183)$$

For all $y \in X$.

Theorem 16.3: Let $f: W \rightarrow B$ be a mapping for which there exists a function $\phi: W^3 \rightarrow [0, \infty)$ with the conditions

$$\lim_{k \rightarrow \infty} \frac{\phi(\tau_i^k x, \tau_i^k y, \tau_i^k z)}{\tau_i^k} = 0 \quad (184)$$

And

$$\lim_{k \rightarrow \infty} \frac{\phi(\tau_i^k x, \tau_i^k y, \tau_i^k z)}{\tau_i^{2k}} = 0 \quad (185)$$

Where

$$\tau_i = \begin{cases} n^2, & i = 0; \\ \frac{1}{n^2}, & i = 1; \end{cases} \quad (186)$$

Such that the functional inequality with

$$\|Df(x, y, z)\| \leq \phi(x, y, z) \quad (187)$$

For all $x, y, z \in W$. If there exist $L = L(i)$ such that the function for all $y \in W$, such that the function

$$y \rightarrow \delta(y) = \frac{1}{2} \phi\left(0, \frac{y}{n^2}, 0\right) \quad (188)$$

Has the properties

$$\frac{1}{\tau_i} \delta(\tau_i y) = L \delta(y), \quad \frac{1}{\tau_i} \delta\left(\tau_i y\right) = L \delta(y)$$

For all $y \in W$. Then there exists a unique additive function $A: W \rightarrow B$ and a unique quadratic function $Q: W \rightarrow B$ satisfying the functional equation (11) and

$$\|f(y) - A(y) - Q(y)\| \leq \frac{L^{1-i}}{1-L} [\delta(y) + \delta(-y)] \quad (189)$$

Holds for all $y \in W$.

Corollary 16.4: Let $f: W \rightarrow B$ be a mapping and there exists a real numbers λ and s such that,

$$\|Df(x, y, z)\| \leq \begin{cases} \lambda, & s \neq 1, 2 \\ \lambda \left\{ \|x\|^s + \|y\|^s + \|z\|^s \right\}, & s \neq \frac{1}{3}, \frac{2}{3} \\ \lambda \left\{ \|x\|^s \|y\|^s \|z\|^s + \left\{ \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \right\} \right\}, & s \neq \frac{1}{3}, \frac{2}{3} \end{cases} \quad (190)$$

For all $x, y, z \in W$. There exists a unique additive mapping $A: W \rightarrow B$ and a unique quadratic mapping $Q: W \rightarrow B$ such that,

$$\|f(y) - A(y) - Q(y)\| \leq \begin{cases} \frac{\lambda}{2} \left[\frac{1}{|n^2 - 1|} + \frac{1}{|n^4 - 1|} \right], \\ \frac{\lambda \|y\|^s}{2} \left[\frac{1}{|n^2 - n^{2s}|} + \frac{1}{2n |n^4 - n^{2s}|} \right], \\ \frac{\lambda \|y\|^{3s}}{2} \left[\frac{1}{|n^2 - n^{6s}|} + \frac{1}{2n |n^4 - n^{6s}|} \right], \end{cases} \quad (191)$$

For all $y \in W$.

Theorem 16.5: Let $j = \pm 1$. Let $f: X \rightarrow Y$ be a mapping for which there exists functions $\phi, \varphi: X^3 \rightarrow [0, \infty)$, satisfying the conditions (177), (178) and

$$\sum_{k=0}^{\infty} \frac{\varphi(n^{2kj}x, n^{2kj}y, n^{2kj}z)}{n^{6kj}} \text{ Converges in } \mathbb{R} \text{ and } \lim_{k \rightarrow \infty} \frac{\varphi(n^{2kj}x, n^{2kj}y, n^{2kj}z)}{n^{6kj}} = 0 \quad (192)$$

$$\sum_{k=0}^{\infty} \frac{\varphi(n^{2kj}x, n^{2kj}y, n^{2kj}z)}{n^{8kj}} \text{ Converges in } \mathbb{R} \text{ and } \lim_{k \rightarrow \infty} \frac{\varphi(n^{2kj}x, n^{2kj}y, n^{2kj}z)}{n^{8kj}} = 0 \quad (193)$$

Such that the functional inequality

$$\|Df(x, y, z)\| \leq \phi(x, y, z) \quad (194)$$

And

$$\|f(xyz) - f(x)yz - xf(y)z - xyf(z)\| \leq \varphi(x, y, z) \quad (195)$$

For all $x, y, z \in X$. There exists a unique additive derivation mapping $A: X \rightarrow Y$ and a unique quadratic derivation mapping $Q: X \rightarrow Y$ satisfying the functional equation (11) and

$$\|f(y) - A(y) - Q(y)\| \leq \frac{1}{2} \left[\frac{1}{2n^2} \sum_{k=\frac{1-j}{2}}^{\infty} \left(\frac{\phi(0, n^{2kj}y, 0)}{n^{2kj}} + \frac{\phi(0, n^{2kj}y, 0)}{n^{2kj}} \right) \right]$$

$$+\frac{1}{4n^4} \sum_{k=\frac{1-j}{2}}^{\infty} \left(\frac{\phi(0, n^{2kj} y, 0)}{n^{4kj}} + \frac{\phi(0, n^{2kj} y, 0)}{n^{4kj}} \right) \quad (196)$$

For all $y \in X$. The mapping $A(y)$ and $Q(y)$ are respectively in (181) for all $y \in Y$.

Corollary 16.6: Let $f : X \rightarrow Y$ be a mapping and there exists a real numbers λ and s such that

$$\|Df(x, y, z)\| \leq \begin{cases} \lambda; & s \neq 1, 2 \\ \lambda \left\{ \|x\|^s + \|y\|^s + \|z\|^s \right\}; & s \neq \frac{1}{3}, \frac{2}{3} \\ \lambda \left\{ \|x\|^s \|y\|^s \|z\|^s + \left\{ \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \right\} \right\}; & s \neq \frac{1}{3}, \frac{2}{3} \end{cases} \quad (197)$$

For all $x, y, z \in X$. There exists a unique additive derivation $A : X \rightarrow Y$ and a unique quadratic derivation $Q : X \rightarrow Y$ such that

$$\|f(y) - A(y) - Q(y)\| \leq \begin{cases} \frac{\lambda}{2} \left[\frac{1}{|n^2 - 1|} + \frac{1}{|n^4 - 1|} \right], \\ \frac{\lambda \|y\|^s}{2} \left[\frac{1}{|n^2 - n^{2s}|} + \frac{1}{2n |n^4 - n^{2s}|} \right], \\ \frac{\lambda \|y\|^{3s}}{2} \left[\frac{1}{|n^2 - n^{6s}|} + \frac{1}{2n |n^4 - n^{6s}|} \right], \end{cases} \quad (198)$$

For all $y \in X$.

Theorem 16.7: Let $j \in \{-1, 1\}$ and $\psi : X^3 \rightarrow [0, \infty)$ be a function satisfying the conditions

$$\sum_{k=0}^{\infty} \frac{\psi(n^{3kj} x, n^{3kj} y, n^{3kj} z)}{n^{3kj}} \text{ Converges to } \mathbb{R} \quad \lim_{k \rightarrow \infty} \frac{\psi(n^{3kj} x, n^{3kj} y, n^{3kj} z)}{n^{3kj}} = 0 \quad (199)$$

$$\sum_{k=0}^{\infty} \frac{\psi(n^{3kj} x, n^{3kj} y, n^{3kj} z)}{n^{6kj}} \text{ Converges to } \mathbb{R} \quad \lim_{k \rightarrow \infty} \frac{\psi(n^{3kj} x, n^{3kj} y, n^{3kj} z)}{n^{6kj}} = 0 \quad (200)$$

For all $x, y, z \in X$. Let $f : X \rightarrow Y$ be a function satisfying the inequality

$$\|Df(x, y, z)\| \leq \psi(x, y, z) \quad (201)$$

For all $x, y, z \in X$. Then there exists a unique additive mapping $A : X \rightarrow Y$ and a unique quadratic mapping $Q : X \rightarrow Y$ which satisfies the functional equation (11) and

$$\|f(z) - A(z) - Q(z)\| \leq \frac{1}{2} \left[\frac{1}{2n^3} \sum_{k=\frac{1-j}{2}}^{\infty} \left(\frac{\psi(0,0,n^{3kj}z)}{n^{3kj}} + \frac{\psi(0,0,n^{3kj}z)}{n^{3kj}} \right) \right. \\ \left. + \frac{1}{4n^6} \sum_{k=\frac{1-j}{2}}^{\infty} \left(\frac{\psi(0,0,n^{3kj}z)}{n^{6kj}} + \frac{\psi(0,0,n^{2kj}z)}{n^{6kj}} \right) \right] \quad (202)$$

For all $z \in X$. The mapping $A(z)$ and $Q(z)$ are defined by

$$A(z) = \lim_{k \rightarrow \infty} \frac{f_a(n^{3kj}z)}{n^{3kj}}, \quad Q(z) = \lim_{k \rightarrow \infty} \frac{f_q(n^{3kj}z)}{n^{6kj}} \quad (203)$$

Respectively for all $z \in X$.

Corollary 16.8: Let λ and s be a nonnegative real numbers. Let a function $f : X \rightarrow Y$ satisfying the inequality

$$\|Df(x, y, z)\| \leq \begin{cases} \lambda; & s \neq 1, 2; \\ \lambda(\|x\|^s + \|y\|^s + \|z\|^s); & s \neq \frac{1}{3}, \frac{2}{3}; \\ \lambda(\|x\|^s \|y\|^s \|z\|^s + \{ \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \}); & s \neq \frac{1}{3}, \frac{2}{3}; \end{cases} \quad (204)$$

For all $x, y, z \in X$. Then there exists a unique additive mapping $A : X \rightarrow Y$ and a unique quadratic mapping $Q : X \rightarrow Y$ such that

$$\|f(z) - A(z) - Q(z)\| \leq \begin{cases} \frac{\lambda}{2} \left[\frac{1}{|n^3 - 1|} + \frac{1}{|n^6 - 1|} \right], \\ \frac{\lambda \|z\|^s}{2} \left[\frac{1}{|n^3 - n^{3s}|} + \frac{1}{2n |n^6 - n^{3s}|} \right], \\ \frac{\lambda \|z\|^{3s}}{2} \left[\frac{1}{|n^3 - n^{6s}|} + \frac{1}{2n |n^6 - n^{9s}|} \right], \end{cases} \quad (205)$$

For all $z \in X$.

Theorem 16.9: Let $f : W \rightarrow B$ be a mapping for which there exists a function $\psi : W^3 \rightarrow [0, \infty)$ with the conditions

$$\lim_{k \rightarrow \infty} \frac{\psi(\kappa_i^k x, \kappa_i^k y, \kappa_i^k z)}{\kappa_i^k} = 0 \quad (206)$$

And

$$\lim_{k \rightarrow \infty} \frac{\phi(\kappa_i^k x, \kappa_i^k y, \kappa_i^k z)}{\kappa_i^{2k}} = 0 \quad (207)$$

Where

$$\kappa_i = \begin{cases} n^3, & i = 0; \\ \frac{1}{n^3}, & i = 1; \end{cases} \quad (208)$$

Such that the functional inequality with

$$\|Df(x, y, z)\| \leq \psi(x, y, z) \quad (209)$$

For all $x, y, z \in W$. If there exist $L = L(i)$ such that the function for all $y \in W$, such that the function

$$z \rightarrow \xi(z) = \frac{1}{2} \psi\left(0, 0, \frac{z}{n^3}\right), \quad (210)$$

Has the properties

$$\frac{1}{\kappa_i} \xi(\kappa_i z) = L \xi(z), \quad \frac{1}{\kappa_i} \xi(\kappa_i z) = L \xi(z)$$

For all $z \in W$. Then there exists a unique additive function $A: W \rightarrow B$ and a unique quadratic function $Q: W \rightarrow B$ satisfying the functional equation (11) and

$$\|f(z) - A(z) - Q(z)\| \leq \frac{L^{1-i}}{1-L} [\xi(z) + \xi(-z)] \quad (211)$$

Holds for all $z \in W$.

Corollary 16.10: Let $f: W \rightarrow B$ be a mapping and there exists a real numbers λ and s such that,

$$\|Df(x, y, z)\| \leq \begin{cases} \lambda, & \\ \lambda \left\{ \|x\|^s + \|y\|^s + \|z\|^s \right\}, & s \neq 1, 2 \\ \lambda \left\{ \|x\|^s \|y\|^s \|z\|^s + \left\{ \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \right\} \right\}, & s \neq \frac{1}{3}, \frac{2}{3} \end{cases} \quad (212)$$

For all $x, y, z \in W$. There exists a unique additive mapping $A: W \rightarrow B$ and a unique quadratic mapping $Q: W \rightarrow B$ such that,

$$\|f(z) - A(z) - Q(z)\| \leq \begin{cases} \frac{\lambda}{2} \left[\frac{1}{|n^3 - 1|} + \frac{1}{|n^6 - 1|} \right], \\ \frac{\lambda \|z\|^s}{2} \left[\frac{1}{|n^3 - n^{3s}|} + \frac{1}{2n |n^6 - n^{3s}|} \right], \\ \frac{\lambda \|z\|^{3s}}{2} \left[\frac{1}{|n^3 - n^{6s}|} + \frac{1}{2n |n^6 - n^{9s}|} \right], \end{cases} \quad (213)$$

For all $z \in W$.

Theorem 16.11: Let $j = \pm 1$. Let $f: X \rightarrow Y$ be a mapping for which there exists functions $\psi, \gamma: X^3 \rightarrow [0, \infty)$, satisfying the conditions (199), (200) and

$$\sum_{k=0}^{\infty} \frac{\gamma(n^{3kj} x, n^{3kj} y, n^{3kj} z)}{n^{9kj}} \text{ converges in } R \text{ and } \lim_{k \rightarrow \infty} \frac{\gamma(n^{3kj} x, n^{3kj} y, n^{3kj} z)}{n^{9kj}} = 0 \quad (214)$$

$$\sum_{k=0}^{\infty} \frac{\gamma(n^{3kj} x, n^{3kj} y, n^{3kj} z)}{n^{12kj}} \text{ converges in } R \text{ and } \lim_{k \rightarrow \infty} \frac{\gamma(n^{3kj} x, n^{3kj} y, n^{3kj} z)}{n^{12kj}} = 0 \quad (215)$$

Such that the functional inequality

$$\|Df(x, y, z)\| \leq \psi(x, y, z) \quad (216)$$

And

$$\|f(xyz) - f(x)yz - xf(y)z - xyf(z)\| \leq \gamma(x, y, z) \quad (217)$$

For all $x, y, z \in X$. There exists a unique additive derivation mapping $A: X \rightarrow Y$ and a unique quadratic derivation mapping $Q: X \rightarrow Y$ satisfying the functional equation (11) and

$$\begin{aligned} \|f(z) - A(z) - Q(z)\| &\leq \frac{1}{2} \left[\frac{1}{2n^3} \sum_{k=\frac{1-j}{2}}^{\infty} \left(\frac{\psi(0, 0, n^{3kj} z)}{n^{3kj}} + \frac{\psi(0, 0, n^{3kj} z)}{n^{3kj}} \right) \right. \\ &\quad \left. + \frac{1}{4n^6} \sum_{k=\frac{1-j}{2}}^{\infty} \left(\frac{\psi(0, 0, n^{3kj} z)}{n^{6kj}} + \frac{\psi(0, 0, n^{2kj} z)}{n^{6kj}} \right) \right] \end{aligned} \quad (218)$$

For all $z \in X$. The mapping $A(z)$ and $Q(z)$ are defined in (198) respectively for all $z \in X$.

Corollary 16.12: Let $f: X \rightarrow Y$ be a mapping and there exists a real numbers λ and s such that

$$\|Df(x, y, z)\| \leq \begin{cases} \lambda; & s \neq 1, 2 \\ \lambda \left\{ \|x\|^s + \|y\|^s + \|z\|^s \right\}; & s \neq \frac{1}{3}, \frac{2}{3} \\ \lambda \left\{ \|x\|^s \|y\|^s \|z\|^s + \left\{ \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \right\} \right\}; & s \neq \frac{1}{3}, \frac{2}{3} \end{cases} \quad (219)$$

For all $x, y, z \in X$. There exists a unique additive derivation $A: X \rightarrow Y$ and a unique quadratic derivation $Q: X \rightarrow Y$ such that

$$\|f(z) - A(z) - Q(z)\| \leq \begin{cases} \frac{\lambda}{2} \left[\frac{1}{|n^3 - 1|} + \frac{1}{|n^6 - 1|} \right], \\ \frac{\lambda \|z\|^s}{2} \left[\frac{1}{|n^3 - n^{3s}|} + \frac{1}{2n |n^6 - n^{3s}|} \right], \\ \frac{\lambda \|z\|^{3s}}{2} \left[\frac{1}{|n^3 - n^{6s}|} + \frac{1}{2n |n^6 - n^{9s}|} \right], \end{cases} \quad (220)$$

For all $z \in X$.

References

- [1] J. Aczel and J. Dhombres, *Functional Equations in Several Variables*, Cambridge University Press, 1989. <http://dx.doi.org/10.1017/CBO9781139086578>.
- [2] T. Aoki, on the stability of the linear transformation in Banach spaces, *J. Math. Soc. Japan*, 2 (1950), 64-66. <http://dx.doi.org/10.2969/jmsj/00210064>.
- [3] M. Arunkumar, S. Jayanthi, S. Hema Latha, Stability of quadratic derivations of Arun- quadratic functional equation, *International Journal Mathematical Sciences and Engineering Applications*, Vol. 5 No. V, Sept. (2011), 433-443.
- [4] M. Arunkumar, S. Karthikeyan, Solution and stability of n -dimensional mixed Type additive and quadratic functional equation, *Far East Journal of Applied Mathematics*, Volume 54, Number 1, (2011), 47-64.
- [5] M. Arunkumar, John M. Rassias, On the generalized Ulam-Hyers stability of an AQ- mixed type functional equation with counter examples, *Far East Journal of Applied Mathematics*, Volume 71, No. 2, (2012), 279-305.
- [6] M. Arunkumar, Solution and stability of modified additive and quadratic functional equation in generalized 2-normed spaces, *International Journal Mathematical Sciences and Engineering Applications*, Vol. 7 No. I (January, 2013), 383-391.
- [7] M. Arunkumar, Generalized Ulam - Hyers stability of derivations of a AQ - functional equation, "Cubo A Mathematical Journal" dedicated to Professor Gaston M.N'Guérékata on the occasion of his 60th Birthday Vol.15, No 01, (2013), 159-169.
- [8] M. Arunkumar, P. Agilan, Additive Quadratic functional equation are Stable in Banach space: A Direct Method, *Far East Journal of Applied Mathematics*, Volume 80, No. 1, (2013), 105 - 121.
- [9] M. Arunkumar, G.Shobana, S. Hemalatha, Ulam - Hyers, Ulam - Trassias, Ulam-Grassias, Ulam - Jrassias Stabilities of A Additive - Quadratic Mixed Type Functional Equation In Banach Spaces, *Proceedings of International Conference on Mathematics and Computing, ICMCE - 2013*, 604 - 611, ISBN 978-93-82338-89-5.
- [10] M. Arunkumar, P. Agilan, C. Devi Shyamala Mary, Permanence of A Generalized AQ Functional Equation In Quasi-Beta Normed Spaces, *International Journal of pure and applied Mathematics(Accepted)*.
- [11] M. Arunkumar, Perturbation of n Dimensional AQ - mixed type Functional Equation via Banach Spaces and Banach Algebra: Hyers Direct and Alternative Fixed Point Methods, *International Journal of Advanced Mathematical Sciences (IJAMS)*, Vol. 2 (1), (2014), 34-56.
- [12] M. Arunkumar, P. Agilan, Random stability of a additive quadratic Functional equation, *Proceedings of International Conference on Applied Mathematical Models, ICAMMA 2014*, 271 - 278.
- [13] M. Arunkumar, P. Agilan, C. Devi Shyamala Mary, Permanence of A Generalized AQ Functional Equation In Quasi-Beta Normed Spaces, A Fixed Point Approach, *Proceedings of the International Conference on Mathematical Methods and Computation, Jamal Academic Research Journal an Interdisciplinary*, (February 2014), 315-324.
- [14] C. Baak, D. Boo, Th.M. Rassias, Generalized additive mapping in Banach modules and isomorphism between C^* -algebras, *J. Math. Anal. Appl.* 314, (2006), 150-161. <http://dx.doi.org/10.1016/j.jmaa.2005.03.099>.
- [15] R. Badora, on approximate derivations, *Math. Inequal. Appl.* 9 (2006), no.1, 167-173.
- [16] C.Borelli, G.L.Forti, on a general Hyers-Ulam stability, *Internat J.Math.Math.Sci.* 18 (1995), 229-236. <http://dx.doi.org/10.1155/S0161171295000287>.
- [17] L. Brown and G. Pedersen, C^* -algebras of real rank zero, *J. Funct. Analysis*, 99, (1991), 138-149. [http://dx.doi.org/10.1016/0022-1236\(91\)90056-B](http://dx.doi.org/10.1016/0022-1236(91)90056-B).
- [18] I.S. Chang, E.H. Lee, H.M. Kim, On the Hyers-Ulam-Rassias stability of quadratic functional equations, *Math. Ineq. Appl.*, 6(1) (2003), 87-95.
- [19] P.W.Cholewa, Remarks on the stability of functional equations, *Aequationes Math.*, 27 (1984), 76-86. <http://dx.doi.org/10.1007/BF02192660>.
- [20] S.Czerwik, on the stability of the quadratic mappings in normed spaces, *Abh.Math.Sem.Univ Hamburg.*, 62 (1992), 59-64. <http://dx.doi.org/10.1007/BF02941618>.
- [21] S. Czerwik, *Functional Equations and Inequalities in Several Variables*, World Scientific, River Edge, NJ, 2002.
- [22] M. Eshaghi Gordji, H. Khodaie, Solution and stability of generalized mixed type cubic, quadratic and additive functional equation in Quasi-Banach spaces, arxiv: 0812. 2939v1 *Math FA*, 15 Dec 2008.
- [23] M. Eshaghi Gordji, H. Khodaei, J.M. Rassias, Fixed point methods for the stability of general quadratic functional equation, *Fixed Point Theory*, 12 (2011), no. 1, 71-82.
- [24] P. Gavruta, A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings, *J. Math. Anal. Appl.*, 184 (1994), 431-436. <http://dx.doi.org/10.1006/jmaa.1994.1211>.
- [25] M.E. Gordji, J.M. Rassias, N. Ghobadipour, Generalized Hyers-Ulam stability of the n -derivations, *Abs. Appl. Anal.* Volume 2009, Article ID 437931.
- [26] M.E. Gordji, S. Kaboli Gharetapeh, M.B. Savadkouhi, M.Aghaei, T.Karimi, On Cubic Derivations, *Int. Journal of Math. Analysis*, Vol.4, 2010, no.51, 2501-2514.
- [27] D. H. Hyers, On the stability of the linear functional equation, *Proc. Natl. Acad. Sci., U.S.A.*, 27 (1941), 222-224. <http://dx.doi.org/10.1073/pnas.27.4.222>.
- [28] D. H. Hyers, G. Isac and Th. M. Rassias, *Stability of functional equations in several variables*, Birkhauser, Basel, 1998. <http://dx.doi.org/10.1007/978-1-4612-1790-9>.
- [29] K.W. Jun and D.W. Park, Almost derivations on the Banach algebra $C_n[0,1]$, *Bull. Korean Math.Soc.* vol 33, No.3 (1996), 359-366.
- [30] K.W. Jun, H.M. Kim, On the Hyers-Ulam-Rassias stability of a generalized quadratic and additive type functional equation, *Bull. Korean Math. Soc.* 42 (1) (2005), 133-148. <http://dx.doi.org/10.4134/BKMS.2005.42.1.133>.
- [31] K.W. Jun and H.M. Kim, on the stability of an n -dimensional quadratic and additive type functional equation, *Math. Ineq. Appl.* 9 (1) (2006), 153-165.
- [32] S.M. Jung, On the Hyers-Ulam stability of the functional equations that have the quadratic property, *J. Math. Anal. Appl.* 222 (1998), 126-137. <http://dx.doi.org/10.1006/jmaa.1998.5916>.
- [33] S.M. Jung, *Hyers-Ulam-Rassias Stability of Functional Equations in Mathematical Analysis*, Hadronic Press, Plan Harbor, 2001.
- [34] Y.S. Jung, The Hyers-Ulam-Rassias Stability of module left derivations, *J. Math. Anal. Appl.*, doi: 10.1016/j.jmaa.2007.07.003, 1-9. <http://dx.doi.org/10.1016/j.jmaa.2007.07.003>.
- [35] Pl. Kannappan, Quadratic functional equation and inner product spaces, *Results Math.*, 27 (1995), 368-372. <http://dx.doi.org/10.1007/BF03322841>.
- [36] Pl. Kannappan, *Functional Equations and Inequalities with Applications*, Springer Monographs in Mathematics, 2009. <http://dx.doi.org/10.1007/978-0-387-89492-8>.
- [37] B.Margoils, J.B.Diaz, A fixed point theorem of the alternative for contractions on a generalized complete metric space, *Bull.Amer. Math. Soc.*, (1968), 305-309.
- [38] Matina J. Rassias, M. Arunkumar, S. Ramamoorthi, Stability of the Leibniz additive-quadratic functional equation in Quasi-Beta normed space: Direct and fixed point methods, *Journal of Concrete and Applicable Mathematics (JCAAM)*, Vol. 14 No. 1-2, (2014), 22 - 46.

- [39] S. Murthy, M. Arunkumar, G. Ganapathy, P. Rajarethinam, Stability of mixed type additive quadratic functional equation in Random Normed space, *International Journal of Applied Mathematics (IJAM)*, Vol. 26. No. 2 (2013), 123-136. <http://dx.doi.org/10.12732/ijam.v26i2.3>.
- [40] A. Najati and M.B. Moghimi, On the Stability of a quadratic and additive functional equation, *J. Math. Anal. Appl.*, 337 (2008), 399-415. <http://dx.doi.org/10.1016/j.jmaa.2007.03.104>.
- [41] C. Park, on the stability of the linear mapping in Banach modules, *J. Math. Anal. Appl.*, 275 (2002), 711-720. [http://dx.doi.org/10.1016/S0022-247X\(02\)00386-4](http://dx.doi.org/10.1016/S0022-247X(02)00386-4).
- [42] C. Park, Linear functional equation in Banach modules over a C*-algebra, *Acta. Appl. Math.* 77 (2003), 125-161. <http://dx.doi.org/10.1023/A:1024014026789>.
- [43] C. Park, Linear derivations on Banach algebras, *Funct. Anal. Appl.* 9 (2004), no.3, 359-368.
- [44] C. Park, Fixed points and Hyers-Ulam-Rassias stability of Cauchy-Jensen functional equation in Banach algebras, *Fixed Point Theory and Applications*, 2007, Art ID 50175.
- [45] C. Park and J.Hou, Homomorphism and derivations in C*-algebras, *Abstract Appl. Anal.* 2007, Art. Id 80630.
- [46] J.M. Rassias, on approximately of approximately linear mappings by linear mappings, *J. Funct. Anal. USA*, 46, (1982) 126-130. [http://dx.doi.org/10.1016/0022-1236\(82\)90048-9](http://dx.doi.org/10.1016/0022-1236(82)90048-9).
- [47] J.M. Rassias, H.M. Kim, Generalized Hyers-Ulam stability for general additive functional equations in quasi- -normed spaces, *J. Math. Anal. Appl.*, 356 (2009), no. 1, 302-309.
- [48] Th. M. Rassias, on the stability of the linear mapping in Banach spaces, *Proc. Amer. Math. Soc.*, 72 (1978), 297-300. <http://dx.doi.org/10.1090/S0002-9939-1978-0507327-1>.
- [49] Th.M.Rassias, on the stability of the functional equations in Banach spaces, *J. Math. Anal. Appl.*, 251, (2000), 264-284. <http://dx.doi.org/10.1006/jmaa.2000.7046>.
- [50] Th. M. Rassias, *Functional Equations, Inequalities and Applications*, Kluwer Academic Publishers, Dordrecht, Boston London, 2003. <http://dx.doi.org/10.1007/978-94-017-0225-6>.
- [51] K. Ravi, M.Arunkumar and J.M. Rassias, On the Ulam stability for the orthogonally general Euler-Lagrange type functional equation, *International Journal of Mathematical Sciences*, 2008 Vol.3, No. 08, 36-47.
- [52] S. M. Ulam, *Problems in Modern Mathematics*, Science Editions, Wiley, New York, 1964.
- [53] G.Zamani Eskandani, Stability of a mixed additive and quadratic functional equation in non-Archimedean Banach modules, *Taiwanese journal of mathematics*, Vol.14, No.4, pp. 1309-1324 August 2010.