International Journal of Advanced Statistics and Probability, 1 (3) (2013) 132-141
©Science Publishing Corporation
www.sciencepubco.com/index.php/IJASP

On a probability distribution with fractional moments arising from generalized Pearson system of differential equation and its characterization

M. Ahsanullah, M. Shakil*, B.M. Golam Kibria

Department of Management Sciences, Rider University, Lawrenceville, NJ 08648, USA
Department of Mathematics, Miami Dade College, Hialeah Campus, Hialeah, FL 33012, USA
Department of Mathematics and Statistics, Florida International University, University Park, Miami, FL 33199, USA
**Corresponding Author's E-mail: mshakil@mdc.edu*

Abstract

This paper derives a probability distribution with fractional moments arising from generalized Pearson system of differential equation. The expressions for the probability density function, cumulative distribution function and moments have been obtained. The plots for the probability density function, cumulative distribution function, and survival and hazard functions are also provided. Some distributional relationships of the proposed distribution have been established. A characterization of the new distribution is given. It is hoped that the findings of the paper will be useful for researchers in different fields like economics, engineering, environmental science, finance, medical sciences and physical sciences among others, where fractional moments become required to be computed if the integer moments $k \geq 1$ do not exist.

Keywords: *Characterization; Distribution; Differential Equations; Moments; Hazard Function; Probability Density Function.*

2010 Mathematics Subject Classifications: *33C15; 60E05; 62E15.*

1 Introduction

Various systems of distributions have been constructed to provide approximations to a wide variety of distributions, see, e.g., Johnson et al. [12]. One of these systems is the Pearson system. A continuous distribution belongs to this system if its probability density function (*pdf*) *f(x)* satisfies a differential equation of the form

$$\frac{1}{f_X(x)}\frac{d\,f(x)}{d\,x} = -\frac{x+a}{bx^2+cx+d}\quad , \tag{1}$$

where a, b, c, and d are real parameters such that *f(x)* is a *pdf*. The shapes of the *pdf*. depend on the values of these parameters based on which Pearson [18-19] classified these distributions into a number of types known as Pearson Types I – VI. Later in another paper, Pearson [20] defined more special cases and subtypes, known as Pearson Types VII - XII. Many well-known distributions are special cases of Pearson Type distributions which include Normal and Student t distributions (Pearson Type VII), Beta distribution (Pearson Type I), Gamma distribution (Pearson Type III), among others.

1.1 Generalized Pearson Distribution

In recent years, some researchers have considered a generalization of (1), known as generalized Pearson differential equation (GPE), given by

$$\frac{1}{f_X(x)}\frac{d\,f(x)}{d\,x} = \frac{\sum_{j=0}^{m} a_j\, x^j}{\sum_{j=0}^{n} b_j\, x^j}\quad , \tag{2}$$

where m, $n \in \mathrm{N}/\{0\}$ and the coefficients a_j and b_j are real parameters. The system of continuous univariate *pdf's* generated by GPE is called a generalized Pearson system, which includes a vast majority of continuous *pdf* '*s*, by proper choices of these parameters. For example:

1) Roy [22] studied GPE, when $m = 2, n = 3, b_0 = 0$, to derive five frequency curves whose parameters depends on the first seven population moments.
2) Dunning and Hanson [8] used GPE in his paper on generalized Pearson distributions and nonlinear programming.
3) Cobb et al. [5] extended Pearson's class of distributions to generate multimodal distributions by taking the polynomial in the numerator of GPE of degree higher than one, and the denominator, say, $v(x)$, having one of the following forms:

(I) $v(x) = 1, \quad -\infty < x < \infty$,

(II) $v(x) = x, \quad 0 < x < \infty$,

(III) $v(x) = x^2, \quad 0 < x < \infty$, (IV) $v(x) = x(1-x), \quad 0 < x < 1$.

4) Chaudhry and Ahmad [3] studied another class of generalized Pearson's distributions when

$$m = 4, n = 3, b_0 = b_1 = b_2 = 0, \frac{a_4}{2b_3} = -2\alpha, \frac{a_0}{2b_3} = 2\beta, b_3 \neq 0 .$$

5) Rossani and Scarfone [21] have studied the GPE in the following form

$$\frac{1}{f_X(x)} \frac{d f(x)}{d x} = -\frac{a_0 + a_1 x + a_2 x^2}{b_0 + b_1 x + b_2 x^2},$$

and used it to generate generalized Pearson distributions in order to study charged particles interacting with an electric and/or a magnetic field.
6) Shakil and Kibria [23], and Shakil et al. [24-25] have defined some new classes of generalized Pearson distributions for different choices of the parameters m , $n \in \mathrm{N} / \{0\}$ and the coefficients a_j and b_j in GPE. Recently, these distributions have been characterized by Hamedani [10-11]. Also, see Ahsanullah et al. [2], Shakil et al. [26], Kibria and Shakil [13], Lahcene [14], and Lee et al. [16], among others.

The organization of the paper is as follows. We give a new class of distribution as a solution of the GPE in section 2. Distributional properties of this is documented in section 3. Characterization of the distribution is provided in section 4. This paper ends up with some conclusions in section 5.

2 Derivation of new probability model

We give a new class of distributions as solutions of the GPE. We consider the following differential equation

$$\frac{df_X(x)}{dx} = \left(\frac{a_1 + a_2 x + a_3 x^2}{b_3 x^2 + b_4 x^3} \right) f_X(x), \tag{3}$$

which is a special case of the generalized Pearson Eq. (2) when $m = 2, n = 3$, Putting $b_3 = 1$, $b_4 = \gamma$, $a_1 = \beta\gamma$, $a_2 = \beta - \gamma + \gamma\nu$, $a_3 = \nu + \mu - 2$, $x > 0$; in (3), we have

$$\frac{1}{f_X(x)} \frac{df_X(x)}{dx} = \frac{\beta\gamma + (\beta - \gamma + \gamma\nu)x + (\mu + \nu - 2)x^2}{x^3 + \gamma x^2} = \frac{\nu - 1}{x} + \frac{\mu - 1}{x + \gamma} + \frac{\beta}{x^2},$$

where we assume that $\beta > 0$, $\gamma > 0$, $0 < \nu < 1$, $0 < \mu < 1$, $1 - \mu > \nu > 0$.
Integrating the above equation, we have

$$f_X(x) = C\, x^{\nu - 1} (x + \gamma)^{(\mu - 1)} \exp\left(-\beta x^{-1}\right), \quad 0 < x < \infty, \tag{4}$$

Using the equation (3.471.7), Page 340 of Gradshteyn and Ryzhik [9], we easily obtain the following normalizing constant as

$$\frac{1}{C} = \beta^{(\nu - 1)/2} \gamma^{\frac{\nu - 1}{2} + \mu} \Gamma(1 - \mu - \nu) \exp\left(\frac{\beta}{2\gamma}\right) W_{\frac{\nu - 1}{2} + \mu, \frac{-\nu}{2}}\left(\frac{\beta}{\gamma}\right), \tag{5}$$

where W(.) denotes the Whittaker function which is defined as the solution of the following differential equation

$$\frac{d^2 W}{dx^2} + \left(-\frac{1}{4} + \frac{\mu}{x} + \frac{1/4 - \mu^2}{x^2} \right) W = 0,$$

(see, for details, page 505, chapter 13, Abramowitz and Stegun [1]). Also, see Lebedev [15], and Chaudhry and Zubair [4], among others. Using the relation between Whittaker function and confluent hypergeometric function, that is,

$$W_{a,b}(z) = e^{-z/2} z^{b+1/2} U(b+1/2-a, 1+2b, z),$$

where $U(b+1/2-a, 1+2b, z)$, sometimes denoted by $\psi(b+1/2-a, 1+2b, z)$, is known as the confluent hypergeometric function of Tricomi, which is another solution of Kummer's differential equation (see page 504, Abramowitz and Stegun [1]), the normalizing function C can be written as

$$\frac{1}{C} = \beta^{(\nu-1)/2} \gamma^{(\nu-1)/2+\mu} \Gamma(1-\mu-\nu) \exp\left(\frac{\beta}{2\gamma}\right) W_{(\nu-1)/2+\mu, \frac{-\nu}{2}}\left(\frac{\beta}{\gamma}\right)$$

$$= \gamma^{\mu+\nu-1} \Gamma(1-\mu-\nu) U\left(1-\mu-\nu; 1-\nu; \frac{\beta}{\gamma}\right)$$

$$= \gamma^{\mu+\nu-1} \Gamma(1-\mu-\nu) \psi\left(1-\mu-\nu; 1-\nu; \frac{\beta}{\gamma}\right) \quad (6)$$

Using the Kummer transformation $U(a; b; z) = z^{1-b} U(1+a-b; 2-b; z)$, Equation 13.1.29, page 505, Abramowitz and Stegun [1], the normalizing function C in Equation (6) can easily be expressed as

$$\frac{1}{C} = \beta^{\nu} \gamma^{\mu-1} \Gamma(1-\mu-\nu) \psi\left(1-\mu; 1+\nu; \frac{\beta}{\gamma}\right) \quad (7)$$

$$F_X(x) = C \sum_{k=0}^{\infty} \left\{ \frac{(-1)^k \beta^{\nu+k} \gamma^{\mu-k-1} (1-\mu)_k \Gamma\left(-(\nu+k), \frac{\beta}{x}\right)}{k!} \right\}, \quad (8)$$

where C is the normalizing constant given by (6), $(1-\mu)_k = \frac{\Gamma(1-\mu+k)}{\Gamma(1-\mu)}$ denotes the Pochhammer symbol, and $\Gamma\left(-(\nu+k), \frac{\beta}{x}\right)$ denotes the incomplete gamma function. By direct differentiation of the c.d.f. in (8), and noting that $\frac{\partial \Gamma(a,t)}{\partial t} = -t^{a-1} e^{-t}$, it can be easily verified that $\frac{dF_X(x)}{dx} = f_X(x)$, where $f_X(x)$ denotes the p.d.f. of the random variable X as given in (4). The possible shapes of the *pdf f(x)* as given in (4) and the *cdf F(x)* as given in (8) are provided for some selected values of the parameters in the following Figures 1(a, b) and 2(a, b), respectively. The proposed distribution is right skewed and the effects of the parameters can easily be seen from these graphs.

3 Distributional properties

In this section, we derive moments for the proposed distribution. Some distributional relationships are established.

3.1 Fractional moments:

Using Eq. (4), the non-integer (or fractional) moment of order k of the proposed distribution is easily obtained as follows.

$$E(X^k) = \int_0^{\infty} x^k f(x) dx = C \int_0^{\infty} x^{\nu+k-1} (x+\gamma)^{(\mu-1)} \exp(-\beta x^{-1}), \quad 0 < x < \infty.$$

Since, $\int_0^{\infty} x^{\nu-1} (x+\gamma)^{(\mu-1)} \exp(-\beta x^{-1}) = \beta^{\nu} \gamma^{\mu-2} \Gamma(1-\mu-\nu) \psi\left(1-\mu-\nu; 1-\nu; \frac{\beta}{\gamma}\right)$,

we will have

$$E(X^k) = \frac{\beta^k \Gamma(1-\mu-\nu-k) \psi\left(1-\mu; 1+\nu+k; \frac{\beta}{\gamma}\right)}{\Gamma(1-\mu-\nu) \psi\left(1-\mu; 1+\nu; \frac{\beta}{\gamma}\right)},$$

or, equivalently,

$$E\left(X^{k}\right)=\frac{\gamma^{k}\,\Gamma\left(1-\mu-\nu-k\right)\psi\left(1-\mu-\nu-k;1-\nu-k;\frac{\beta}{\gamma}\right)}{\Gamma\left(1-\mu-\nu\right)\psi\left(1-\mu-\nu;1-\nu;\frac{\beta}{\gamma}\right)},$$

where $\beta>0$, $\gamma>0$, $0<\nu<1$, $0<\mu<1$, $1-\mu>\nu>0$, $\left(\nu\neq\frac{1}{2},\mu\neq\frac{1}{2}\right)$ (at the same time). Evidently, only non-integer moments (NIM), that is, fractional moments, in $0<k<1$ exist because $k<1-\mu-\nu$. One of the earliest examples in which non-integer moments (NIM) were calculated related to the spans of random walks, but more recently the properties of non-integer moments have found application in the study of random resistor networks, chaos, and diffusion-limited aggregation, see Weiss et al. [27], and references therein. Also, see Cottone and Paola [6], and Cottone et al. [7] for recent developments on fractional moments. Fractional Moments for different parameters are given in Table 3.1 and presented in Figure 3. From Table 3.1 and Figure 3, it is observed that the fractional moments $E(X^k)$ is an increasing function in k and ν for the given values of parameters.

Table 3.1: table for fractional moments

k	ν	$\mu=0.2,\ \beta=2,\ \gamma=1$	$\nu=0.2,\ \beta=2,\ \gamma=1$
0.1			
	0.1	1.261431458	1.265137937
	0.2	1.309431046	1.309431046
	0.3	1.377644178	1.373385208
	0.4	1.485576009	1.476137246
	0.5	1.691455140	1.674866820
	0.6	2.285613328	2.254809943
0.2			
	0.1	1.651757512	1.661559089
	0.2	1.803930056	1.803930056
	0.3	2.046595138	2.033849652
	0.4	2.512785173	2.480667190
	0.5	3.866012410	3.789845477
0.3			
	0.1	2.275534118	2.295986971
	0.2	2.679875211	2.679875211
	0.3	3.461723864	3.429153427
	0.4	5.743255284	5.632500198
0.4			
	0.1	3.380478894	3.421388272
	0.2	4.532888700	4.532888700
	0.3	7.912162208	7.812192904
0.5			
	0.1	5.717928397	5.805272281
	0.2	10.36043082	10.36043082
0.6			
	0.1	13.06897336	13.31088520




Figure 1(a): PDF for $\nu = 0.1, 0.3, 0.5, 0.7$ when $\beta = 2,\ \gamma = 1,\ \mu = 0.2$




Figure 1(b): PDF for $\beta = 1, 3, 5, 7$ when $\gamma = 3,\ \mu = 0.3,\ \nu = 0.4$




Figure 2(a): CDF for $\nu = 0.1, 0.3, 0.5, 0.7$ when $\beta = 2, \gamma = 1, \mu = 0.2$




Figure 2(b): CDF for $\nu = 0.1, 0.3, 0.5, 0.6$ when $\beta = 2, \gamma = 2, \mu = 0.3$




Figure 3: Plot of $E(X^{k})$, when $\mu = 0.045, \nu = 0.05, \beta = 3, \gamma = 2$.

3.2 Survival and Hazard functions

The survival and hazard functions for our newly proposed distribution are respectively given by $S(x) = 1 - F_X(x)$, and $h(x) = \dfrac{f_X(x)}{1 - F_X(x)}$, where f(x) and F(x) are given by the equations (4) and (8) respectively. The possible shapes of the survival $S(x)$ and hazard rate $h(x)$ functions corresponding to the *pdf* (4) are provided for some selected values of the parameters in the following Figures 4(a) and 4(b), respectively. The effects of the parameters can easily be seen from these graphs.




Figure 4(a) $S(x)$ for $\nu = 0.1, 0.3, 0.5, 0.7$ when $\beta = 2, \gamma = 1, \mu = 0.2$




Figure 4(b) $h(x)$ for $\nu = 0.1, 0.2, 0.3, 0.5$ when $\beta = 0.2$, $\gamma = 1$, $\mu = 0.3$

3.3 Distributional relationships

The following probability distributions are a special case of the proposed distribution with the *pdf* given by (4).

Inverted Kummer-Gamma Distribution: The Kummer-gamma distribution is defined by the probability density function (see NG and Kotz [17])

$$f_X(x) = x^{\nu-1}\left(1+x\right)^{-\mu}\exp\left(-\beta x\right),\ x>0,\beta>0,\ \nu>0,-\infty<\mu<\infty \tag{9}$$

Let$Y = \frac{1}{X}$, then it is easy to see from (9) that the inverted Kummer-gamma distribution is defined by the following probability density function

$$g_Y(y) = C\,y^{\mu-\nu-1}\left(1+y\right)^{-\mu}\exp\left(-\frac{\beta}{y}\right),\ y>0,\beta>0,\ \nu>0,-\infty<\mu<\infty \tag{10}$$

where $C = \beta^{\frac{\mu-\nu-1}{2}}\,\Gamma(1+\nu)\exp\left(\frac{\beta}{2}\right)W_{\frac{1-\mu-\nu}{2},-\frac{\mu-\nu}{2}}(\beta)$ is the normalizing constant, and $W_{\frac{1-\mu-\nu}{2},-\frac{\mu-\nu}{2}}(\beta)$ denotes the Whittaker's Function, Gradshteyn & Ryzhik [9], Equation 3.471.7, Page 340. Comparing Equations (4) and (10), it follows that the inverted Kummer-Gamma distribution (10) is a special case of the proposed distribution (4) when $\gamma = 1$, $\beta > 0$, $0 < \nu < 1$, and $0 < \mu < 1$, in Equation (10).

4 A characterization

To prove the characterization theorem (Theorem (4.1)) we need the following Lemma.

Lemma 4.1 *Suppose that a non-negative random variable X has an absolutely continuous (with respect to Lebesgue measure) cumulative distribution function (CDF) F(x) and probability density function (PDF) f(x). We assume that f'(x) exits for all x and 0 < E(X) < ∞. Further, if*

$$E(X\ |X\ \le t) = g(t)\tau(t),\ \text{where}\ \tau(t) = \frac{f(t)}{F(t)},\ \text{for all t > 0 and}\ g'(t)\ \text{exists for all t > 0, then}$$

$$f(x) = ce^{\int \frac{x-g'(x))}{g(x)}dx},\ \text{for all x > 0, where c is a constant and}\ c = \int_0^\infty f(x)dx\ .$$

Proof: It is well known that

$$\mathrm{E}\,(\mathrm{X}|\mathrm{X}\leq \mathrm{t}) = \frac{\int_0^t xf\,(x)dx}{F(t)}\,.$$

Thus

$$\int_0^t xf\,(x) = g(t)f\,(t)\,.$$

Differentiating both sides of the equation, we obtain

$$t\,f\,(t) = g'(t)f\,(t) + g(t)f'(t).$$

On simplification we obtain

$$\frac{f'(t)}{f\,(t)} = \frac{t - g'(t)}{g(t)}$$

Integrating the above expression, we obtain

$$f\,(x) = ce^{\int \frac{x-g'(x)}{g(x)}dx}\,.$$

Remark 1: In the Lemma, the left truncated conditional expectation of X considers a product of reverse hazard rate and another function of the truncated point.

Theorem 4.1: *Suppose that a non-negative random variable X has an absolutely continuous (with respect to Lebesgue measure) cumulative distribution function F(x) and probability density function f(x). If f(x) is as given in (4) if and only if*

$$g(t) = t^{1-\nu}\gamma^{1-\mu}\exp(\beta x^{-1})\sum_{k=0}^{\infty}\frac{(-1)^k\,\beta^{\nu+k+1}(1-\eta)_k\,\Gamma(-(\nu+k+1),\frac{\beta}{x})}{k\,!}. \tag{11}$$

Proof:

We have $g(t) = \dfrac{\int_0^t xf\,(x)dx}{f\,(x)}$, then substituting f(x) from (4), it is easy to show that g(x) is as given in (11).

Suppose

$$g(t) = t^{1-\nu}\gamma^{1-\mu}\exp(\beta x^{-1})\sum_{k=0}^{\infty}\frac{(-1)^k\,\beta^{\nu+k+1}(1-\eta)_k\,\Gamma(-(\nu+k+1),\frac{\beta}{x})}{k\,!},$$

Then

$$g'(t) = t - g(t)[\frac{\nu-1}{t} + \frac{\mu-1}{x+\gamma} + \frac{\beta}{x^2}]\,.$$

Thus

$$\frac{f'(t)}{f\,(t)} = \frac{t - g'(t)}{g(t)} = [\frac{\nu-1}{t} + \frac{\mu-1}{x+\gamma} + \frac{\beta}{x^2}]$$

On integrating both sides of the above expression from 0 to x, we obtain

$$\mathrm{f_X}\,(\mathrm{x}) = \mathrm{c}\;e^{\int\{\frac{\nu-1}{x}+\frac{\mu-1}{x+\gamma}+\frac{\beta}{x})dx}$$

$$= cx^{\nu-1}(x+\gamma)^{\mu-1}\exp(-\frac{\beta}{x}),$$

where c is a normalizing constant such that

$$\frac{1}{c} = \int_0^{\infty} x^{\nu-1}(x+\gamma)^{\mu-1}\exp\left(-\frac{\beta}{x}\right)dx\,.$$

This completes the proof of Theorem 4.1.

5 Conclusion

This paper derives a new class of generalized Pearson distribution. The expressions for the PDF, CDF, and moments have been obtained. The plots for the PDF, CDF, and hazard rate and survival functions are given. Some distributional relationships and the characterization of the proposed distribution have been established. For the proposed distribution,

evidently, only non-integer moments (NIM), that is, fractional moments of order k , where $0 < k < 1$, exist. We hope that the findings of the paper will be useful for researchers in different fields like economics, engineering, environmental science, finance, life testing, medical sciences, reliability theory, physical sciences, traffic data, among others, where fractional moments become required to be computed if the integer moments $k \geq 1$ do not exist. Also, it is hoped that the proposed attempt will be helpful in designing a new approach of unifying different families of distributions based on generalized Pearson differential equation.

Acknowledgment

The authors would like to thank the editor for helpful suggestions which improved the presentation of the paper.

References

[1] Abramowitz, M., and Stegun, I. A. (1970). Handbook of Mathematical Functions, with Formulas, Graphs, and Mathematical Tables. Dover, New York. Bondesson, L. (1979). A general result on infinite divisibility, Annals of Probability, 7 965-979.

[2] Ahsanullah, M., Hamedani, G. G. and Shakil, M. (2011). A Note on Distributions arising from Generalized Pearsonian System of Differential Equation. *Technical Report No. 481*, Department of Mathematics, Statistics, and Computer Science, Marquette University, 1-9.

[3] Chaudhry, M. A., and Ahmad, M. (1993). On a probability function useful in size modeling. Canadian Journal of Forest Research, 23(8), 1679–1683.

[4] Chaudhry, M. A., and Zubair, S. M. (2002). On A Class of Incomplete Gamma Functions, with Applications. Chapman & Hall/CRC, Boca Raton.

[5] Cobb, L., Koppstein, P., and Chen, N. H. (1983). Estimation and moment recursion relations for multimodal distributions of the exponential family. Journal of the American Statistical Association, 78(381), 124-130.

[6] Cottone, G., and Di Paola, M. (2009). On the use of fractional calculus for the probabilistic characterization of random variables. Probabilistic Engineering Mechanics, 24(3), 321-330.

[7] Cottone, G., Di Paola, M., and Metzler, R. (2010). Fractional calculus approach to the statistical characterization of random variables and vectors. Physica A: Statistical Mechanics and its Applications, 389(5), 909-920.

[8] Dunning, K., and Hanson, J. N. (1977). Generalized Pearson distributions and nonlinear programming. Journal of Statistical Computation and Simulation, Volume 6, Issue 2, 115 – 128.

[9] Gradshteyn, I. S. and Ryzhik, I. M. (1980). Table of Integrals, Series, and Products (6th edition). Academic Press, San Diego.

[10] Hamedani, G. G. (2011). Characterizations of the Shakil-Kibria-Singh Distribution. Austrian Journal of Statistics, Volume 40, Number 3, 201–207.

[11] Hamedani, G. G. (2013). Characterizations of Certain Continuous Distributions., Part III Statistical Analysis – Chapter 13, Page 297, in Multiscale Signal Analysis and Modeling, Shen, Xiaoping; Zayed, Ahmed I. (Eds.), 2013, 2013, XVII, 378 p. 88 illus.

[12] Johnson, N. L., Kotz, S., and Balakrishnan, N. (1994). Continuous Univariate Distributions, (Volume 1, Second edition). John Wiley & Sons, New York.

[13] Kibria, B. M. G. and Shakil, M. (2011). A New Five-Parameter Burr System of Distributions based on Generalized Pearson Differential Equation. 2011 Joint Statistical Meetings Proceedings, Section on Physical and Engineering Sciences, Miami, Fl., U.S.A., July 30-August 4.

[14] Lahcene, B. (2013). On Pearson families of distributions and its applications. African Journal of Mathematics and Computer Science Research, 6(5), 108-117.

[15] Lebedev, N. N. (1972). Special Functions and their Applications. Dover, New York.

[16] Lee, C., Famoye, F., and Alzaatreh, A. Y. (2013). Methods for generating families of univariate continuous distributions in the recent decades, WIREs Comput Stat 2013, 5, 219-238. doi: 10.1002/wics.1255.

[17] NG, K.W. and Kotz, S. (1995). Kummer-Gamma and Kummer-Beta Univariate and Multivariate Distributions. Research Report, Department of Statistics, The University of Hong Kong, Hong Kong.

[18] Pearson, K. (1895). Contributions to the mathematical theory of evolution, II: Skew variation in homogeneous material. Philosophical Transactions of the Royal Society of London, A186, 343-414.

[19] Pearson, K. (1901). Mathematical contributions to the theory of evolution X: Supplement to a memoir on skew of variation. Philosophical Transactions of the Royal Society of London. Series A, Containing Papers of a Mathematical or Physical Character, 197, 343-414.

[20] Pearson, K. (1916). Mathematical contributions to the theory of evolution, XIX: Second supplement to a memoir on skew of variation. Philosophical Transactions of the Royal Society of London. Series A, Containing Papers of a Mathematical or Physical Character, 216, 429-457.

[21] Rossani, A., and Scarfone, A. M. (2009). Generalized Pearson distributions for charged particles interacting with an electric and/or a magnetic field. Physica, A, 388, 2354-2366.

[22] Roy, L. K. (1971). An extension of the Pearson system of frequency curves. Trabajos de estadistica y de investigacion operativa, 22 (1-2), 113-123.

[23] Shakil, M., and Kibria, B. M. G. (2010). On a family of life distributions based on generalized Pearson differential equation with applications in health statistics. Journal of Statistical Theory and Applications, 9 (2), 255-282.

[24] Shakil, M., Singh, J. N., and Kibria, B. M. G. (2010a). On a family of product distributions based on the Whittaker functions and generalized Pearson differential equation. Pakistan Journal of Statistics, 26(1), 111-125.

[25] Shakil, M., Kibria, B. M. G., and Singh, J. N. (2010b). A new family of distributions based on the generalized Pearson differential equation with some applications. Austrian Journal of Statistics, 39 (3), 259–278.

[26] Shakil, M., Kibria, B. M. G. and Singh, J. N. (2011). On Some New Classes of Pearsonian Distributions, *2011 Joint Statistical Meetings Proceedings, Section on Physical and Engineering Sciences*, Miami, Fl., U.S.A., July 30-August 4.

[27] Weiss, G. H., Havlin, S., and Matan, O. (1989). Properties of noninteger moments in a first passage time problem. Journal of Statistical Physics, 55(1-2), 435-439.