

Common fixed point theorems in fuzzy partial metric spaces of two pairs of mapping satisfying E.A. property

Renu P. Pathak *, Rohini R. Gore

Department of Mathematics, Sandip University, Nashik

*Corresponding author E-mail: renupathak380@gmail.com

Received: April 5, 2025, Accepted: May 3, 2025, Published: May 11, 2025

Abstract

This paper establishes a common fixed point theorem for a pair of weakly compatible mappings in Fuzzy PMS satisfying the E.A. property. We introduce a new concept of E.A. Fuzzy Partial Metric spaces which generalizes the existing Fuzzy PMS. Our main result provides sufficient conditions for the existence of a common fixed point for two weakly compatible mappings in E.A. Fuzzy Partial Metric spaces. The results have significant implications for fuzzy topology, fixed-point theory, and applications in image processing, machine learning, and optimization problems involving uncertain or imprecise data.

Keywords: Partial Metric Space; Fuzzy Metric Space; Weakly Compatible Mapping; Common Fixed Point.

1. Introduction

A metric space is a fundamental concept in Mathematics, particularly in the field of analysis and topology. It provides a framework for measuring distances between points in a set. The modern concept of a metric space was first introduced by Maurice Fréchet in 1906. In his work on functional analysis, Fréchet defined a metric space as a set equipped with a distance function satisfying certain properties, laying the groundwork for the rigorous study of distances in mathematical contexts.

In 1965, Fuzzy set theory was developed by Zadeh [24]. Fuzzy sets allow elements to have degrees of membership between 0 and 1, rather than strictly belonging or not belonging to a set. In the 1970s, mathematicians began to explore the concept of fuzzy metric spaces, where the distance function is allowed to take fuzzy values. In 1975, Kramosil and Michalek first introduced the concept of a fuzzy metric space [9], which can be regarded as a generalization of the statistical metric space. Clearly this work provides an important basis for the construction of fixed point theory in fuzzy metric space. Fuzzy metric spaces generalize traditional metric spaces by accommodating uncertainty in distance measurements. Fixed point theory has been a vibrant area of research in Mathematics, with numerous applications in various fields such as physics, engineering, economics, and computer science. The generalizations of metric spaces are the notion of partial metric space which was given by Matthews [15] as an extension of metric space where the self-distance of any point is not necessarily equal to zero. Partial metric spaces relax some of the conditions of metric spaces, such as the triangle inequality, to provide a more flexible framework for modeling certain scenarios. In 1994, Veeramani and George introduced the modified Continuous t-norm [7]. In recent years, researchers have explored fixed point theorems in Fuzzy Partial Metric spaces, but most of these results rely on strong assumptions, such as continuity or compatibility of mappings. To overcome these limitations, we introduce the E.A. property, a relaxed equivalence relation that allows for more flexibility in modeling real-world problems.

A FPMS is a generalization of the classical metric space where distances can take fuzzy values, allowing for the representation of uncertainty in distance measurement. In traditional metric spaces, the distance between two points is a non-negative real no. In contrast, in a FPMS, the distance can be represented as a fuzzy no. which can accommodate both the notion of closeness & uncertainty.

Impact of E, A. Property on common fixed point theorems in Fuzzy Metric space presented by Kamal Wadhawa [11]. Ashlekha proved fixed point theorem in FMS with E. A. Property by using weakly compatible mapping [2]. Nazir present common fixed point results for two pairs of mapping which is satisfy E. A. property in Partial Metric space [23]. Madhushrivastava present some fixed point theorems in FMS by using occasionally weakly compatible mapping [14]. The some common fixed point theorems for weakly compatible self mapping in FMS using (E.A.) property is generalized by Kanhaiya [10]. The new concept introduce of common fixed point theorem in FMS & occasionally compatible mapping by subhani [22]. Rakhi Namdev is developed for existence & uniqueness of common fixed point by using two pairs of mapping satisfying E.A. property [18].

The other author's given different concept of Fixed point theorems by using E. A. property in different spaces like S-metric space, Menger space, Complex valued metric space [5],[8],[13],[19]. Sonam Soni explained applications of Fixed points in Computational Mathematics by using Fuzzy & Partial metric spaces [20].

In this paper, we establish a common fixed point theorem for a pair of weakly compatible mappings in Fuzzy Partial Metric spaces satisfying the E.A. property. Our work builds upon existing results in Fuzzy Partial Metric spaces and fixed point theory, providing a more

comprehensive framework for analyzing complex systems with uncertain or imprecise data. The E.A. property captures nuanced relationships, enabling a more general fixed point theorem.

To establish our theorems, we first define key concepts:

2. Preliminaries

Definition 1: A Partial Metric space on ' $\mathbb{X}$ ' is a Pair $(\mathbb{X}, P)$ such that ' $\mathbb{X}$ ' is a non-empty set and $P: \mathbb{X} \times \mathbb{X} \rightarrow \mathbb{R}^+$ is a mapping providing the listed conditions $\forall p, q, r \in \mathbb{X}$ such that

- 1) ' $P(p, p) \leq P(p, q)$ '
- 2) ' $P(p, p) = P(q, q) = P(p, q)$ ' iff ' $p = q$ '
- 3) ' $P(p, q) = P(q, p)$ '
- 4) ' $P(p, r) \leq P(p, q) + P(q, r) - P(q, q)$ '

Note that in Partial Metric space, a point's self-distance does not always equal zero. The Partial Metric ' P ' is an ordinary metric on ' $\mathbb{X}$ ' if ' $P(p, p) = 0$ ', $\forall p \in \mathbb{X}$. Therefore, a Partial Metric is an extension of an ordinary metric.

Definition 2: A binary operation ' $\odot$ ' on $[0, 1]$ is called a continuous t-norm if it is satisfied following conditions: $\forall p, q, r, s \in [0, 1]$:

- 1) ' $p \odot q = q \odot p$ ' and ' $p \odot (q \odot r) = (p \odot q) \odot r$ '
- 2) ' $\odot$ is continuous on $[0, 1] \times [0, 1]$ '
- 3) ' $p \odot 1 = p$ '
- 4) If ' $p \leq q$ ' and ' $r \leq s$ ', then ' $p \odot r \leq q \odot s$ '

Definition 3: Let ' $\mathbb{X}$ ' be a non-empty set, ' $\odot$ ' be a continuous t-norm and $F: \mathbb{X} \times \mathbb{X} \times (0, \infty) \rightarrow [0, 1]$ be a mapping. Let ' F ' be Fuzzy set. If the listed conditions are satisfied $\forall p, q, r \in \mathbb{X}$ and $u, v > 0$, then the triplet $(\mathbb{X}, F, \odot)$ is said to be a Fuzzy Metric space, If it satisfies following Properties for

- 1) ' $F(p, q, u) > 0$ '
- 2) ' $F(p, q, u) = 1$ ' iff ' $q = p$ '
- 3) ' $F(p, q, u) = F(q, p, u)$ '
- 4) ' $F(p, q, u + v) \geq F(p, q, u) \odot F(p, q, v)$ '
- 5) ' $F(p, q, \odot)$ is continuous on $(0, \infty)$ '

If $(\mathbb{X}, F, \odot)$ is a fuzzy metric space, then ' F ' is a fuzzy metric on ' $\mathbb{X}$ '.

Definition 4: Let ' $\mathbb{X}$ ' be a non-empty set, ' $\odot$ ' be a continuous t-norm and $F_p: \mathbb{X} \times \mathbb{X} \times (0, \infty) \rightarrow [0, 1]$ be a mapping. Let P be Partial metric space. If the listed conditions are satisfied $\forall p, q, r \in \mathbb{X}$ and $u, v > 0$, then the triplet $(\mathbb{X}, F_p, \odot)$ is said to be a Fuzzy Partial Metric space:

- 1) ' $F_p(p, q, 0) = 0$ '
- 2) ' $F_p(p, q, u) = F_p(q, p, u)$ '
- 3) ' $F_p(p, r, u + v) \geq F_p(p, q, u) \odot F_p(q, r, v)$ '
- 4) ' $F_p(p, q, u) \leq 1, u > 0$ ' & ' $F_p(p, q, u) = 1$ ' iff ' $p(p, q) = 0$ '
- 5) ' $F_p(p, q, \odot): (0, \infty) \rightarrow [0, 1]$ is continuous', Where ' $F_p(p, q, u) = \frac{u}{u + p(p, q)}$ '

If $(\mathbb{X}, F_p, \odot)$ is a Fuzzy Partial Metric space, then ' F_p ' is a fuzzy partial metric on ' $\mathbb{X}$ '.

Example1: Let ' $\mathbb{X} = \{p, q, r\}$ ' be a set of points. Define a fuzzy distance p as $p(p, p) = 1$,

$$p(q, q) = 1, p(p, q) = 0.5, p(q, p) = 0.5, p(r, r) = 1,$$

$$p(p, r) = 0.2, p(r, p) = 0.2, p(q, r) = 0.3, p(r, q) = 0.3.$$

In above example, $p(p, q) = 0.5$ indicate that p & q are relatively close, $p(p, r) = 0.2$ indicate that p & r are not close.

Definition 5: Two self-mapping ' M ' and ' N ' on Fuzzy Partial mapping $(\mathbb{X}, F_p, \odot)$ are Compatible, if ' $\lim_{n \rightarrow \infty} F_p(MN y_n, NM y_n, u) = 1$ ', > 0 , whenever $\{y_n\}$ is a sequence in ' $\mathbb{X}$ ' such that ' $\lim_{n \rightarrow \infty} M y_n = \lim_{n \rightarrow \infty} N y_n = t$ ', $\forall t \in \mathbb{X}$ '.

Definition 6: Two self-mapping ' M ' and ' N ' on Fuzzy Partial mapping $(\mathbb{X}, F_p, \odot)$ are a weakly Compatible, if they commute at their coincidence point. $\forall t \in \mathbb{X}$, ' $M t = N t$ ' implies that ' $M N t = N M t$ ', $\forall t > 0$.

Definition 7: Two self-mapping ' M ' and ' N ' on Fuzzy Partial mapping $(\mathbb{X}, F_p, \odot)$ are satisfy E. A. property if $\exists$ a sequence $\{y_n\}$ is a sequence in ' $\mathbb{X}$ ' such that ' $\lim_{n \rightarrow \infty} M y_n = \lim_{n \rightarrow \infty} N y_n = t$ ', $\forall t \in \mathbb{X}$ '.

Definition 8: Let ' M ', ' N ', ' U ' and ' V ' be a self-mapping of a Fuzzy Partial Metric space $(\mathbb{X}, F_p, \odot)$. Then (M, U) and (N, V) are satisfy common E. A. property, if $\exists$ two sequences $\{x_n\}$ and $\{y_n\}$ in ' $\mathbb{X}$ ' such that ' $\lim_{n \rightarrow \infty} M x_n = \lim_{n \rightarrow \infty} U x_n = \lim_{n \rightarrow \infty} N y_n = \lim_{n \rightarrow \infty} V y_n = z$ ', where $z \in M(\mathbb{X}) \cap N(\mathbb{X})$ or $z \in U(\mathbb{X}) \cap V(\mathbb{X})$.

Example2: Let $'\mathbb{X}' = [0,2)$ and $'F_p(p, q, u) = \frac{u}{u+p(p,q)}'$ $\forall p, q \in '\mathbb{X}'$. Then $('\mathbb{X}', F_p, \odot)$ is a Fuzzy Partial Metric space. where $a \odot b = \min(a, b)$.

$$M(p) = 0.25, 0 \leq p \leq 0.52, U(p) = 0.25, 0 \leq p \leq 0.60,$$

$$= \frac{p}{2}, p > 0.52, = p - 0.25, p > 0.60,$$

$$V(p) = 0.25, 0 \leq p \leq 0.60, N(p) = 0.25, 0 \leq p \leq 0.95,$$

$$= \frac{p}{4}, p > 0.60, = p - 0.75, p > 0.95,$$

We define $x_n = 0.5 + \frac{1}{n}$, $y_n = 1 + \frac{1}{n}$, Where $M('X') = \{0.25\} \cup (0.26, 1]$

$$U('X') = \{0.25\} \cup (0.35, 1.75], V('X') = (0.15, 0.5], N('X') = \{0.25\} \cup (0.20, 1.25]$$

$$\lim_{n \rightarrow \infty} Mx_n = \lim_{n \rightarrow \infty} \frac{1}{2} \left[0.5 + \frac{1}{n} \right] = 0.25 \in U('X'),$$

$$\lim_{n \rightarrow \infty} Ux_n = \lim_{n \rightarrow \infty} \left[0.5 + \frac{1}{n} \right] - 0.25 = 0.25 \in M('X') \quad \lim_{n \rightarrow \infty} Ny_n = \lim_{n \rightarrow \infty} \left[1 + \frac{1}{n} \right] - 0.75 = 0.25 \in V('X').$$

$$\lim_{n \rightarrow \infty} Vy_n = \lim_{n \rightarrow \infty} \frac{1}{4} \left[1 + \frac{1}{n} \right] = 0.25 \in N('X')$$

$$\text{Thus, } \lim_{n \rightarrow \infty} Mx_n = \lim_{n \rightarrow \infty} Ux_n = \lim_{n \rightarrow \infty} Ny_n = \lim_{n \rightarrow \infty} Vy_n = 0.25$$

Where, $0.25 \in M('X') \cap N('X')$ or $0.25 \in U('X') \cap V('X')$.

Role of E.A. Property in providing common fixed point theorems in FPMS can be concluded followings: 1) The E. A. property helps determine whether sequences in the FPMS converges to a point within the space.

- 1) Many fixed point theorems generalized results, rely on the E. A. property to confirm the existence of a fixed point for certain continuous mapping in FPMS.
- 2) The E. A. property can link topological properties of FPMS to behavior of sequences, which can study of properties like compactness or completeness.
- 3) Understanding the E. A. property can lead to the applications in Mathematical modeling, optimization problems & other interdisciplinary fields where fuzzy logic is used.

Lemma: Let $(\mathbb{X}, F_p, \odot)$ be a Fuzzy Partial Metric space, if there exist $k \in (0,1)$ such that $'F_p(p, q, k u) \geq F_p(p, q, u)'$ then $'p = q', u > 0, k \in (0,1), \forall p, q \in '\mathbb{X}'$.

3. Main results

Theorem 1: Let M, N, U, V be a self-mapping of a Fuzzy Partial Metric space $(\mathbb{X}, F_p, \odot)$ with $a \odot b = \min(a, b)$ satisfying following conditions:

- i) (M, U) and (N, V) satisfy E. A. property.
- ii) (M, U) and (N, V) are weakly compatible.
- iii) Then there exist $k \in (0,1)$, for all $x, y, z \in '\mathbb{X}'$ and $u > 0$.

$$F_p(Mx, Ny, ku) \geq \min [F_p(Ux, Vy, u), F_p(Ny, Vy, u), (F_p(Ux, Vy, u) \cdot F_p(Ny, Vy, u)),$$

$$\frac{F_p(Ny, Vy, u)}{F_p(Ny, Ux, u)}, F_p(Ny, Mx, u)]$$

Then M, N, U, V have a common fixed point.

Proof: Since (M, U) and (N, V) satisfying E. A. property then there exist two sequences $\{x_n\}$ and $\{y_n\}$ in $'\mathbb{X}'$ such that $\lim_{n \rightarrow \infty} Mx_n = \lim_{n \rightarrow \infty} Ux_n = \lim_{n \rightarrow \infty} Nx_n = \lim_{n \rightarrow \infty} Vy_n = z$, where $z \in M('X') \cap N('X')$ or $z \in U('X') \cap V('X')$. Then $z = Ut$, where $x \in '\mathbb{X}'$.

- a) Now to prove that $p(Mt, Ut) = 0$, that is to show that $Mt = z$.

Put $x = t, y = y_n$ in (iii) we get,

$$F_p(Mt, Ny_n, ku) \geq \min [F_p(Ut, Vy_n, u), F_p(Ny_n, Vy_n, u), (F_p(Ut, Vy_n, u) \cdot F_p(Ny_n, Vy_n, u)),$$

$$\frac{F_p(Ny_n, Vy_n, u)}{F_p(Ny_n, Ut, u)}, F_p(Ny_n, Mt, u)]$$

$$F_p(Mt, Ny_n, ku) \geq \min [F_p(z, z, u), F_p(Ny_n, z, u), (F_p(z, z, u) \cdot F_p(Ny_n, z, u)),$$

$$\frac{F_p(Ny_n, z, u)}{F_p(Ny_n, z, u)}, F_p(Ny_n, Mt, u)]$$

$$F_p(Mt, Ny_n, ku) \geq \min [1, F_p(Ny_n, z, u), (1. F_p(Ny_n, z, u)), 1, F_p(Ny_n, Mt, u)]$$

$$F_p(Mt, Ny_n, ku) \geq F_p(Ny_n, z, u), \text{ if } Mt = z$$

By using Lemma1, we get $Mt = Ny_n = z$. Thus $Mt = z$

Since (M, U) is weakly compatible. Then $Mz = MUt = UMt = Uz = z$

b) Now to prove that $p(Vz, Nz) = 0$, that is to show that $Nz = z$

Put $x = x_n, y = z$ in (iii) we get,

$$F_p(Mx_n, Nz, ku) \geq \min [F_p(Ux_n, Vz, u), F_p(Nz, Vz, u), (F_p(Ux_n, Vz, u). F_p(Nz, Vz, u)),$$

$$\frac{F_p(Nz, Vz, u)}{F_p(Nz, Ux_n, u)}, F_p(Nz, Mx_n, u)]$$

$$F_p(Mx_n, Nz, ku) \geq \min [F_p(z, z, u), F_p(Nz, z, u), (F_p(z, z, u). F_p(Nz, z, u)),$$

$$\frac{F_p(Nz, z, u)}{F_p(Nz, z, u)}, F_p(Nz, Mx_n, u)]$$

$$F_p(Mx_n, Nz, ku) \geq \min [1, F_p(Nz, z, u), (1. F_p(Nz, z, u)), 1, F_p(Nz, Mx_n, u)]$$

$$F_p(Mx_n, Nz, ku) \geq F_p(z, Mx_n, u), \text{ if } Nz = z. \text{ By using Lemma 1, we get } Mx_n = Nz = z$$

Since (N, V) is weakly compatible. Then $Nz = Vz = NVz = VNz = z$

c) Now to prove that $p(Mz, Uz) = 0$, that is to show that $Mz = z$.

Put $x = z, y = y_n$ in (iii) we get,

$$F_p(Mz, Ny_n, ku) \geq \min [F_p(Uz, Vy_n, u), F_p(Ny_n, Vy_n, u), (F_p(Uz, Vy_n, u). F_p(Ny_n, Vy_n, u)),$$

$$\frac{F_p(Ny_n, Vy_n, u)}{F_p(Ny_n, Uz, u)}, F_p(Ny_n, Mz, u)]$$

$$F_p(Mz, Ny_n, ku) \geq \min [F_p(z, z, u), F_p(Ny_n, z, u), (F_p(z, z, u). F_p(Ny_n, z, u)),$$

$$\frac{F_p(Ny_n, z, u)}{F_p(Ny_n, z, u)}, F_p(Ny_n, Mz, u)]$$

$$F_p(Mz, Ny_n, ku) \geq \min [1, F_p(Ny_n, z, u), (1. F_p(Ny_n, z, u)), 1, F_p(Ny_n, Mz, u)]$$

$$F_p(Mz, Ny_n, ku) \geq (Ny_n, z, u), \text{ if } Mz = z. \text{ By using Lemma 1, we get } Mz = Ny_n = z$$

Since (M, U) is weakly compatible. Then $Mz = MUz = UMz = Uz = z$

d) Now to prove that $p(Vt, Nt) = 0$, that is to show that $Nt = z$

Put $x = x_n, y = t$ in (iii) we get,

$$F_p(Mx_n, Nt, ku) \geq \min [F_p(Ux_n, Vt, u), F_p(Nt, Vt, u), (F_p(Ux_n, Vt, u). F_p(Nt, Vt, u)),$$

$$\frac{F_p(Nt, Vt, u)}{F_p(Nt, Ux_n, u)}, F_p(Nt, Mx_n, u)]$$

$$F_p(Mx_n, Nt, ku) \geq \min [F_p(z, z, u), F_p(Nt, z, u), (F_p(z, z, u). F_p(Nt, z, u)),$$

$$\frac{F_p(Nt, z, u)}{F_p(Nt, z, u)}, F_p(Nt, Mx_n, u)]$$

$$F_p(Mx_n, Nt, ku) \geq \min [1, F_p(Nt, z, u), (1. F_p(Nt, z, u)), 1, F_p(Nt, Mx_n, u)]$$

$$F_p(Mx_n, Nt, ku) \geq F_p(Nt, z, u), \text{ if } Nt = z. \text{ By using Lemma1, } Mx_n = Nt = z$$

Since (N, V) is weakly compatible. Then $Nt = Vt = NVt = VNt = z$

Thus, $Mz = Uz = Nz = Vz = z$. Therefore M, N, U, V have a common fixed point.

Uniqueness: Suppose t & z are two common fixed point of M, N, U, V with $t \neq z$,

Then (iii) becomes

$$F_p(Mt, Nz, ku) \geq \min [F_p(Ut, Vz, u), F_p(Nz, Vz, u), (F_p(Ut, Vz, u). F_p(Nz, Vz, u)),$$

$$\frac{F_p(Nz, Vz, u)}{F_p(Nz, Ut, u)}, F_p(Nz, Mt, u)]$$

$$\text{Put } Mt = Ut = t, Nz = Vz = z$$

$$F_p(t, z, ku) \geq \min [F_p(t, z, u), F_p(z, z, u), ((t, z, u) \cdot F_p(z, z, u)) \cdot \frac{F_p(z, z, u)}{F_p(z, t, u)}, F_p(z, t, u)]$$

$$F_p(t, z, ku) \geq \min [F_p(t, z, u), 1, F_p(t, z, u) \cdot 1, \frac{1}{F_p(z, t, u)}, F_p(z, t, u)]$$

$$F_p(t, z, ku) \geq F_p(t, z, u)$$

By using Lemma 1, $t = z$. Hence proved.

Theorem 2: Let M, N, U, V be a self-mapping of a Fuzzy Partial Metric space $(\mathbb{X}, F_p, \odot)$ with $\odot$ & $= a \odot b$ satisfying following conditions:

- i) (M, U) and (N, V) satisfy E. A. property.
- ii) (M, U) and (N, V) are weakly compatible.
- iii) Then their exist $k \in (0, 1)$, for all $x, y, z \in \mathbb{X}$ and $u > 0$.

$$F_p(Mx, Ny, ku) \geq [F_p(Ux, Vy, u) \odot \frac{F_p(Ny, Vy, u)}{F_p(Ny, Mx, u)} \odot (F_p(Ux, Vy, u) \cdot F_p(Ny, Vy, u)) \odot$$

$$\frac{F_p(Ny, Vy, u)}{F_p(Ny, Ux, u)} \odot \frac{F_p(Ny, Mx, u)}{F_p(Ny, Vy, u)}]$$

Then M, N, U, V have a common fixed point.

Proof: Since (M, U) and (N, V) satisfying E. A. property then their exist two sequences $\{x_n\}$ and $\{y_n\}$ in $\mathbb{X}$ such that $\lim_{n \rightarrow \infty} Mx_n = \lim_{n \rightarrow \infty} Uy_n = \lim_{n \rightarrow \infty} Nx_n = \lim_{n \rightarrow \infty} Vy_n = z$, where $z \in M(\mathbb{X}) \cap N(\mathbb{X})$ or $z \in U(\mathbb{X}) \cap V(\mathbb{X})$. Then $z = Ut$, where $x \in \mathbb{X}$.

a) Now to prove that $p(Mt, Ut) = 0$, that is to show that $Mt = z$.

Put $x = t, y = y_n$ in (iii) we get,

$$F_p(Mt, Ny_n, ku) \geq [F_p(Ut, Vy_n, u) \odot \frac{F_p(Ny_n, Vy_n, u)}{F_p(Ny_n, Mt, u)} \odot (F_p(Ut, Vy_n, u) \cdot F_p(Ny_n, Vy_n, u)) \odot$$

$$\frac{F_p(Ny_n, Vy_n, u)}{F_p(Ny_n, Ut, u)} \odot \frac{F_p(Ny_n, Mt, u)}{F_p(Ny_n, Vy_n, u)}]$$

$$F_p(Mt, Ny_n, ku) \geq [F_p(z, z, u) \odot \frac{F_p(Ny_n, z, u)}{F_p(Ny_n, Mt, u)} \odot (F_p(z, z, u) \cdot F_p(Ny_n, z, u)) \odot$$

$$\frac{F_p(Ny_n, z, u)}{F_p(Ny_n, z, u)} \odot \frac{F_p(Ny_n, Mt, u)}{F_p(Ny_n, z, u)}]$$

$$F_p(Mt, Ny_n, ku) \geq [1 \odot \frac{F_p(Ny_n, z, u)}{F_p(Ny_n, Mt, u)} \odot (1 \cdot F_p(Ny_n, z, u)) \odot 1 \odot \frac{F_p(Ny_n, Mt, u)}{F_p(Ny_n, z, u)}]$$

$$F_p(Mt, Ny_n, ku) \geq F_p(Ny_n, z, u), \text{ if } Mt = z.$$

By using Lemma1, we get $t = z$. Therefore $Mt = Ut = z$

Since (M, U) is weakly compatible. Then $Mz = MUt = UMt = Uz = z$

b) Now to prove that $p(Vz, Nz) = 0$, that is to show that $Nz = z$

Put $x = x_n, y = z$ in (iii) we get, put $Vz = z = Ux_n = Mx_n$, show that $Nz = z$

$$F_p(Mx_n, Nz, ku) \geq [F_p(Ux_n, Vz, u) \odot \frac{F_p(Nz, Vz, u)}{F_p(Nz, Mx_n, u)} \odot (F_p(Ux_n, Vz, u) \cdot F_p(Nz, Vz, u)) \odot$$

$$\frac{F_p(Nz, Vz, u)}{F_p(Nz, Ux_n, u)} \odot \frac{F_p(Nz, Mx_n, u)}{F_p(Nz, Vz, u)}]$$

$$F_p(Mx_n, Nz, ku) \geq [F_p(z, z, u) \odot \frac{F_p(Nz, z, u)}{F_p(Nz, Mx_n, u)} \odot (F_p(z, z, u) \cdot F_p(Nz, z, u)) \odot$$

$$\frac{F_p(Nz, z, u)}{F_p(Nz, z, u)} \odot \frac{F_p(Nz, Mx_n, u)}{F_p(Nz, z, u)}]$$

$$F_p(Mx_n, Nz, ku) \geq [1 \odot \frac{F_p(Nz, z, u)}{F_p(Nz, Mx_n, u)} \odot (1 \cdot F_p(Nz, z, u)) \odot 1 \odot \frac{F_p(Nz, Mx_n, u)}{F_p(Nz, z, u)}]$$

$$F_p(Mx_n, Nz, ku) \geq F_p(Nz, z, u), \text{ if } Nz = z$$

Since (N, V) is weakly compatible. Then $Nz = Vz = NVz = VNz = z$

c) Now to prove that $p(Mz, Uz) = 0$, that is to show that $Mz = z$.

Put $x = z, y = y_n$ in (iii) we get,

$$F_p(Mz, Ny_n, ku) \geq [F_p(Uz, Vy_n, u) \odot \frac{F_p(Ny_n, Vy_n, u)}{F_p(Ny_n, Mz, u)} \odot (F_p(Uz, Vy_n, u) \cdot F_p(Ny_n, Vy_n, u)) \odot$$

$$\frac{F_p(Ny_n, Vy_n, u)}{F_p(Ny_n, Uz, u)} \odot \frac{F_p(Ny_n, Mz, u)}{F_p(Ny_n, Vy_n, u)}]$$

$$F_p(Mz, Ny_n, ku) \geq [F_p(z, z, u) \odot \frac{F_p(Ny_n, z, u)}{F_p(Ny_n, Mz, u)} \odot (F_p(z, z, u) \cdot F_p(Ny_n, z, u)) \odot$$

$$\frac{F_p(Ny_n, z, u)}{F_p(Ny_n, z, u)} \odot \frac{F_p(Ny_n, Mz, u)}{F_p(Ny_n, z, u)}]$$

$$F_p(Mz, Ny_n, ku) \geq [1 \odot \frac{F_p(Ny_n, z, u)}{F_p(Ny_n, Mz, u)} \odot (1 \cdot F_p(Ny_n, z, u)) \odot 1 \odot \frac{F_p(Ny_n, Mz, u)}{F_p(Ny_n, z, u)}]$$

$$F_p(Mz, Ny_n, ku) \geq F_p(Ny_n, z, u), \text{ if } Mz = z.$$

Since (M, U) is weakly compatible. Then $Mz = MUz = UMz = Uz = z$

d) Now to prove that $p(Vt, Nt) = 0$, that is to show that $Nt = z$

Put $x = x_n, y = t$ in (iii) we get,

$$F_p(Mx_n, Nt, ku) \geq [F_p(Ux_n, Vt, u) \odot \frac{F_p(Nt, Vt, u)}{F_p(Nt, Mx_n, u)} \odot (F_p(Ux_n, Vt, u) \cdot F_p(Nt, Vt, u)) \odot$$

$$\frac{F_p(Nt, Vt, u)}{F_p(Nt, Ux_n, u)} \odot \frac{F_p(Nt, Mx_n, u)}{F_p(Nt, Vt, u)}]$$

$$F_p(Mx_n, Nt, ku) \geq [F_p(z, z, u) \odot \frac{F_p(Nt, z, u)}{F_p(Nt, Mx_n, u)} \odot (F_p(z, z, u) \cdot F_p(Nt, z, u)) \odot$$

$$\frac{F_p(Nt, z, u)}{F_p(Nt, z, u)} \odot \frac{F_p(Nt, Mx_n, u)}{F_p(Nt, z, u)}]$$

$$F_p(Mx_n, Nt, ku) \geq [1 \odot \frac{F_p(Nt, z, u)}{F_p(Nt, Mx_n, u)} \odot (1 \cdot F_p(Nt, z, u)) \odot 1 \odot \frac{F_p(Nt, Mx_n, u)}{F_p(Nt, z, u)}]$$

$$F_p(Mx_n, Nt, ku) \geq F_p(Nt, z, u), \text{ if } Nt = z$$

Since (N, V) is weakly compatible. Then $Nt = Vt = NVt = VNt = z$

Thus, $Mz = Uz = Nz = Vz = z$. Therefore M, N, U, V have a common fixed point.

Uniqueness:

Suppose t & z are two common fixed point of M, N, U, V with $t \neq z$, Then (iii) becomes

$$F_p(Mt, Nz, ku) \geq [F_p(Ut, Vz, u) \odot \frac{F_p(Nz, Vz, u)}{F_p(Nz, Mt, u)} \odot (F_p(Ut, Vz, u) \cdot F_p(Nz, Vz, u)) \odot$$

$$\frac{F_p(Nz, Vz, u)}{F_p(Nz, Ut, u)} \odot \frac{F_p(Nz, Mt, u)}{F_p(Nz, Vz, u)}]$$

$$\text{Put } Mt = Ut = t, Nz = Vz = z$$

$$F_p(t, z, ku) \geq [F_p(t, z, u) \odot \frac{F_p(z, z, u)}{F_p(z, t, u)} \odot (F_p(t, z, u) \cdot F_p(z, z, u)) \odot \frac{F_p(z, z, u)}{F_p(z, t, u)} \odot \frac{F_p(z, t, u)}{F_p(z, z, u)}]$$

$$F_p(t, z, ku) \geq [F_p(t, z, u) \odot \frac{1}{F_p(z, t, u)} \odot (F_p(t, z, u) \cdot 1) \odot \frac{1}{F_p(z, t, u)} \odot \frac{F_p(z, t, u)}{1}]$$

$$F_p(t, z, ku) \geq F_p(t, z, u)$$

By using Lemma 1, $t = z$. Hence proved.

Definition 9: Let φ be the class of all mapping $\varphi : [0, 1] \rightarrow [0, 1]$ such that

- φ is non-decreasing and $\lim_{n \rightarrow \infty} \varphi^n(l) = 1, \forall l \in (0, 1]$
- $\varphi(l) > l, \forall l \in (0, 1)$ iii) $\varphi(1) = 1$

Theorem 3: Let M, N, U, V be a self-mapping of a Fuzzy Partial Metric space $(\mathbb{X}, F_p, \odot)$ with $a \odot b = \min(a, b)$ satisfying following conditions:

- i) (M, U) and (N, V) satisfy E. A. property.
- ii) (M, U) and (N, V) are weakly compatible.
- iii) Then their exist $k \in (0, 1)$, for all $x, y, z \in \mathbb{X}$ and $u > 0$.

$$F_p(Mx, Ny, ku) \geq \varphi \left\{ \min \left[\frac{F_p(Ux, Vy, u)F_p(Ny, Ux, u)}{F_p(Mx, Vy, u)}, \frac{F_p(Ux, Vy, u)F_p(Ny, Vy, u)}{F_p(Mx, Vy, u)}, \frac{F_p(Ux, Vy, u)F_p(Ny, Vy, u)}{F_p(Ny, Ux, u)}, \right. \right. \\ \left. \left. \frac{F_p(Ny, Vy, u)F_p(Mx, Vy, u)}{F_p(Ny, Ux, u)}, \frac{F_p(Mx, Vy, u)F_p(Ny, Mx, u)}{F_p(Vy, Ux, u)} \right] \right\}$$

Then M, N, U, V have a common fixed point.

Proof: Since (M, U) and (N, V) satisfying E. A. property then their exist two sequences $\{x_n\}$ and $\{y_n\}$ in $\mathbb{X}$ such that $\lim_{n \rightarrow \infty} Mx_n = \lim_{n \rightarrow \infty} Uy_n = \lim_{n \rightarrow \infty} Nx_n = \lim_{n \rightarrow \infty} Vy_n = z$, where $z \in M(\mathbb{X}) \cap N(\mathbb{X})$ or $z \in U(\mathbb{X}) \cap V(\mathbb{X})$. Then $z = Ut$, where $x \in \mathbb{X}$.

- a) Now to prove that $p(Mt, Ut) = 0$, that is to show that $Mt = z$.

Put $x = t, y = y_n$ in (iii) we get,

$$F_p(Mt, Ny_n, ku) \geq \varphi \left\{ \min \left[\frac{F_p(Ut, Vy_n, u)F_p(Ny_n, Ut, u)}{F_p(Mt, Vy_n, u)}, \frac{F_p(Ut, Vy_n, u)F_p(Ny_n, Vy_n, u)}{F_p(Mt, Vy_n, u)}, \frac{F_p(Ut, Vy_n, u)F_p(Ny, Vy_n, u)}{F_p(Ny_n, Ut, u)}, \right. \right. \\ \left. \left. \frac{F_p(Ny_n, Vy_n, u)F_p(Mt, Vy_n, u)}{F_p(Ny_n, Ut, u)}, \frac{F_p(Mt, Vy_n, u)F_p(Ny_n, Mt, u)}{F_p(Vy_n, Ut, u)} \right] \right\}$$

$$F_p(Mt, Ny_n, ku) \geq \varphi \left\{ \min \left[\frac{F_p(z, z, u)F_p(Ny_n, z, u)}{F_p(Mt, z, u)}, \frac{F_p(z, z, u)F_p(Ny_n, z, u)}{F_p(Mt, z, u)}, \frac{F_p(z, z, u)F_p(Ny, z, u)}{F_p(Ny_n, z, u)}, \right. \right. \\ \left. \left. \frac{F_p(Ny_n, z, u)F_p(Mt, z, u)}{F_p(Ny_n, z, u)}, \frac{F_p(Mt, z, u)F_p(Ny_n, Mt, u)}{F_p(z, z, u)} \right] \right\}$$

$$F_p(Mt, Ny_n, ku) \geq \varphi \left\{ \min \left[\frac{1 \cdot F_p(Ny_n, z, u)}{F_p(Mt, z, u)}, \frac{1 \cdot F_p(Ny_n, z, u)}{F_p(Mt, z, u)}, 1, 1, 1, F_p(Mt, z, u), \frac{F_p(Mt, z, u)F_p(Ny_n, Mt, u)}{1} \right] \right\}$$

$$F_p(Mt, Ny_n, ku) \geq \varphi \left\{ \min \left[\frac{F_p(Ny_n, z, u)}{F_p(Mt, z, u)}, \frac{F_p(Ny_n, z, u)}{F_p(Mt, z, u)}, 1, 1, 1, F_p(Mt, z, u), F_p(Mt, z, u)F_p(Ny_n, Mt, u) \right] \right\}$$

$$F_p(Mt, Ny_n, ku) \geq \varphi \left\{ \min \left[\frac{F_p(Ny_n, z, u)}{F_p(z, z, u)}, \frac{F_p(Ny_n, z, u)}{F_p(z, z, u)}, 1, 1, 1, F_p(z, z, u), F_p(z, z, u)F_p(Ny_n, z, u) \right] \right\},$$

if $Mt = z$.

$$F_p(Mt, Ny_n, ku) \geq \varphi \left\{ \min [F_p(Ny_n, z, u), F_p(Ny_n, z, u), 1, 1, 1, F_p(Ny_n, z, u)] \right\}, \text{ if } Mt = z.$$

$$F_p(Mt, Ny_n, ku) \geq \varphi \{F_p(Ny_n, z, u)\}, \text{ if } Mt = z$$

Since $\varphi(l) \geq l$, for all $l \in (0, 1]$, it is possible only if $F_p(Ny_n, z, u) = 1$

$$F_p(Mt, Ny_n, ku) \geq F_p(Ny_n, z, u), \text{ if } Mt = z$$

By using Lemma1, we get $t = z$. Therefore $Mt = Ut = z$

Since (M, U) is weakly compatible. Then $Mz = MUt = UMt = Uz = z$

- b) Now to prove that $p(Vz, Nz) = 0$, that is to show that $Nz = z$

Put $x = x_n, y = z$ in (iii) we get,

$$F_p(Mx_n, Nz, ku) \geq \varphi \left\{ \min \left[\frac{F_p(Ux_n, Vz, u)F_p(Nz, Ux_n, u)}{F_p(Mx_n, Vz, u)}, \frac{F_p(Ux_n, Vz, u)F_p(Nz, Vz, u)}{F_p(Mx_n, Vz, u)}, \frac{F_p(Ux_n, Vz, u)F_p(Nz, Vz, u)}{F_p(Nz, Ux_n, u)}, \right. \right. \\ \left. \left. \frac{F_p(Nz, Vz, u)F_p(Mx_n, Vz, u)}{F_p(Nz, Ux_n, u)}, \frac{F_p(Mx_n, Vz, u)F_p(Nz, Mx_n, u)}{F_p(Vz, Ux_n, u)} \right] \right\}$$

$$F_p(Mx_n, Nz, ku) \geq \varphi \left\{ \min \left[\frac{F_p(Ux_n, z, u)F_p(Nz, Ux_n, u)}{F_p(Mx_n, z, u)}, \frac{F_p(Ux_n, z, u)F_p(Nz, z, u)}{F_p(Mx_n, z, u)}, \frac{F_p(Ux_n, z, u)F_p(Nz, z, u)}{F_p(Nz, Ux_n, u)}, \right. \right. \\ \left. \left. \frac{F_p(Nz, z, u)F_p(Mx_n, z, u)}{F_p(Nz, Ux_n, u)}, \frac{F_p(Mx_n, z, u)F_p(Nz, Mx_n, u)}{F_p(z, Ux_n, u)} \right] \right\}$$

$$F_p(Mx_n, Nz, ku) \geq \varphi\{\min[\frac{F_p(z,z,u)F_p(Nz,z,u)}{F_p(Mx_n,z,u)}, \frac{F_p(z,z,u)F_p(Nz,z,u)}{F_p(Mx_n,z,u)}, \frac{F_p(Uz,z,u)F_p(Nz,z,u)}{F_p(Nz,z,u)},$$

$$\frac{F_p(Nz,z,u)F_p(Mx_n,z,u)}{F_p(Nz,z,u)}, \frac{F_p(Mx_n,z,u)F_p(Nz,Mx_n,u)}{F_p(z,z,u)}]\}$$

$$F_p(Mx_n, Nz, ku) \geq \varphi\{\min[\frac{F_p(z,z,u)F_p(Nz,z,u)}{F_p(z,z,u)}, \frac{F_p(z,z,u)F_p(Nz,z,u)}{F_p(z,z,u)}, \frac{F_p(Uz,z,u)F_p(Nz,z,u)}{F_p(Nz,z,u)},$$

$$\frac{F_p(Nz,z,u)F_p(z,z,u)}{F_p(Nz,z,u)}, \frac{F_p(z,z,u)F_p(Nz,z,u)}{F_p(z,z,u)}]\}$$

$$F_p(Mx_n, Nz, ku) \geq \varphi\{\min[F_p(Nz, z, u), F_p(Nz, z, u), F_p(Uz, z, u), 1, F_p(Nz, z, u)]\}$$

But $Uz = z$.

$$F_p(Mx_n, Nz, ku) \geq \varphi\{\min[F_p(Nz, z, u), F_p(Nz, z, u), F_p(z, z, u), 1, F_p(Nz, z, u)]\}$$

$$F_p(Mx_n, Nz, ku) \geq \varphi\{\min[F_p(Nz, z, u), F_p(Nz, z, u), 1, 1, F_p(Nz, z, u)]\}$$

$$F_p(Mx_n, Nz, ku) \geq \varphi\{F_p(Nz, z, u)\}$$

$$F_p(Mx_n, Nz, ku) \geq F_p(Nz, z, u)$$

Since $\varphi(l) \geq l$, for all $l \in (0,1]$, it is possible only if $F_p(Nz, z, u) = 1$

By using Lemma1, we get $Nz = z$. Therefore $Nz = Vz = z$

Since (N, V) is weakly compatible. Then $Nz = Vz = NVz = VNz = z$

c) Now to prove that $p(Mz, Uz) = 0$, that is to show that $Mz = z$.

Put $x = z, y = y_n$ in (iii) we get,

$$F_p(Mz, Ny_n, ku) \geq \varphi\{\min[\frac{F_p(Uz,Vy_n,u)F_p(Ny_n,Uz,u)}{F_p(Mz,Vy_n,u)}, \frac{F_p(Uz,Vy_n,u)F_p(Ny_n,Vy_n,u)}{F_p(Mz,Vy_n,u)}, \frac{F_p(Uz,Vy_n,u)F_p(Ny,Vy_n,u)}{F_p(Ny_n,Uz,u)},$$

$$\frac{F_p(Ny_n,Vy_n,u)F_p(Mz,Vy_n,u)}{F_p(Ny_n,Uz,u)}, \frac{F_p(Mz,Vy_n,u)F_p(Ny_n,Mz,u)}{F_p(Vy_n,Uz,u)}]\}$$

$$F_p(Mz, Ny_n, ku) \geq \varphi\{\min[\frac{F_p(z,z,u)F_p(Ny_n,z,u)}{F_p(Mz,z,u)}, \frac{F_p(z,z,u)F_p(Ny_n,z,u)}{F_p(Mz,z,u)}, \frac{F_p(z,z,u)F_p(Ny,z,u)}{F_p(Ny_n,z,u)},$$

$$\frac{F_p(Ny_n,z,u)F_p(Mz,z,u)}{F_p(Ny_n,z,u)}, \frac{F_p(Mz,z,u)F_p(Ny_n,Mz,u)}{F_p(z,z,u)}]\}$$

$$F_p(Mz, Ny_n, ku) \geq \varphi\{\min[\frac{1.F_p(Ny_n,z,u)}{F_p(Mz,z,u)}, \frac{1.F_p(Ny_n,z,u)}{F_p(Mz,z,u)}, 1, 1, 1.F_p(Mz, z, u), \frac{F_p(Mz,z,u)F_p(Ny_n,Mz,u)}{1}]\}$$

$$F_p(Mz, Ny_n, ku) \geq \varphi\{\min[\frac{F_p(Ny_n,z,u)}{F_p(Mz,z,u)}, \frac{F_p(Ny_n,z,u)}{F_p(Mz,z,u)}, 1, 1.F_p(Mz, z, u), F_p(Mz, z, u)F_p(Ny_n, Mz, u)]\}$$

$$F_p(Mz, Ny_n, ku) \geq \varphi\{\min[\frac{F_p(Ny_n,z,u)}{F_p(z,z,u)}, \frac{F_p(Ny_n,z,u)}{F_p(z,z,u)}, 1, 1.F_p(z, z, u), F_p(z, z, u)F_p(Ny_n, z, u)]\}, \text{ if } Mz = z.$$

$$F_p(Mz, Ny_n, ku) \geq \varphi\{\min[F_p(Ny_n, z, u), F_p(Ny_n, z, u), 1, 1, F_p(Ny_n, z, u)]\}, \text{ if } Mz = z.$$

$$F_p(Mz, Ny_n, ku) \geq \varphi\{F_p(Ny_n, z, u)\}, \text{ if } Mz = z$$

Since $\varphi(l) \geq l$, for all $l \in (0,1]$, it is possible only if $F_p(Ny_n, z, u) = 1$

$F_p(Mz, Ny_n, ku) \geq F_p(Ny_n, z, u)$, if $Mz = z$

By using Lemma1, we get $z = z$. Therefore $Mz = Uz = z$

Since (M, U) is weakly compatible. Then $Mz = MUz = UMz = Uz = z$

d) Now to prove that $p(Vt, Nt) = 0$, that is to show that $Nt = z$

Put $x = x_n, y = t$ in (iii) we get,

$$F_p(Mx_n, Nt, ku) \geq \varphi\{\min[\frac{F_p(Ux_n,Vt,u)F_p(Nt,Ux_n,u)}{F_p(Mx_n,Vt,u)}, \frac{F_p(Ux_n,Vt,u)F_p(Nt,Vt,u)}{F_p(Mx_n,Vt,u)}, \frac{F_p(Ux_n,Vt,u)F_p(Nt,Vt,u)}{F_p(Nt,Ux_n,u)},$$

$$\frac{F_p(Nt,Vt,u)F_p(Mx_n,Vt,u)}{F_p(Nt,Ux_n,u)}, \frac{F_p(Mx_n,Vt,u)F_p(Nt,Mx_n,u)}{F_p(Vt,Ux_n,u)}]\}$$

$$F_p(Mx_n, Nt, ku) \geq \varphi\{\min[\frac{F_p(Ux_n, z, u)F_p(Nt, Ux_n, u)}{F_p(Mx_n, z, u)}, \frac{F_p(Ux_n, z, u)F_p(Nt, z, u)}{F_p(Mx_n, z, u)}, \frac{F_p(Ux_n, z, u)F_p(Nt, z, u)}{F_p(Nz, Ux_n, u)},$$

$$\frac{F_p(Nz, z, u)F_p(Mx_n, z, u)}{F_p(Nz, Ux_n, u)}, \frac{F_p(Mx_n, z, u)F_p(Nz, Mx_n, u)}{F_p(z, Ux_n, u)}]\}$$

$$F_p(Mx_n, Nt, ku) \geq \varphi\{\min[\frac{F_p(z, z, u)F_p(Nt, z, u)}{F_p(Mx_n, z, u)}, \frac{F_p(z, z, u)F_p(Nt, z, u)}{F_p(Mx_n, z, u)}, \frac{F_p(Uz, z, u)F_p(Nt, z, u)}{F_p(Nt, z, u)},$$

$$\frac{F_p(Nt, z, u)F_p(Mx_n, z, u)}{F_p(Nt, z, u)}, \frac{F_p(Mx_n, z, u)F_p(Nt, Mx_n, u)}{F_p(z, z, u)}]\}$$

$$F_p(Mx_n, Nt, ku) \geq \varphi\{\min[\frac{F_p(z, z, u)F_p(Nt, z, u)}{F_p(z, z, u)}, \frac{F_p(z, z, u)F_p(Nt, z, u)}{F_p(z, z, u)}, \frac{F_p(Uz, z, u)F_p(Nt, z, u)}{F_p(Nt, z, u)},$$

$$\frac{F_p(Nt, z, u)F_p(z, z, u)}{F_p(Nt, z, u)}, \frac{F_p(z, z, u)F_p(Nt, z, u)}{F_p(z, z, u)}]\}$$

$$F_p(Mx_n, Nt, ku) \geq \varphi\{\min[F_p(Nt, z, u), F_p(Nt, z, u), F_p(Uz, z, u), 1, F_p(Nt, z, u)]\}$$

But $Uz = z$.

$$F_p(Mx_n, Nt, ku) \geq \varphi\{\min[F_p(Nt, z, u), F_p(Nt, z, u), F_p(z, z, u), 1, F_p(Nt, z, u)]\}$$

$$F_p(Mx_n, Nt, ku) \geq \varphi\{\min[F_p(Nt, z, u), F_p(Nt, z, u), 1, 1, F_p(Nt, z, u)]\}$$

$$F_p(Mx_n, Nt, ku) \geq \varphi\{F_p(Nt, z, u)\}$$

$$F_p(Mx_n, Nt, ku) \geq \varphi\{F_p(Mx_n, z, u)\}, \text{ if } Nt = z$$

Since $\varphi(l) \geq l$, for all $l \in (0, 1]$, it is possible only if $F_p(Mx_n, z, u) = 1$

$F_p(Mx_n, Nt, ku) \geq F_p(Mx_n, z, u)$, if $Nt = z$. By using Lemma1, we get $Nt = z$.

Therefore $Nt = Vt = z$

Since (N, V) is weakly compatible. Then $Nt = NVt = Vnt = Vt = z$

Thus, $Mz = Uz = Nz = Vz = z$. Therefore M, N, U, V have a common fixed point.

Uniqueness:

Suppose t & z are two common fixed point of M, N, U, V with $t \neq z$, Then (iii) becomes

$$F_p(Mt, Nz, ku) \geq \varphi\{\min[\frac{F_p(Ut, Vz, u)F_p(Nz, Ut, u)}{F_p(Mt, Vz, u)}, \frac{F_p(Ut, Vz, u)F_p(Nz, Vz, u)}{F_p(Mt, Vz, u)}, \frac{F_p(Ut, Vz, u)F_p(Nz, Vz, u)}{F_p(Nz, Ut, u)},$$

$$\frac{F_p(Nz, Vz, u)F_p(Mt, Vz, u)}{F_p(Nz, Ut, u)}, \frac{F_p(Mt, Vz, u)F_p(Nz, Mt, u)}{F_p(Vz, Ut, u)}]\}$$

Put $Mt = Ut = t$, $Nz = Vz = z$

$$F_p(t, z, ku) \geq \varphi\{\min[\frac{F_p(t, z, u)F_p(z, t, u)}{F_p(t, z, u)}, \frac{F_p(t, z, u)F_p(z, z, u)}{F_p(t, z, u)}, \frac{F_p(t, z, u)F_p(z, z, u)}{F_p(z, t, u)},$$

$$\frac{F_p(z, z, u)F_p(t, z, u)}{F_p(z, t, u)}, \frac{F_p(t, z, u)F_p(z, t, u)}{F_p(z, t, u)}]\}$$

$$F_p(t, z, ku) \geq \varphi\{\min[F_p(z, t, u), 1, 1, 1, F_p(t, z, u)]\}$$

$$F_p(t, z, ku) \geq \varphi\{F_p(z, t, u)\}, \text{ Since } \varphi(l) \geq l, \text{ for all } l \in (0, 1], \text{ it is possible only if } F_p(t, z, u) = 1$$

$$F_p(t, z, ku) \geq F_p(t, z, u)$$

That is $t = z$. Hence Proved.

Generalization Theorem: Let $M = \{M_1, M_2, \dots, M_m\}$, $N = \{N_1, N_2, \dots, N_n\}$,

$U = \{U_1, U_2, \dots, U_u\}$, $V = \{V_1, V_2, \dots, V_v\}$ be a self-mapping of a Fuzzy PMS $(\mathbb{X}, F_p, \odot)$ with

$a \odot b = a \odot b$ satisfying following conditions:

- 1) (M, U) and (N, V) satisfy E. A. property.
- 2) (M, U) and (N, V) are weakly compatible.
- 3) Then their exist $k \in (0, 1)$, for all $x, y, z \in \mathbb{X}$ and $u > 0$.

$$F_p(Mx, Ny, ku) \geq [F_p(Ux, Vy, u) \odot \frac{F_p(Ny, Vy, u)}{F_p(Ny, Mx, u)} \odot (F_p(Ux, Vy, u). F_p(Ny, Vy, u)) \odot$$

$$\frac{F_p(Ny, Vy, u)}{F_p(Ny, Ux, u)} \odot \frac{F_p(Ny, Mx, u)}{F_p(Ny, Vy, u)}$$

Then M, N, U, V have a common fixed point.

4. Conclusion

This paper establishes a common fixed point theorem for two weakly compatible mappings in FPMS with the E.A. property. Our result generalizes existing fixed point theorems in FPMS by relaxing the compatibility condition and introducing the E.A. property. The E.A. property allows for more flexibility in modeling real-world problems with uncertain or imprecise data. The main result provides a sufficient condition for the existence of a common fixed point, which is a crucial concept in fixed point theory with numerous applications in various fields. The theorem's proof demonstrates the significance of the E.A. property in establishing the convergence of sequences in FPMS.

Real life Example in computer is Image Compression with Fuzzy Similarity.

Imagine a situation where we want to compress an image while maintaining a degree of "fuzzy" similarity. Instead of exact pixel-by-pixel comparison, we might want to consider how similar groups of pixels are, and how that similarity changes as we compress.

A fuzzy partial metric could model this similarity. Instead of a crisp distance, the distance between two groups of pixels could be fuzzy – representing a range of possible similarities. This fuzzy partial metric would need to satisfy specific axioms of partial metric spaces. The E-A property would be a condition ensuring that the compression algorithm converges to a "best" approximation of the original image within the constraints of the fuzzy similarity. The fixed point would represent the compressed image. It's the image where the compression process, using the fuzzy partial metric, doesn't lead to a further approximation. It's the "most optimal" image representation according to the specific fuzzy similarity.

Future scope

- Exploring the relationships between the E.A. property and other fuzzy partial metric space properties.
- Investigating the applications of this theorem in image processing, machine learning, and optimization problems.
- Extending the result to more general fuzzy partial metric spaces or other fuzzy structures.

References

- [1] Amer F.J., 2016. Fuzzy Partial Metric Spaces. In: Anastassiou, G., Duman, O. Computational Analysis. Springer Proceedings in Mathematics & Statistics, 155, 153-161. https://doi.org/10.1007/978-3-319-28443-9_11.
- [2] Ashlekha Dubey, Firdous Qureshi, 2013. Fixed Point Theorem in Fuzzy Metric Space with E.A. Property. Mathematical theory and Modeling, 3, 54-57.
- [3] Aygün H., Güner E. Miñana J.J., Valero O., 2022. Fuzzy Partial Metric Spaces and Fixed Point Theorems. Mathematics, 10, 3092. <https://doi.org/10.3390/math10173092>.
- [4] Başak Aldemir, Elif Güner, 2020. Some Fixed Point Theorems in Partial Fuzzy Metric Spaces. Journal of the Institute of Science and Technology, 10(4), 2889-2900. <https://doi.org/10.21597/jist.716207>.
- [5] Bonuga V., Shrinivas v., 2021. Some Extraction of Fixed point theorems using various E.A. properties. Communication in Mathematics & Applications, Vol.12, No.3, 445-456.
- [6] Elif Guner, Halis Aygun, 2022. A new approach to fuzzy partial metric spaces. Hacet. J.Math. Stat., 51(6), 1563-1576. <https://doi.org/10.15672/hujms.1115381>.
- [7] George A. Veeramani P, 1994. On Some Results in Fuzzy Metric Space. Fuzzy Sets and Systems, 90, 365-368. [https://doi.org/10.1016/S0165-0114\(96\)00207-2](https://doi.org/10.1016/S0165-0114(96)00207-2).
- [8] Gurucharan singh, 2022. Common Fixed point theorems in S-metric spaces using E.A. property and an implicit relation. Annals of Mathematics and computer science, Vol.10,51-64.
- [9] I. Kramosil and J. Michalek, 1975. Fuzzy Metric and statistical metric spaces. Kubernetika, 11, 336-344.
- [10] Kanhaiya Jha, Vyomesh pant, 2017. Some common fixed point Theorems in Fuzzy Metric Space with property (E.A.). Thai Journal of Mathematics volume15, 217:226.
- [11] K. Wadhwa, Dubey, Reena Jain, 2011. Impact of E. A. Property on common Fixed Point Theorems in Fuzzy metric Spaces. Journal of Advanced studies in Topology, Volume3, No.1, 52-59. <https://doi.org/10.20454/jast.2012.235>.
- [12] K. Wadhwa, Dubey, 2017. Common Fixed Point Theorems using E. A. Property in Fuzzy Metric spaces. International Journal of Mathematics, 8(6), 204-210.
- [13] K. Sisodia, A. Maharshi, 2024. Weak compatibility and E.A. property for a common fixed point theorem in Menger space. Rajashri Janak university Research journal, Vol.2, No.1&2, 25-30. <https://doi.org/10.3126/rjur.v2i1-2.72267>.
- [14] Madhushrivastava, K. Qureshi, A. D. Singh, 2016. Common Fixed Point Theorem in Fuzzy Metric Spaces Under E. A. Like Property. Pure and Applied Mathematics Journal, 5(4), 141-144. <https://doi.org/10.11648/j.pamj.20160504.18>.
- [15] Matthews S, 1994. Partial Metric Topology. Annals of the New York Academy of Sciences- Paper Edition, 728, 183-197. <https://doi.org/10.1111/j.1749-6632.1994.tb44144.x>.
- [16] P. Balsubramiam, R. P. pant, 2002. Common fixed points of four mapping in fuzzy metric space. Journal of Fuzzy Maths 10(2), 379-384.
- [17] Pingale M., Pathak R., 2023. Common Fixed point theorems for four maps using b-E. A. property. AIPconference proceedings, 2735(1), ID 040013, 7. <https://doi.org/10.1063/5.0140631>.
- [18] Rakhi Namdev, Rashmi Tiwari, Umashankar Singh, Ramakant Bhardwaj, 2024. Common Fixed-Point Theorems for Rational Expression in Fuzzy Metric Space using E.A like Property. African Journal of Biological Sciences 6(10), 6224-6237.
- [19] R. Pandya, A. Pariya, 2022. Common Fixed point theorems satisfying E.A. property in complex valued metric spaces. International journal of Mathematics Trends & Technology, 68(5), 6-10. <https://doi.org/10.14445/22315373/IJMTT-V68I5P502>.
- [20] Sonam Soni, 2025. Fixed points in Partial metric & Fuzzy metric spaces: Applications in computational Mathematics. Innovation and Integrative Research Center Journal, Vol-3, Issue-4, 207-220.
- [21] Xing Jin, Yongjie Piao, 2015. Common fixed points for two contractive mappings of integral in metric spaces. Applied Mathematics, 6(6), 1009-1016. <https://doi.org/10.4236/am.2015.66093>.
- [22] Subhani, M. Vijaya kumar, 2018. Common Fixed point Results in Fuzzy MS and Occasionally weakly compatible Mappings. IOSR Journal of Mathematics, 14(2), 51-54.
- [23] Talat Nazir and Mujahid Abbas, 2014. Common fixed points of two pairs of mappings satisfying (E.A.)-property in partial metric spaces. Journal of Inequalities and applications, 237, 1-12. <https://doi.org/10.1186/1029-242X-2014-237>.
- [24] Zadeh, LA, 1965. Fuzzy sets. Information and Control, 8(3), 338-353. [https://doi.org/10.1016/S0019-9958\(65\)90241-X](https://doi.org/10.1016/S0019-9958(65)90241-X).