

# The new solitary solutions of the foam drainage & (2+1) dimensional breaking soliton equations

Emad H. M. Zahran<sup>1\*</sup>, Maha S. M. Shehat<sup>2</sup>

<sup>1</sup>Department of Mathematical and physical Engineering, Benha University, Faculty of Engineering, Shubra, Egypt

<sup>2</sup>Department of Mathematics, Zagazig University, Faculty of Science, Zagazig, Egypt

\*E-mail: [e\\_h\\_zahran@hotmail.com](mailto:e_h_zahran@hotmail.com)

## Abstract

In this study, the modified extended tanh-function method is handling to obtain many new solitary wave solutions of two important models in nonlinear physics. The first one is the foam drainage equation which is a simple model for describing the flow of liquid through channels and nodes between the bubbles, driven by gravity and capillarity. The second is (2+1)-dimensional breaking soliton equation which describe the interaction of a Riemann wave propagating along the y-axis with along the x-axis. The obtained results are compared with that obtained in previous work.

**Keywords:** Foam Drainage Equation; the (2+1)-Dimensional Breaking Soliton Equation; the Modified Extended Tanh-Function Method; Riccati Equation; Travelling Wave Solution.

## 1. Introduction

In fluid mechanics trooping pockets of gas in liquid will form what is mean by the foam whose influence in the properties of liquid is important. The foam drainage in fluid mechanics according to the accessible configuration is modeled analytically by one-dimensional nonlinear partial differential equation. Also the (2+1)-dimensional breaking soliton equation which contributes to understanding most experiments and complex phenomena to many nonlinear branches of physics These two important models which used to represent the wave motion in different branches of nonlinear physics (fluid, plasma,...,etc) are investigated mathematically. The modified extended tanh-function method is a good technique to obtain the exact travelling wave solutions (which contain some parameters) for these equations if these parameters take definite values the solitary wave solution can be derived from it Several methods are applied successively to solve the nonlinear partial equations through many authors [1-22] such as extended Jacobi elliptic function method, the F-expansion method, the modified simple equation-method, the modified simple equation-method, the  $(\frac{G'}{G})$ -expansion method ,the modified  $(\frac{G'}{G}, \frac{1}{G})$ -expansion method , the  $\exp(-\phi(\zeta))$ -expansion method,modified extended  $\exp(-\phi(\zeta))$ -expansion method, theRiccati-Bernoulli Sub-ODE method, exp-function method, the tanh-function method, the extended tanh-function method, modified extended tanh-function method and so on. The main idea of this method is finding the exact solution of any models which can be expressed by a polynomial of  $\phi(\zeta)$  which satisfies the Riccati differential equation  $\phi' = b + \phi^2(\zeta)$  where  $\zeta = x - ct$  while b, c are arbitrary constants to be determined later .The degree of the polynomial can be calculated by the homogenous balance between the highest order derivative term and the nonlinear term. Equating the coefficients of the different power of  $\phi(\zeta)$  to zero , we get the system of algebraic

equations .The constants of the polynomial can be determined by solving this system of algebraic equation by Maple or any other computer program. The remainder of this paper is organized as follows: In section 2, we provide a description of the modified extended tanh-function method .In section 3 we use this method to find the exact and the solitary wave solutions of the nonlinear evolution equations given above. In section 4 our conclusion is presented.

## 2. Description of the method

Consider the following nonlinear evolution equation

$$f(u, u_x, u_{xx}, u_{xxx}, \dots) = 0 \quad (1)$$

Where f is a polynomial in u(x, t) and its partial derivatives in which the highest order derivatives and nonlinear terms are involved. In the following, we give the main steps of this method. Step1. We use the wave transformation,

$$u(x, t) = u(\zeta), \quad \zeta = x - ct \text{ and } c \text{ is cons.} \quad (2)$$

To reduce Eq. (1) to the following ODE:

$$p(u, u', u'', \dots) = 0 \quad (3)$$

Where P is a polynomial in  $u(\xi)$  and its total derivative, while  $(\xi = \frac{d'}{d\zeta})$

Step2. Suppose the solution of Eq. (3) has the form:

$$u(\xi) = a_0 + \sum_{i=1}^m (a_i \phi_i + \frac{b_i}{\phi_i}) \quad (4)$$

Where  $a_i, b_i$  are constants to be determined, such that  $a_m \neq 0$  or  $b_m \neq 0$  and  $\phi$  satisfies the

$$\text{Riccati equation } \phi' = b + \phi^2 \quad (5)$$

Eq. (5) admits several types of solutions according to the value of the constant  $b$  namely:

Case1. If  $b < 0$ , then

$$\phi = -\sqrt{-b} \tanh(\sqrt{-b}\zeta), \text{ or } \phi = -\sqrt{-b} \coth(\sqrt{-b}\zeta). \quad (6)$$

Case2. If  $b > 0$ , then

$$\phi = \sqrt{b} \tan(\sqrt{b}\zeta), \text{ or } \phi = -\sqrt{b} \cot(\sqrt{b}\zeta). \quad (7)$$

Case3. If  $b = 0$ , then

$$\phi = -\frac{1}{\zeta} \quad (8)$$

Step3. Determine the positive integer  $m$  in Eq. (4) by balancing the highest order derivative term and the nonlinear term Step4. Substitute Eq.(4) along Eq.(5) into Eq.(3) and collecting all the terms of the same power of  $\phi^i$ ,  $i = 0, \pm 1, \pm 2, \pm 3, \dots$  and by equating the coefficient of different power of  $\phi$  to zero, we obtain a system of algebraic equations, which can be solved by Maple or Mathematica to get the values of  $a_i, b_i$ .

Step5. Substituting these values and the solutions of Eq. (5) into Eq. (4) we obtain the exact solutions of Eq. (1).

### 3. Application

Example (1): The foam drainage equation [2]

The drainage of foam liquid involves the interplay of gravity, surface tension and viscous force. According to the accessible configuration the dynamic of the foam drainage in fluid mechanics phenomena are modeled analytically using one -dimensional nonlinear partial differential equation namely the foam drainage equation.

Here, we will apply the modified extended tanh -function method described in sec.2 to find the exact traveling wave solutions and then the solitary wave solutions for the foam drainage equation.

Consider the foam drainage equation [28]

$$u_t + (u^2 - \frac{\sqrt{u}}{2} u_x)_x = 0 \quad (9)$$

Where  $x$  and  $t$  scaled position and time coordinates, respectively. Foam is central to a number of everyday activities, both natural and industrial. As such foam has been of great interest for academic research. In the process industries, foam can be a desirable and even essential element of a process. An example is in the case of the forth flotation separation of minerals and coal [28]. Foaming occurs in many distillation and absorption processes. Foams are very important in many technological processes and applications. Their properties are subject to intensive investigational efforts from both practical developers and scientific researchers [28]. In this study; we show the effectiveness and convenience of the method by obtaining the exact solution of Eq. (9).

Using the wave variable  $\zeta = k(x + ct)$ , Eq. (9) is carried to an ODE

$$cku' + k(u^2 - \frac{k}{2}\sqrt{uu'})' = 0 \quad (10)$$

Integrating Eq. (10) with respect to  $\zeta$  and considering the zero constants for integration we obtain.

$$cku + k(u^2 - \frac{k}{2}\sqrt{uu'}) = 0 \quad (11)$$

Then we use the transformation

$$u(\zeta) = v^2(\zeta), \quad (12)$$

That will convert Eq. (11) to

$$kcv^2 + kv^4 - k^{\frac{3}{2}}v^{\frac{3}{2}}v' = 0 \quad (13)$$

Or equivalently

$$c + v^2 - kv' = 0 \quad (14)$$

Balancing  $v', v^2$  yields  $m+1 = 2m \Rightarrow m = \text{one}$ . Consequently, we obtain the solution

$$u(\zeta) = a_0 + a_1\phi(\zeta) + \frac{b_1}{\phi(\zeta)}, \quad (15)$$

Sub. , about  $v$  and  $v^2$  at Eq. (14), and equating different power of  $\phi(\zeta)$  to zero, we obtain algebraic system of equation

$$\begin{aligned} a_1^2 - ka_1 &= 0 \\ 2a_1a_0 &= 0 \\ b_1^2 + kbb_1 &= 0 \\ 2a_1b_1 &= 0 \\ c + a_0^2 + 2ab_1 - kab_1 - kab + kb_1 &= 0 \end{aligned} \quad (16)$$

Solving this system by Maple or Mathematica, we get

$$\begin{aligned} (1) \quad a_0 &= 0, \quad a_1 = k, \quad b_1 = -\frac{c}{4k}, \quad b = \frac{c}{4k^2} \\ (2) \quad a_0 &= 0, \quad a_1 = k, \quad b_1 = 0, \quad b = \frac{c}{a_1^2} \\ (3) \quad a_0 &= 0, \quad a_1 = 0, \quad b_1 = -\frac{c}{k}, \quad b = \frac{c}{k^2} \end{aligned}$$

Case (1):

$$\text{For, } a_0 = 0, \quad a_1 = k, \quad b_1 = -\frac{c}{4k}, \quad b = \frac{c}{4k^2}$$

The solution is

$$v(\zeta) = k\phi - \frac{c}{4k} \left(\frac{1}{\phi}\right)$$

When  $b < 0$ , we get the solitary wave solution

$$v(\zeta) = \sqrt{\frac{-c}{4k^2}} \tanh\left(\sqrt{\frac{-c}{4k^2}}\zeta\right) - \frac{c}{4k} \sqrt{\frac{-c}{4k^2}} \cot\left(\sqrt{\frac{-c}{4k^2}}\zeta\right) \quad (17)$$

$$v(\zeta) = \sqrt{\frac{-c}{4k^2}} \coth\left(\sqrt{\frac{-c}{4k^2}}\zeta\right) - \frac{c}{4k} \sqrt{\frac{-c}{4k^2}} \tanh\left(\sqrt{\frac{-c}{4k^2}}\zeta\right) \quad (18)$$

When  $b > 0$ , we get the solitary wave solution

$$v(\zeta) = \sqrt{\frac{c}{4k^2}} \tan\left(\sqrt{\frac{c}{4k^2}}\zeta\right) - \frac{c}{4k} \sqrt{\frac{c}{4k^2}} \cot\left(\sqrt{\frac{c}{4k^2}}\zeta\right) \quad (19)$$

$$v(\zeta) = \sqrt{\frac{c}{4k^2}} \cot\left(\sqrt{\frac{c}{4k^2}} \zeta\right) - \frac{c}{4k} \sqrt{\frac{c}{4k^2}} \tan\left(\sqrt{\frac{c}{4k^2}} \zeta\right) \tag{20}$$

When  $b=0$ , we get the trivial solution.

Case (2):

$$(2) a_0 = 0, a_1 = k, b_1 = 0, b = \frac{c}{a_1^2}$$

The solution is  $v(\zeta) = k \phi$

When  $b < 0$ , we obtain the solitary wave solution

$$v(\zeta) = -k \sqrt{\frac{-c}{k^2}} \tanh\left(\sqrt{\frac{-c}{4k^2}} \zeta\right) \tag{21}$$

$$v(\zeta) = -k \sqrt{\frac{-c}{k^2}} \coth\left(\sqrt{\frac{-c}{k^2}} \zeta\right) \tag{22}$$

When  $b > 0$ , we obtain the solitary wave solution

$$v(\zeta) = k \sqrt{\frac{c}{k^2}} \tan\left(\sqrt{\frac{c}{k^2}} \zeta\right) \tag{23}$$

$$v(\zeta) = k \sqrt{\frac{c}{k^2}} \cot\left(\sqrt{\frac{c}{k^2}} \zeta\right) \tag{24}$$

When  $b=0$ , we get the solitary wave solution.

$$v(\zeta) = -\frac{k}{\zeta} \tag{25}$$

Case (3)

$$(3) a_0 = 0, a_1 = 0, b_1 = \frac{-c}{k}, b = \frac{c}{k^2}$$

The solution is  $v(\zeta) = bk / \phi$

When  $b < 0$ , we obtain the solitary wave solution

$$v(\zeta) = \frac{c}{k} \sqrt{\frac{-c}{k^2}} \tanh\left(\sqrt{\frac{-c}{k^2}} \zeta\right) \tag{26}$$

$$v(\zeta) = \frac{c}{k} \sqrt{\frac{-c}{k^2}} \coth\left(\sqrt{\frac{-c}{k^2}} \zeta\right) \tag{27}$$


When  $b > 0$ , we obtain

$$v(\zeta) = \frac{-c}{k} \sqrt{\frac{c}{k^2}} \tan\left(\sqrt{\frac{c}{k^2}} \zeta\right) \tag{28}$$


$$v(\zeta) = \frac{-c}{k} \sqrt{\frac{c}{k^2}} \cot\left(\sqrt{\frac{c}{k^2}} \zeta\right) \tag{29}$$

When  $b=0$ , we obtain the trivial solution


Eq. (17).




Eq. (19).




Eq. (21).




Eq. (23).




Eq. (25).



Eq. (26).



Eq. (28).



Example (2): The (2+1)-Dimensional Breaking Soliton Equation [29].

Let us consider the (2+1)-dimensional breaking soliton equation,

$$u_t + \alpha u_{xy} + 4\alpha u v_x + 4\alpha u_x v = 0$$

$$u_y = v_x \tag{30}$$

Where  $\alpha$  is a well-known constant, physically this equation describes the (2+1)-dimensional interaction of a Riemann wave propagating along the y-axis with along wave along the x-axis. Many authors [30 - 33] study this equation by different manner to get soliton-like solutions, a class of periodic wave solutions, and two classes of exact solutions.

According to our proposed method, let us introduce the transformation  $\zeta = x + y - ct$ .

Sub.at (30) we get,

$$-cu' + \alpha u''' + 4\alpha uv' + 4\alpha u'v = 0$$

$$u' = v' \tag{31}$$

Integrating the second part of Eq. (31) with taking the constant of integration in account,

Sub.at the first part of Eq. (31), we obtain

$$-cu' + \alpha u''' + 4\alpha uv' + 4\alpha u'(u + c) = 0 \tag{32}$$

Integrating the last Eq., we get

$$(4\alpha c_1 - c)u + \alpha u''' + 4\alpha u^2 = 0 \tag{33}$$

Balancing the nonlinear term with the highest order derivative term, we find  $m+1 = 2m \Rightarrow m=2$ .

Consequently according to the constructed method the solution is

$$u(\zeta) = a_0 + a_1\varphi(\zeta) + \frac{b_1}{\varphi(\zeta)} + a_2\varphi^2(\zeta) + \frac{b_2}{\varphi^2(\zeta)}, \tag{34}$$

Sub.,about  $u, u'$  and  $u''$  at Eq.(33) and equating the coefficients of the different powers of  $\varphi$  to zero, we get the system of algebraic equations namely,

$$\begin{aligned} 6\alpha a_2 + 4\alpha a_1^2 &= 0 \\ 2\alpha a_1 + 8\alpha a_1 a_2 &= 0 \\ 6\alpha a_1 + 4\alpha a_2^2 &= 0 \end{aligned} \tag{35}$$

Solving this system of algebraic equations by Maple or any computer program, we get the case

$$a_0 = \left(\frac{c}{16\alpha} - \frac{c_1}{4}\right), a_1 = 0, a_2 = -3/2, b = a_0, b_1 = 0, b_2 = \frac{-3a_0^2}{2} \tag{36}$$

The solution is

$$u(\zeta) = \left(\frac{c}{16\alpha} - \frac{c_1}{4}\right) - (3/2)\varphi^2(\zeta) - \frac{3a_0^2}{2} \frac{1}{\varphi^2(\zeta)} \tag{37}$$

When  $b < 0$

$$u(\zeta) = \left(\frac{c}{16\alpha} - \frac{c_1}{4}\right) - \frac{3}{2} \left(\frac{c}{16\alpha} - \frac{c_1}{4}\right) \tanh^2 \left( \sqrt{\frac{c}{4} - \frac{c_1}{16\alpha}} \zeta \right) - \frac{3}{2} \left(\frac{c}{16\alpha} - \frac{c_1}{4}\right)^2 \coth^2 \left( \sqrt{\frac{c}{4} - \frac{c_1}{16\alpha}} \zeta \right) \tag{38}$$

$$u(\zeta) = \left(\frac{c}{16\alpha} - \frac{c_1}{4}\right) - \frac{3}{2} \left(\frac{c}{16\alpha} - \frac{c_1}{4}\right) \coth^2 \left( \sqrt{\frac{c}{4} - \frac{c_1}{16\alpha}} \zeta \right) - \frac{3}{2} \left(\frac{c}{16\alpha} - \frac{c_1}{4}\right)^2 \tanh^2 \left( \sqrt{\frac{c}{4} - \frac{c_1}{16\alpha}} \zeta \right)$$


When  $b > 0$ , we get the solitary wave solution

$$u(\zeta) = \left(\frac{c}{16\alpha} - \frac{c_1}{4}\right) - \frac{3}{2} \left(\frac{c}{16\alpha} - \frac{c_1}{4}\right) \tan^2 \left( \sqrt{\frac{c}{16\alpha} - \frac{c_1}{4}} \zeta \right) - \frac{3}{2} \left(\frac{c}{16\alpha} - \frac{c_1}{4}\right)^2 \cot^2 \left( \sqrt{\frac{c}{16\alpha} - \frac{c_1}{4}} \zeta \right) \tag{39}$$


$$u(\zeta) = \left(\frac{c}{16\alpha} - \frac{c_1}{4}\right) - \frac{3}{2} \left(\frac{c}{16\alpha} - \frac{c_1}{4}\right) \cot^2 \left( \sqrt{\frac{c}{16\alpha} - \frac{c_1}{4}} \zeta \right) - \frac{3}{2} \left(\frac{c}{16\alpha} - \frac{c_1}{4}\right)^2 \tan^2 \left( \sqrt{\frac{c}{16\alpha} - \frac{c_1}{4}} \zeta \right) \tag{40}$$

When  $b=0$ , we get


$$u(\zeta) = \left(\frac{c}{16\alpha} - \frac{c_1}{4}\right) - \frac{3}{2\zeta^2} - \frac{3}{2} \left(\frac{c}{16\alpha} - \frac{c_1}{4}\right)^2 \zeta^2 \tag{41}$$



Eq. (39).



Eq. (40).



#### 4. Competing interests

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sector. The author did not have any competing interests in this research.

#### 5. Authors contributions

All parts contained in this research carried out by the researcher through hard work and a review of the various references and contributions in the field of mathematics, physics and medical science. It represents the research work of the author.

#### Acknowledgment

The author is grateful to the anonymous referee for a careful checking of the details and for helpful comments that improved this research.

#### 6. Conclusion

In this study, after comparing our solutions (17-28) for the foam equation and (36) for the (2+1)-dimensional breaking soliton with that obtained by other authors, we find that many new solitary solutions according to the proposed method are introduced which are consistent in some cases but not in other. The proposed method give a large numbers of solitary solutions which give a good interpretation for the foam drainage and the (2+1)-dimensional breaking soliton phenomenon. Also, these large numbers of solitary solutions will help the Scientists to treat these phenomenon experimentally. And this well makes a forward effect on the application in fluid mechanics and plasma physics. Also, the obtained solutions can be considered as benchmarks against the numerical simulations in the fluid mechanics and plasma physics.

#### References

- [1] Younis, Muhammad. A new approach for the exact solutions of nonlinear equations of fractional order via modified simple equation method. *Applied Mathematics* 5.13 (2014):1927.
- [2] Emad H.M. Zahran., Mostafa M. A. Khater. The modified simple equation method and its applications for solving some nonlinear evolution equations in mathematical physics. *Jokull journal* 64.5 (2014): 297-312.
- [3] Wang, Mingliang, Xiangzheng Li, and Jinliang Zhang. The  $(G'/G)$ -expansion method and travelling wave solutions of nonlinear evolution equations in mathematical physics. *Letters A* 372.4 (2008): 417-423.
- [4] Ahmet Bekir, Ferat Uygun, Exact travelling wave solutions of some nonlinear evolution equations by using  $(G'/G)$ -expansion method. *Arab Journal of Mathematical Sciences* 18 (2012) 73-85.
- [5] Zhang, Sheng, Jing-Lin Tong, and Wei Wang. A generalized expansion method for the mKdV equation with variable coefficients, *Physics Letters a*, 37213 (2008): 2254-2257.
- [6] Fan Engui and Jian Zhang. Applications of the Jacobi elliptic function method to special type nonlinear equations. *Physics Letters a* 305.6 (2002): 383-392.
- [7] Emad H.M. Zahran. , Mostafa M. A. Khater. Exact Travelling wave solutions for the system of shallow water wave Equation Modified Liouville Equation using Extended Jacobi Elliptic function Expansion Method. *American Journal of computational Mathematics*, 4 455-463. (December 2014).
- [8] Maha S. M. Shehata, Extended Jacobi Elliptic function Expansion Method and its Applications for solving some Nonlinear Evolution Equations in Mathematical Physics. *International Journal of Computer Applications*, Vol.109-No.12. Jan. 2015.
- [9] Emad H. M. Zahran. Exact traveling wave solutions for Nano-solitons of Ionic waves propagation along Microtubules in living cells and Nano-Ionic currents of MTs, *World journal of Nano science and engineering*, 5,78-87,2015.
- [10] Emad H. M. Zahran .Exact traveling wave solutions for Nano-Ionic solitons and Nano-Ionic currents of MTs using  $\exp(-\varphi(\xi))$ -expansion method, *Advances in Nano particles*, 4,25-36, 2015.
- [11] Emad H. M. Zahran, Exact traveling wave solutions for nonlinear fractional partial differential equations arising in soliton using the  $\exp(-\varphi(\xi))$ -expansion method. *International journal of computer applications*. Vol.109, N.13, Jan. 2015.
- [12] Maha S. M. Shehata, The  $\exp(-\varphi(\xi))$ - Method and Its Applications for Solving some Nonlinear Evolution Equations in Mathematical Physics , *American Journal of Computational Mathematics*, 2015, 5,468-480.
- [13] Emad H. M. Zahran , Travelling wave solutions of Non Linear Evolution Equation Via Modified  $\exp(-\varphi(\xi))$ -Expansion method, *Journal of Computational and Theoretical Nanoscience*, Vol.12, 5716-5724, (2016).
- [14] Yang, Xiao-Feng, Zi-Chen Deng, and Yi Wei. A Riccati-Bernoulli sub-ODE method for nonlinear partial differential equations and its application. *Advances in Difference Equations*. 11 17, (2015).
- [15] Maha S. M. Shehata, A new solitary wave solution of the perturbed nonlinear Schrodinger equation using a Riccati Bernoulli Sub-ODE Method , *International Journal of Physical Sciences*, Vol.11 (6) pp.80-84, March.2016.
- [16] Bekir, Ahmet, and Ahmet Boz. Exact solutions for a class of nonlinear partial differential Equations using exp-function method. In-

- ternational Journal of Nonlinear Sciences and Numerical Simulation 8.4 (2007): 505-512.
- [17] Fan, Engui Extended tanh-function method and its applications to nonlinear equations. *Physics Letters. A.* 277.4(2000): 212-218.
- [18] Maha S. M. Shehata. Exact Traveling Wave Solutions for Nonlinear Evolutions Equation *Journal of Computational and Theoretical Nano science* (2016) Vol.13.No.1, pp.534-538.2016.
- [19] Elwakil, S. A., et al. Modified extended tanh-function method for solving nonlinear partial differential equations. *Physics Letters A* 299.2 (2002): 179-188.
- [20] Emad H.M Zahran, and Mostafa M. A. Khatter, New solitary wave solution of the generalized Hirota-Satsuma couple KdV system, *International Journal of scientific & Engineering Research*, Vol 6, Issue 8, August-2015 Wang, G. W.
- [21] Emad H.M Zahran, and Mostafa M. A. Khater. Modified extended tanh-function method and its applications to the Bogoyavlenskii equation. *Applied Mathematical Modeling* 40.3 (2016): 1769-1775.
- [22] Emad H.M Zahran, Dianchin Lu and Mostafa M. A. Khater. Solitary wave solution of the Benjamin-Bona-Mahoney-Burgers Equation with Dual Power-Law Nonlinearity, *Appl. Math., Inf. Sci.*, 11, No.5, 1-5 (2017)
- [23] S. D. Stoyanov, V. N. Paunov, E.S. Basheva, I. B. Ivanov, A. Mehreteab, G. Broze, Motion of the front between thick and thin film: Hydrodynamic theory and experiment with vertical foam films, *Langmuir B* (1997) 1400.
- [24] R. Hirta, Y. Ohta, Hierarchies of coupled soliton equations I, *J. Phys. Soc. Jpn.* 60. (1991) 798.
- [25] S. Zhang, New exact non-traveling wave and coefficient function solutions of the (2+1) dimensional breaking soliton equations, *Phys. Lett. A.* 368(2007) 470.
- [26] Y. Cheng, B. Li, Symbolic computation and construction of soliton-like solutions of the (2+1) dimensional breaking soliton equation, *Commun. Theor. Phys. (Beijing, China)* 40 (2003) 137.
- [27] Y. Z. Peng, New exact solutions for the (2+1) - dimensional breaking soliton equation. *Commun. Theor. Phys. (Beijing, China)* 43 (2005) 205.
- [28] Y. Z. Peng, E. V. Krishna, Two classes of new exact solutions to the (2+1) dimensional breaking soliton equation, *Commun. Theor. Phys. (Beijing, China)* 44 (2005) 807.