International Journal of Engineering & Technology, 7 (3.31) (2017) 160-162

International Journal of Engineering & Technology

Website: www.sciencepubco.com/index.php/IJET

Research paper

A Study on Fuzzy TΓ-Ideals in Ternary Γ-Semiring

K. Revathi[1, 2], D. Madhusudhana Rao[3]*, P. Sundarayya[4] andT. Satish[5]

[1]Research Scholar, Department of Mathematics, GITAM University, Vishkhapatnam, A.P. India.
[2]Asst. Professor, Department of Mathematics, Adikavi Nannaya University, Rajamandri, A.P.
[3] Associate Professor, Department of Mathematics, VSR & NVR College, Tenali, A.P. India.
[4] Asst. Professor, Department of Mathematics, GITAM University, Visakhapatnam, A.P. India.
[5] Asst. Professor, Department of Mathematics, S. R. K. R. Engineering College, Bhimavaram, A.P. India.
**Corresponding AuthorE-mail: dmrmaths@gmail.com , dmr04080@gmail.com*

Abstract

The concepts of fuzzy theory in TΓ-semiring in terms of fuzzy TΓ-ideals in TΓ-semirings are introduced and also we made a study on some properties of fuzzy TΓ-Ideals in TΓ-semiring.

Keywords: *TΓ-Semiring, TΓ-ideal, fuzzy TΓ-ideal, TΓ-subsemiring, regular TΓ-semiring.*

1. Introduction

The concept of TΓ-semiring was introduced by S. Lavanya, D. M. Rao and syam Julius Rajendra[2] In the year 2015, Later they [3], [4] studied about the characteristics of TΓ-ideals in TΓ-semirings. In the year 2005 T. K. Dutta and T. Chanda[1] investigated about fuzzy Γ-ideals in Γ-rings. In 2004, K. Syam Prasad and Bh. Satyanarayana [5] studied about prime fuzzy ideals in Γ-near rings. In this paper fuzzy concept in TΓ-semiring was introduced.

2. Preliminaries

Note 2.8: For preliminaries refer the references [2], [3].

3. Fuzzy TΓ-ideals:

Def 3.1: A fuzzy set ξ of a TΓ-semiring is said to be a ***fuzzy TΓ-sub semiring*** of Q if

(i) $\xi(u + v) \geq \min\{\xi(u), \xi(v)\}$

(ii) $\xi(u\gamma v\delta w) \geq \min\{\xi(u), \xi(v), \xi(w)\}$ $\forall$ $u, v, w \in M$ and $\gamma, \delta \in \Gamma$.

Ex: **3.2**: Let M be the set of rational numbers and Γ is the set of natural numbers. Define a mapping from M× Γ × M × Γ × M to M by usual addition and ternary multiplication defined by $a\alpha b\beta c$ = usual product of a, α, b, β, c; for $a, b, c \in Q$, $\alpha, \beta \in \Gamma$. Then M is a T Γ-semi ring. Define μ:M → [0, 1] as $\mu(x) = \begin{cases} 1 & \text{if } x \in Q \\ 0, & \text{otherwise} \end{cases}$

Then μ is a fuzzy T Γ-semi ring of M.

Example 3.4: Consider the set Z = { 0, 1, -1, 2, -2, ……} and Γ be the set of even natural numbers. Then with respect to usual addition Z and multiplication is infinite T Γ-semiring. Clearly 2Z is a proper Γ-subsemiring of Z. Define $\mu : z \to [0,1]$ by $\xi(x) = \begin{cases} 0.9 & \text{if } x \in 2Z \\ 0.8, & \text{if } x \in 2Z+1 \end{cases}$ It is easy to verify that ξ is a fuzzy T Γ-sub semi ring of the TΓ-semi ring Z.

Def 3.5: A fuzzy T Γ -sub semi ring μ of a TΓ-semi ring M is called ***improper*** if μ is constant on the T Γ-semi ring T, otherwise μ is termed as ***proper*.**

The 3.6 : Let M be a TΓ-semiring and F(M) be the set of all non-empty fuzzy subsets of TΓ-semiring M. If *e, f, g, h* ∈ F(M), then

(i) $e \cap h \cap (f \cup g) = (e \cap h \cap f) \cup (f \cap h \cap g)$

(ii) $(e \cup f) \cap g \cap h = (e \cap f \cap g) \cup (f \cap g \cap h)$

(iii) $g \cap (e \cup f) \cap h = (g \cap e \cap h) \cup (g \cap f \cap h)$

The 3.7: Suppose Q be a TΓ-semi ring and ϱ a fuzzy sub set of M. Then (i) $\varrho(d\chi m\psi k) = \min\{\varrho(d), \varrho(m), \varrho(k)\}$ $\forall$ *d, m, k* ∈ Q and $\chi, \psi \in \Gamma$ and $\varsigma'(d\chi m\psi k) = \vee\{\varsigma'(d), \varsigma'(m), \varsigma'(k)\}$ are equivalent $\forall$ *d, m, k* ∈ Q and $\chi, \psi \in \Gamma$.

***Proof*:** Assume that $\varrho(d\chi m\psi k) = \min\{\varrho(d), \varrho(m), \varrho(k)\}$ $\forall$ *d, m, k* ∈ Q and $\chi, \psi \in \Gamma$. We may assume that $\varrho(d\chi m\psi k) = \varrho(d)$. Then $\varrho(d) \leq \varrho(m)$ and $\varrho(d) \leq \varrho(k)$, so $\varsigma'(p\gamma f\delta k) = 1 - \varsigma(d\chi m\psi k) = 1 - \varsigma(d) = \varsigma'(d)$ and $\varsigma'(d) = 1 - \varsigma(d) \geq 1 - \varsigma(m) = \varsigma'(m)$ and $\varsigma'(m) = 1 - \varsigma(d) \geq 1 - \varsigma(k) = \varsigma'(k)$. Therefore, $\varsigma'(d\chi m\psi k) = \vee\{\varsigma'(d), \varsigma'(m), \varsigma'(k)\}$ $\forall$ *d, m, k* ∈ Q and $\chi, \psi \in \Gamma$.

Conversely, suppose that $\varsigma'(d\chi m\psi k) = \vee\{\varsigma'(d), \varsigma'(m), \varsigma'(k)\}$ $\forall$ *d, m, k* ∈ Q and $\chi, \psi \in \Gamma$. We may assume that $\varsigma'(d\chi m\psi k) = \varsigma'(m)$. Then $\varsigma'(d) \geq \varsigma'(m)$ and $\varsigma'(d) \geq \varsigma'(k)$ so 1 - $\varsigma(d\chi m\psi k)$= 1 - $\varsigma(d)$, 1 - $\varsigma(d) \geq$ 1 - $\varsigma(m)$ and 1 - $\varsigma(m) \geq$ 1 - $\varsigma(k)$. Thus $\varsigma(d\chi m\psi k) = \varsigma(d)$, $\varsigma(d) \leq \varsigma(m)$ and $\varsigma(d) \leq \varsigma(k)$. Hence $\varsigma(d\chi m\psi k) = \min\{\varrho(d), \varrho(m), \varrho(k)\}$ $\forall$ *d, m, k* ∈ Q and $\chi, \psi \in \Gamma$.

Copyright © 2018 Authors. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Th 3.8: Let R be a TΓ-semi ring and K $\subseteq$ R, K $\neq \emptyset$. Then K is a TΓ-sub semi ring of R iff the fuzzy subset π_K is a fuzzy TΓ-sub-semiring of R.

***Proof* :** Obviously, π_K is a fuzzy subset of R. Let $u, v, w \in R$ and $\gamma, \delta \in \Gamma$. If $u \notin K$ or $v \notin K$ or $w \notin K$, then $\pi_K(u) = 0$ or $\pi_K(v) = 0$ or $\pi_K(w) = 0$, then $\pi_K(u+v) \geq 0 = \min\{\pi_K(u), \pi_K(v)\}$ and $\pi_K(u\gamma v\delta w) \geq 0 = \min\{\pi_K(u), \pi_K(v), \pi_K(w)\}$. Let $u, v, w \in K$. Then $\pi_K(u) = \pi_K(v) = \pi_K(w) = 1$. Since $u+v \in K$, $u\gamma v\delta w \in K\Gamma K\Gamma K \subseteq K$, we have $\pi(u + v) = 1 \geq 1 = \min\{\pi_K(u), \pi_K(v)\}$ and $\pi_K(u\gamma v\delta w) = 1$. Thus $\pi_K(u\gamma v\delta w) = 1 \geq 1 = \min\{\pi_K(u), \pi_K(v), \pi_K(w)\}$. Therefore the fuzzy subset π_K is a fuzzy TΓ-sub-semi ring of R.

Conversely, suppose that $u, v, w \in K$ and $\gamma, \delta \in \Gamma$. Then $\pi_K(u) = \pi_K(v) = \pi_K(w) = 1$. Since π_K is a fuzzy TΓ-sub semiring of R, we have $\pi_K(u+v) \geq \min\{\pi_K(u), \pi_K(v)\} = \min\{1, 1\} = 1$. Thus $u + v \in K$ and $\pi_K(u\gamma v\delta w) \geq \min\{\pi_K(u), \pi_K(v), \pi_K(w)\} = \min\{1, 1, 1\} = 1$. Thus $\pi_K(u\gamma v\delta w) = 1$ and hence $u\gamma v\delta w \in K$. Therefore K is a TΓ-sub semi ring of R.

Def 3.9: A non-empty fuzzy sub-set ϱ of a TΓ-semiring Q is called a ***fuzzy L(La, R) TΓ-ideal*** or simply ***fuzzy left TΓ-ideal*** of Q if

(i) $\varrho(u + v) \geq \wedge\{\varrho(u), \varrho(v)\}$

(ii) $\varrho(u\gamma v\delta w) \geq \varrho(w)[\ \varrho(u\gamma v\delta w) \geq \varrho(v),\ \varrho(u\gamma v\delta w) \geq \varrho(u)]\ \forall$ $u,v, w \in Q, \forall \gamma, \delta \in \Gamma$.

Ex 3.10: Let Q = {0, *s*, *t*, *u*} and Γ be the non-empty set of binary operations such that $\alpha, \beta \in \Gamma$ is defined below:

+	0	*s*	*t*	*u*
0	0	*s*	*t*	*u*
s	*s*	*t*	*u*	*s*
t	*t*	*u*	*u*	*s*
u	*u*	*t*	*s*	*t*

α	*0*	*s*	*s*	*u*
0	0	0	0	0
s	*0*	*s*	*t*	*u*
t	0	0	0	0
u	*0*	*s*	*t*	*u*

β	*0*	*s*	*t*	*u*
0	0	0	0	0
s	*s*	*s*	*s*	*s*
t	0	0	0	*t*
u	*s*	*s*	*s*	*u*

Clearly Q is a TΓ-semi ring. Define a fuzzy subset π: Q $\rightarrow$ [0, 1] by $\pi(x) = \begin{cases} 1 \text{ if } x=0, a \\ 0, \text{ otherwise} \end{cases}$. Clearly, π is a fuzzy L-TΓ-ideal of Q.

Def 3.11: A non-empty fuzzy sub-set ϖ of a TΓ-semi ring Q is called a ***fuzzy TΓ-ideal*** of Q if

(i) $\varpi(f + g) \geq \min\{\pi(f), \pi(g)\}$

(ii) $\varpi(f\gamma g\delta h) \geq \varpi(f) \vee \varpi(g) \vee \varpi(h)$ for any $f, g, h \in Q$ and $\gamma, \delta \in \Gamma$.

Ex 3.12: Let the set of -ve integers with 0 be P and the set of negative even integers with 0 be Γ. Then P is a TΓ-semi ring if $u + v$, $u\gamma v\delta w$ as well as $\alpha u\beta v\gamma$ denote the usual multiplication of integers u, γ, v, δ, w as well as $\alpha, u, \beta, v, \gamma$ respectively where $u, v, w \in Q$ as well as $\alpha, \beta, \gamma, \delta \in \Gamma$. Let η be a fuzzy sub-set of Q, defined as follows:

$$\eta(m) = \begin{cases} 1, & \text{if } m = 0 \\ 0.1, & \text{if } m = -1, -2 \\ 0.2, & \text{if } m < -2 \end{cases}$$

Then η is a fuzzy TΓ-ideal of Q.

Ex 3.13: Let M = {0, *u*, *v*, *w*} and Γ = {α, β} be the non-empty set of binary operations defined below:

+	0	*u*	*v*	*w*
0	0	*u*	*v*	*w*
u	*u*	*v*	*w*	*u*
v	*v*	*w*	*w*	*u*
w	*w*	*v*	*u*	*v*

α	0	*u*	*v*	*w*
0	0	0	0	0
u	0	*v*	0	*u*
v	0	*v*	0	*w*
w	0	0	0	*v*

β	0	*u*	*v*	*w*
0	0	0	0	0
u	*u*	*u*	*u*	*u*
v	0	0	0	0
w	*u*	*u*	*u*	*w*

Obviously, M is a TΓ-semiring. Moreover the fuzzy set π: M $\rightarrow$ [0, 1] defined by $\pi(0) = 0.7$, $\pi(u) = 0.6$, $\pi(v) = \pi(w) = 0.3$ is a fuzzy TΓ-ideal of Q.

Th 3.14: Suppose $\{\pi_i : i \in I\}$ is a family of fuzzy L-(La, R) TΓ-ideals of TΓ-semi ring T, then $(\bigcap_{i\in I} \pi_i)(u)$ is a fuzzy L-(La, R, TΓ-ideal) TΓ-ideal of Q.

***Proof* :** Let $\{\pi_i : i \in I\}$ be a family of fuzzy L-TΓ-ideals of TΓ-semi ring T. Let $\pi = \bigcap_{i\in I} \pi_i$. Let $u, v, w \in Q$, $\gamma, \delta \in \Gamma$. Since $(\bigcap_{i\in I} \pi_i)(u) = \min\{\pi_i(u) : i \in I\}$ and each $\pi_i(u)$ is a fuzzy L-TΓ-ideal of Q. Therefore,

$$(\bigcap_{i\in I} \pi_i)(k + v) \geq \min\{\pi_i(k + v)\}$$
$$= \min\{\min\{\pi_i(k), \pi_i(v)\}$$
$$= \min\{\min\{\pi_i(k)\}, \min\{\{\pi_i(v)\}\}. \text{ and}$$

$\pi_i(k\gamma v\delta o) \geq \pi_i(o)$, $i \in I$ and $\forall k, v, o \in Q$, $\forall \gamma, \delta \in \Gamma$.

Now $\pi(k\gamma v\delta o) = (\bigcap_{i\in I} \pi_i)(k\gamma v\delta o)$

$$= \min\{(\pi_i)(k\gamma v\delta o) : i \in I\}$$
$$\geq \min\{\pi_i(o) : i \in I\}$$
$$= (\bigcap_{i\in I} \pi_i)(o) = \pi(o).$$

Therefore $\pi(k\gamma v\delta o) \geq \pi(o)$ and hence $\pi = \bigcap_{i\in I} \pi_i$ is a L- fuzzy TΓ-ideal of Q. Similarly, we can prove the remaining two parts.

Theorem 3.15: If $\{\pi_i : i \in I\}$ is a family of fuzzy left (lateral, right) TΓ-ideals of Γ-semi ring T, then $(\bigcup_{i\in I} \pi_i)(g)$ is a fuzzy left (lateral, right, TΓ-ideal) TΓ-ideal of H.

***Proof* :** Let $\{\pi_i : i \in I\}$ be a family of fuzzy left TΓ-ideals of TΓ-semi ring H. Let $\pi = \bigcup_{i\in I} \pi_i$. Let $k, v, o \in H$, $\gamma, \delta \in \Gamma$. Since $(\bigcup_{i\in I} \pi_i)(k) = \max\{\pi_i(k) : i \in I\}$ and $\pi_i(k)$ is a fuzzy left TΓ-ideal of S for some $i \in I$. Therefore,

$$\bigcup_{i\in I}(\pi_i)(k + v) \geq \max\{\pi_i(k + v)\}$$
$$= \max\{\min\{\pi_i(k), \pi_i(v)\}$$
$$= \max\{\min\{\pi_i(k)\}, \min\{\{\pi_i(v)\}\} \text{and}$$

$\pi_i(k\gamma v\delta o) \geq \pi_i(o)$, $i \in I$ and $\forall k, v, o \in H$, $\forall \gamma, \delta \in \Gamma$.

Now $\pi(k\gamma v\delta o) = (\bigcup_{i\in I} \pi_i)(k\gamma v\delta o)$

$$= \max\{(\pi_i)(k\gamma v\delta o) : i \in I\} \geq \max\{\pi_i(o) : i \in I\}$$
$$= (\bigcup \pi_i)(o) \text{ for some } i \in I = \pi(w).$$

Therefore $\pi(k\gamma v\delta o) \geq \pi(o)$ and hence $\pi = \bigcup_{i\in I} \pi_i$ is a L- fuzzy TΓ-ideal of Q. Similarly, prove the other parts.

Th 3.16: Let P(≠∅) be a subset of a TΓ-semiring Q and π_I be the characteristic function of I, then I is a L(La, R) TΓ-ideal of Q iff π_I is a fuzzy L(La, R) TΓ-ideal of Q.

***Proof*:** Let P be a L-TΓ-ideal of a TΓ-semi ring Q. Let $x, y, z \in Q$ and $\gamma, \delta \in \Gamma$, then $x + y \in I$ and $x\gamma y\delta z \in I$ if $z \in I$. It follows that $\chi_I(x + y) = 1$ and $\chi_I(x\gamma y\delta z) = 1 = \chi_I(z)$. If $y \notin I$, then $\chi_I(z) = 0$. In this case $\chi_I(x\gamma y\delta) \geq 0 = \chi_I(z)$. Therefore χ_I is a fuzzy L-TΓ-ideal of Q.

Conversely, suppose that χ_I be a fuzzy left Γ-ideal of S. Let $x, y \in I$, then if $x, y \in I$, then $\chi_I(x) = \chi_I(y) = 1$ and $\chi_I(x + y) \geq \min\{\chi_I(x), \chi_I(y)\} = \min\{1, 1\} = 1$. Thus $x + y \in I$. $\chi_I(x) = \chi_I(y) = 1$. Now let $x \in I$ and $s, t \in Q$, , $\delta \in \Gamma$. Then $\chi_I(x) = 1$. Also $\chi_I(s\gamma t\delta x) \geq \chi_I(x) = 1$. Thus $s\gamma t\delta x \in I$. So I is a L-TΓ-ideal of Q.

In the similar manner one can prove remaining two parts.

Th 3.17: Suppose X be a L(La, R) TΓ-ideal of a TΓ-semi ring V and $\gamma \leq \theta \neq 0$ be any 2 elements in [0, 1], then the fuzzy subset π of X, defined by $\pi(x) = \begin{cases} \theta \text{ if } x \in I \\ \gamma \text{ otherwise} \end{cases}$ is a fuzzy L(La, R) TΓ-ideal of X.

Proof: Let X be a L-TΓ-ideal of a TΓ-semi ring V and $\alpha, \beta \in [0,1]$. Let $s, t, x, y \in T$ and $\gamma, \delta \in \Gamma$. If $x, x + y \in I$, $\pi(x + y) \geq \min\{\pi(x), \mu(x)\} = \alpha \Rightarrow \alpha \leq \mu(x + y)$ and if $x \notin I$, $x + y \notin I$, then $\alpha = \mu(x + y) \geq \min\{\pi(x), \pi(x)\} = \alpha$. Therefore, $\pi(x + y) \geq \min\{\pi(x), \pi(x)\}$. Now $x \in X$, then $s\gamma t\delta x \in X$ and $\pi(x) = \theta = \pi(s\gamma t\delta x)$. Therefore $\pi(x) = \pi(s\gamma t\delta x)$. If $x \notin X$ then $\mu(x) = \gamma \leq \theta = \pi(s\gamma t\delta x)$ and then $\pi(s\gamma t\delta x) \geq \pi(x)$. Thus $\pi(s\gamma t\delta x) \geq \pi(x)$ $\forall$ $s, t, x \in V$ and $\gamma, \delta \in \Gamma$. Hence π is a fuzzy L-TΓ-ideal of V. Similarly, prove the other parts.

Th 3.18: Let V be a TΓ-semi ring and π be a non-empty fuzzy subset of V, then π is a fuzzy L(La. R) TΓ-ideal of V if and only if π_t's are L(La. R) TΓ-ideal of V for all $t \in \text{Im}(\pi)$ where $\pi_t = \{x \in V : \mu(x) \geq t\}$.

***Proof*:** Let π be a fuzzy L-TΓ-ideal of V. Let $t \in \text{Im}(\pi)$, then $\exists\ \alpha \in V \ni \pi(\alpha) = t$ and so $\alpha \in \pi_t$. Thus $\pi_t \neq \emptyset$. Let $a, q \in \pi_t$ then $a + q \geq t$ implies that $\pi(a + q) \geq \min\{\pi(a), \pi(q)\} \geq t$ and hence $\pi(a + p) \geq t$. Therefore $a + q \in \pi_t$. Now let $a \in \pi_t$ then $\pi(a) \geq t$. Again let $s, t \in V$, $e \in \pi_t$ and $\gamma, \delta \in \Gamma$. Now $\pi(s\gamma t\delta e) \geq \pi(e) \geq t$. Therefore, $s\gamma t\delta e \in \pi_t$. Thus π_t is a L-TΓ-ideal of V.

Conversely, suppose that π_t's are L-TΓ-ideals of V for all $t \in \text{Im}(\pi)$. Again let $a, q \in V$, $s, t \in V$ and $\gamma, \delta \in \Gamma$, then $\pi(a) = \pi(q) = t$. Since π_t is a L-TΓ-ideal of V and hence $a + q, s\gamma t\delta a \in \pi_t$. Therefore π $\pi(a + q) \geq \min\{\pi(a), \pi(q)\} = t$ and $\pi(s\gamma t\delta a) \geq t = \pi(a)$. Hence π is a fuzzy L-TΓ-ideal of V. Similarly, prove the other parts.

Def 3.19: Let V be a TΓ-semi ring and π be a fuzzy L(La, R) TΓ-ideal of a TΓ-semi ring V. Then the TTΓ-ideals π_t's are know as ***level L*(La, R) *TΓ-ideal*** of π where $t \in \text{Im}(\pi)$.

Th 3.20: Let π be a fuzzy L(La, R) TΓ-ideal of a TΓ-semiring V and $t_1 > t_2$. Then $\eta_{t_1} \subseteq \eta_{t_2}$ where $t_1, t_2 \in \text{Im}(\eta)$, equality occurs iff there is no $x \in V \ni t_1 \leq \eta(x) < t_2$.

***Proof*.** The 1st section of the theorem follows easily. Now let η be a fuzzy L-TΓ-ideal of $V \ni \eta_{t_1} = \eta_{t_2}$. If possible $\exists\ u \in V \ni t_1 \leq \eta(x) < t_2$. Then $x \in \eta_{t_1}$ but $x \notin \eta_{t_1}$, it is contradiction. So $\exists$ no $x \in V \ni t_1 \leq \eta(x) \leq t_2$.

Conversely, Let η is a fuzzy L-TΓ-ideal of $V \ni$ there does not exist $x \in V$ with $t_1 \leq \eta(x) < t_2$. $\because t_1 < t_2$, then $\eta_{t_1} \subseteq \eta_{t_2}$ (by Definition 3.15). If possible $\eta_{t_1} \neq \eta_{t_2}$. Then there is some $y \in V \ni y \in \eta_{t_1}$ but $y \notin \eta_{t_2}$, i.e, $\eta(y) \geq t_1$ but $\eta(y) < t_2$, i.e, $t_1 \leq \eta(y) < t_2$, it is a contradiction to our assumption. Hence, $\eta_{t_1} = \eta_{t_2}$. Similarly, prove the other sections of the statement.

Th 3.21 : Suppose V be a TΓ-semi ring. Every L(R)-Γ-ideal of V is two-sided iff every fuzzy L(R)-Γ-ideal of V is fuzzy two-sided.

Th 3.22: Let ς be a fuzzy subset of a TΓ-semi ring M. Then ς is a fuzzy TΓ-ideal of M iff $\forall\ t \in [0, 1]$, if $\varsigma_t^S \neq \emptyset$, then ς_t^S is a TΓ-ideal of M.

***Proof* :** Assume ς is a fuzzy TΓ-ideal of M. Then $\varsigma(a+p) \geq \wedge\{\varsigma(a), \varsigma(p)\}$ and $\varsigma(a\gamma p\delta w) \geq \varsigma(a) \vee \varsigma(p) \vee \varsigma(w)$ for any $a, p, w \in M$ and $\gamma, \delta \in \Gamma$. Thus $\varsigma(a) > t$. Since ς is a fuzzy TΓ-ideal of M, $\varsigma(a + p) \geq \min\{\varsigma(a), \varsigma(p)\} > t$ and $\varsigma(a\gamma p\delta w) \geq \varsigma(a) \vee \varsigma(p) \vee \varsigma(w) \geq \varsigma(a) > t$.

Therefore, $a + p,\ a\gamma p\delta w \in \varsigma_t^S$.

Similarly, $a + p,\ a\gamma p\delta w \in \varsigma_t^S$ and $a + p,\ a\gamma p\delta w \in \varsigma_t^S \Rightarrow \varsigma_t^S$ is a TTΓ-ideal of M.

Conversely, suppose that for all $t \in [0, 1]$, if $\varsigma_t^S \neq \emptyset$, then ς_t^S is a TΓ-ideal of M. Let $a, p, w \in M$ and $\gamma, \delta \in \Gamma$.

Case 1: $\varsigma(a) \geq \varsigma(p) \geq \xi(w)$. Thus $a \in \varsigma_t^S\ \forall\ t < \varsigma(a)$.

By our supposition, we have ς_t^S is a TΓ-ideal of M $\forall\ t < \varsigma(a)$.

So $a\gamma p\delta w \in \varsigma_t^S\ \forall\ t < \varsigma(a)$.

Then $\varsigma(a\gamma p\delta w) > t\ \forall\ t < \varsigma(a)$.

Then $\varsigma(a\gamma p\delta w) \geq \varsigma(a) = \varsigma(a) \vee \varsigma(p) \vee \varsigma(w)$.

Case 2: $\varsigma(a) < \varsigma(p) < \varsigma(w)$. Then, $p \in \varsigma_t^S\ \forall\ t < \varsigma(w)$.

By our supposition, we have ς_t^S is a TΓ-ideal $\forall\ t < \varsigma(w)$.

So $a\gamma p\delta w \in \varsigma_t^S\ \forall\ t < \varsigma(w)$.

Then, $\varsigma(a\gamma p\delta w) > t\ \forall\ t < \varsigma(w)$.

Then $\varsigma(a\gamma p\delta w) \geq \varsigma(w) = \varsigma(a) \vee \varsigma(p) \vee \varsigma(w)$.

Therefore ς is a fuzzy TΓ-ideal of M.

4. Conclusion

many of the researchers were studied about fuzzy ideals in semigroups, γ-semigroups, ternary semigroups, semirings and near rings. here, we extended those concepts to fuzzy tγ-ideals in ternary γ-semirings.

References

[1] T. K. Dutta and Chanda. T., *Structres of Fuzzy Ideals of Γ-Ring*; Bull. Malays. Math. Sci. Soc. (2) 28(1) (2005), 9-18.

[2] M. Sajani Lavanya, Dr. D. Madhusudhana Rao, and V. Syam Julius Rajendra; *On Lateral Ternary Γ-Ideals of Ternary Γ-Semirings*-American International Journal of Research in Science, Technology, Engineering &Mathematics (AIJRSTEM), 12(1), September-November, 2015, pp: 11-14.

[3] M. Sajani Lavanya, Dr. D. Madhusudhana Rao, and Prof. K. Panduranga Rao; *On simple Ternary* Γ-*Smiring*, International Journal of Engineering Research and Application, Vol.7, Issue 2, (Part-5) February 2017, pp: 01-06.

[4] M. Sajani Lavanya, Dr. D. Madhusudhana Rao, and VB Subrahmanyeswara rao Seetam Raju; *On Right Ternary* Γ-*Ideals of Ternary* Γ-*Semiring*, International Journal of Research in Applied Natural and Social Sciences, Vol. 4, Issue 5, May 2016, 107-114.

[5] K. Syam Prasad and Bh. Satyanarayana., *Fuzzy Prime Ideal of a Gamma Near-Ring*; Soochow Journal of Mathematics, Volume 31, No. 1, pp: 121-129.