



# Oscillation Criteria for Generalized Second Kind Sublinear Delay Difference Equations

A.Benevatho Jaison<sup>1\*</sup>, Sk.Khadar Babu<sup>2</sup>, V.Chandrasekar<sup>3</sup>

<sup>1,2</sup>Vellore Institute of Technology, Vellore – 632 014, Tamil Nadu, India

<sup>3</sup>Thiruvalluvar University College of Arts and Science, Thennangur - 604 408, Tamil Nadu, India

\*Corresponding author E-mail: benejaison@gmail.com

## Abstract

The Using Riccati transformation techniques, we present some new oscillation criteria for generalized second kind nonlinear difference equation

$$\Delta_{\ell_2} (p(k\ell_2 + j_2)(\Delta_{\ell_1} (u(k\ell_1 + j_1)))^\gamma) + q(k\ell_2 + j_2)u^\beta((k - \sigma)\ell_2 + j_2) = 0,$$

when  $0 < \beta < 1$  is a quotient of odd positive integers,  $k \in [0, \infty)$ ,  $\ell_1, \ell_2 \in (0, \infty)$ ,  $j = k - \left\lfloor \frac{k}{\ell} \right\rfloor \ell$ .

**Keywords:** Generalized Difference Equation; Second Kind; Oscillation.

## 1. Introduction

Difference equations is a captivating mathematical area on its own as well as a rich field of the applications in such various disciplines such as population dynamics, operations research, ecology, economics, biology etc. For general background as difference equations with numerous examples from various fields, one can refer to [1]. The study of difference equations is based on the operator  $\Delta$  defined as  $\Delta u(k) = u(k+1) - u(k), k \in \mathbb{N} = \{0, 1, 2, \dots\}$ . Eventhough many authors [1],[16] have suggested the definition of  $\Delta$  as

$$\Delta u(k) = u(k + \ell) - u(k), \quad \ell \in (0, \infty), \quad (1)$$

no major progress have taken place on this line.

In [14], authors took up the definition of  $\Delta$  as given in (1), they produced some important results. The results obtained can be found in [14]. They labeled the operator  $\Delta$  defined by (1) as  $\Delta_\ell$  and by defining its inverse  $\Delta_\ell^{-1}$ . Recently, there has been increasing study in the asymptotic behavior of second order difference equations by many authors [2-12,17-24].

In the sequel we will be concerned the generalized second order sublinear delay difference equations of the form

$$\Delta_{\ell_2} (p(k\ell_2 + j_2)(\Delta_{\ell_1} (u(k\ell_1 + j_1)))^\gamma) + q(k\ell_2 + j_2)u^\beta((k - \sigma)\ell_2 + j_2) = 0, \quad (2)$$

where  $\Delta_\ell$  denotes the generalized difference operator for any real valued function  $u(k\ell + j)$ ,  $k, \ell_1, \ell_2 \in (0, \infty)$ ,  $j = k - \left\lfloor \frac{k}{\ell} \right\rfloor \ell$ ,  $\gamma$

and  $0 < \beta < 1$  are quotient of odd positive integers,  $\sigma$  is a fixed

nonnegative integer,  $p(k\ell_2 + j_2) > 0$ , and  $q(k\ell_2 + j_2) \geq 0$  are real valued functions, and

$$\sum_{k=k_0}^{\infty} \left( \frac{1}{p(k\ell_2 + j_2)} \right)^{1/\gamma} = \infty, \quad \text{for some } k_0 > 0. \quad (3)$$

By a solution of (2) we mean a nontrivial real valued function  $u(k\ell_1 + j_1)$  defined for  $k \geq -\sigma$ , and satisfies equation (2) for  $k \in [0, \infty)$ . Suppose

$$u(k\ell_1 + j_1) = A(k\ell_1 + j_1) \text{ for } k \in [-\sigma, 0] \quad (4)$$

then equation (2) has a unique solution satisfying the initial condition (4).

A solution  $u(k\ell + j)$  of equation (2) is said to be oscillatory if for every  $k_1 > 0$  there exists an  $k \geq k_1$  such that  $u(k\ell + j)u((k+1)\ell + j) \leq 0$ , otherwise it is nonoscillatory. Equation (2) is said to be oscillatory if all its solutions are oscillatory.

## 2. Main Results

### Theorem 2.1

Assume that (3) holds. Also, assume that there exists a real valued functions  $\rho(k\ell_2 + j_2)$  such that for every  $\eta \geq 1$  and positive number M

$$\limsup_{k \rightarrow \infty} \sum_{r=k_0}^k [\rho(r\ell_2 + j_2)q(r\ell_2 + j_2) \frac{(p((r - \sigma)\ell_2 + j_2))^{1/\gamma} \eta^{1-\beta} ((r - \sigma + 1)\ell_2 + j_2)^{1-\beta} (\Delta_{\ell_2} \rho(r\ell_2 + j_2))^2}{4\beta(M)^{(\gamma-1)/\gamma} \rho(r\ell_2 + j_2)}] \quad (5)$$

Then every solution of equation (2) oscillates.

**Proof.**

Suppose to the contrary that  $u(k\ell_1 + j_1)$  is an eventually nonoscillatory solution of (2) such that  $u((k - \sigma)\ell_2 + j_2) > 0$  for all  $k \geq k_0 > 0$ . Since the substitution  $v(k\ell_1 + j_1) = -u(k\ell_1 + j_1)$  transforms equation (2) into an equation of the same form, we shall consider only this case. From equation (2), we have

$$\Delta_{\ell_2} (p(k\ell_2 + j_2)(\Delta_{\ell_1} u(k\ell_1 + j_1))^\gamma) = -q(k\ell_2 + j_2)u^\beta((k - \sigma)\ell_2 + j_2) \leq 0, \tag{6}$$

for  $k \geq k_0$  and so  $p(k\ell_2 + j_2)(\Delta_{\ell_1} u(k\ell_1 + j_1))^\gamma$  is an eventually nonincreasing sequence. We first show that  $p(k\ell_2 + j_2)(\Delta_{\ell_1} u(k\ell_1 + j_1))^\gamma \geq 0$  for  $k \geq k_0$ . In fact, if there exists an real  $k_1 \geq k_0$  such that  $p(k_1\ell_2 + j_2)(\Delta_{\ell_1} u(k_1\ell_1 + j_1))^\gamma = c < 0$ , then (6) implies that  $p(k\ell_2 + j_2)(\Delta_{\ell_1} u(k\ell_1 + j_1))^\gamma \leq c$  for  $k \geq k_1$  that is

$$\Delta_{\ell_1} u(k\ell_1 + j_1) \leq \left( \frac{c}{p(k\ell_2 + j_2)} \right)^{1/\gamma}$$

and, hence as  $k \rightarrow \infty$

$$u(k\ell_1 + j_1) \leq u(k_1\ell_1 + j_1) + c^{1/\gamma} \sum_{i=0}^{k-1} \left( \frac{1}{p(k_1\ell_2 - i\ell_1 + j_2)} \right)^{1/\gamma} \rightarrow -\infty \tag{7}$$

which contradicts the fact that  $u(k\ell_1 + j_1) > 0$  for  $k \geq k_0$ , then  $p(k\ell_2 + j_2)(\Delta_{\ell_1} u(k\ell_1 + j_1))^\gamma \geq 0$ . Also we claim that  $\Delta_{\ell_1}^2 u(k\ell_1 + j_1) \leq 0$ . If not there exists  $k_1 \geq k_0$  such that  $\Delta_{\ell_1}^2 u(k\ell_1 + j_1) > 0$  for  $k \geq k_1$  and this implies that  $\Delta_{\ell_1} u((k+1)\ell_1 + j_1) > \Delta_{\ell_1} u(k\ell_1 + j_1)$ , so that since  $\Delta_{\ell_2} p(k\ell_2 + j_2) \geq 0$ ,

$$p((k+1)\ell_2 + j_2)(\Delta_{\ell_1} u((k+1)\ell_1 + j_1))^\gamma > p((k+1)\ell_2 + j_2)(\Delta_{\ell_1} u(k\ell_1 + j_1))^\gamma \geq p(k\ell_2 + j_2)(\Delta_{\ell_1} u(k\ell_1 + j_1))^\gamma$$

and this contradicts the fact that  $p(k\ell_2 + j_2)(\Delta_{\ell_1} u(k\ell_1 + j_1))^\gamma$  is nonincreasing sequence, then  $\Delta_{\ell_1}^2 u(k\ell_1 + j_1) \leq 0$ , and therefore we have

$$u(k\ell_1 + j_1) > 0, \Delta_{\ell_1} u(k\ell_1 + j_1) \text{ and } \Delta_{\ell_1}^2 u(k\ell_1 + j_1) \leq 0 \text{ for } k \geq k_0. \tag{8}$$

Define the sequence  $z(k\ell_2 + j_2)$  by

$$z(k\ell_2 + j_2) = \rho(k\ell_2 + j_2) \frac{p(k\ell_2 + j_2)(\Delta_{\ell_1} u(k\ell_1 + j_1))^\gamma}{u^\beta((k - \sigma)\ell_2 + j_2)} \tag{9}$$

then  $z(k\ell_2 + j_2) > 0$ , and

$$\Delta_{\ell_2} z(k\ell_2 + j_2) = p((k+1)\ell_2 + j_2)(\Delta_{\ell_1} u(k\ell_1 + \ell_2 + j_1)) \Delta_{\ell_2} \left( \frac{\rho(k\ell_2 + j_2)}{u^\beta((k - \sigma)\ell_2 + j_2)} \right)$$

$$+ \frac{\rho(k\ell_2 + j_2)\Delta_{\ell_2} (p(k\ell_2 + j_2)(\Delta_{\ell_1} u(k\ell_1 + j_1))^\gamma)}{u^\beta((k - \sigma)\ell_2 + j_2)} \tag{10}$$

From (2) and (10), we have

$$\Delta_{\ell_2} z(k\ell_2 + j_2) = -\rho(k\ell_2 + j_2)q(k\ell_2 + j_2) + \frac{\Delta_{\ell_2} \rho(k\ell_2 + j_2)}{\rho((k+1)\ell_2 + j_2)} z((k+1)\ell_2 + j_2) - \left( \frac{\rho(k\ell_2 + j_2)p((k+1)\ell_2 + j_2)}{u^\beta((k - \sigma + 1)\ell_2 + j_2)} \times \frac{(\Delta_{\ell_1} u(k\ell_2 + \ell_2 + j_1))^\gamma \Delta_{\ell_2} u^\beta((k - \sigma)\ell_2 + j_2)}{u^\beta((k - \sigma)\ell_2 + j_2)} \right) \tag{11}$$

From (6) and (8), we get

$$p((k - \sigma)\ell_2 + j_2)(\Delta_{\ell_1} u(k\ell_1 - \sigma\ell_2 + j_1))^\gamma \geq p((k+1)\ell_2 + j_2)(\Delta_{\ell_1} u(k\ell_1 + \ell_2 + j_1))^\gamma \text{ and } u((k+1 - \sigma)\ell_2 + j_2) \geq u((k - \sigma)\ell_2 + j_2) \tag{12}$$

and then from (11) and (12), we have

$$\Delta_{\ell_2} z(k\ell_2 + j_2) = -\rho(k\ell_2 + j_2)q(k\ell_2 + j_2) + \frac{\Delta_{\ell_2} \rho(k\ell_2 + j_2)}{\rho((k+1)\ell_2 + j_2)} z((k+1)\ell_2 + j_2) - \frac{\rho(k\ell_2 + j_2)p((k+1)\ell_2 + j_2)}{(u^\beta((k - \sigma + 1)\ell_2 + j_2))^2} \times (\Delta_{\ell_1} u(k\ell_2 + \ell_2 + j_1))^\gamma \Delta_{\ell_2} u^\beta((k - \sigma)\ell_2 + j_2) \tag{13}$$

Now, by using the inequality in [13]

$$u^\beta - v^\beta \geq \beta u^{\beta-1}(u - v) \text{ for all } u \neq v > 0 \text{ and } 0 < \beta \leq 1.$$

Then, we have

$$\Delta_{\ell_2} u^\beta((k - \sigma)\ell_2 + j_2) = \beta(u((k - \sigma + 1)\ell_2 + j_2))^{\beta-1} \Delta_{\ell_2} u((k - \sigma)\ell_2 + j_2). \tag{14}$$

Substitute from (14) in (13), we have

$$\Delta_{\ell_2} z(k\ell_2 + j_2) = -\rho(k\ell_2 + j_2)q(k\ell_2 + j_2) + \frac{\Delta_{\ell_2} \rho(k\ell_2 + j_2)}{\rho((k+1)\ell_2 + j_2)} z((k+1)\ell_2 + j_2) - \left( \frac{\rho(k\ell_2 + j_2)p((k+1)\ell_2 + j_2)\beta(u((k+1 - \sigma)\ell_2 + j_2))^{\beta-1}}{(u^\beta((k - \sigma + 1)\ell_2 + j_2))^2} \right) \times \Delta_{\ell_2} u((k - \sigma)\ell_2 + j_2)(\Delta_{\ell_1} u(k\ell_1 + \ell_2 + j_1))^\gamma \tag{15}$$

From (12) and (15), we have

$$\Delta_{\ell_2} z(k\ell_2 + j_2) = -\rho(k\ell_2 + j_2)q(k\ell_2 + j_2) + \frac{\Delta_{\ell_2} \rho(k\ell_2 + j_2)}{\rho((k+1)\ell_2 + j_2)} z((k+1)\ell_2 + j_2) - \frac{\beta \rho(k\ell_2 + j_2)p((k+1)\ell_2 + j_2)^{1/\gamma}}{(p((k - \sigma)\ell_2 + j_2))^{1/\gamma}} \times \frac{p((k+1)\ell_2 + j_2)(\Delta_{\ell_1} u(k\ell_1 + \ell_2 + j_1))^{\gamma+1}}{(u((k - \sigma + 1)\ell_2 + j_2))^{1-\beta} (u^\beta((k - \sigma + 1)\ell_2 + j_2))}$$

$$\begin{aligned}
 &= -\rho(k\ell_2 + j_2)q(k\ell_2 + j_2) + \frac{\Delta_{\ell_2}\rho(k\ell_2 + j_2)}{\rho((k+1)\ell_2 + j_2)}z((k+1)\ell_2 + j_2) \\
 &\left[ \frac{\beta\rho(k\ell_2 + j_2)(p((k+1)\ell_2 + j_2))^{(1/\gamma)-1}(p((k+1)\ell_2 + j_2))^2}{(p((k+1)\ell_2 + j_2))^2(p((k-\sigma)\ell_2 + j_2))^{1/\gamma}(u((k-\sigma+1)\ell_2 + j_2))^{1-\beta}} \right. \\
 &\left. \times \frac{(\rho((k+1)\ell_2 + j_2))^2(\Delta_{\ell_1}u(k\ell_1 + \ell_2 + j_1))^{2\gamma}}{(u^\beta((k-\sigma+1)\ell_2 + j_2))^2(\Delta_{\ell_1}u(k\ell_1 + \ell_2 + j_1))^{\gamma-1}} \right] \quad (16)
 \end{aligned}$$

From (8), we conclude that

$$u(k\ell_2 + j_2) \leq u(k_0\ell_2 + j_2) + \Delta_{\ell_2}u(k_0\ell_2 + j_2)((k - k_0)\ell_2 + j_2), k \geq k_0$$

consequently there exists a  $k_1 \geq k_0$  and appropriate constant  $\eta \geq 1$  such that

$$u(k\ell_2 + j_2) \leq \eta(k\ell_2 + j_2) \text{ for } k \geq k_1$$

and this implies that

$$u((k - \sigma + 1)\ell_2 + j_2) \leq \eta((k - \sigma + 1)\ell_2 + j_2) \text{ for } k \geq k_2 = k_1 + \sigma - 1$$

and, hence

$$\frac{1}{(u((k - \sigma + 1)\ell_2 + j_2))^{1-\beta}} \geq \frac{1}{(\eta((k - \sigma + 1)\ell_2 + j_2))^{1-\beta}} \quad (17)$$

Since  $p(k\ell_2 + j_2)(\Delta_{\ell_1}u(k\ell_1 + j_1))^\gamma$  is a positive and increasing function, there exists a  $k_2 \geq k_1$  sufficiently large such that  $p(k\ell_2 + j_2)(\Delta_{\ell_1}u(k\ell_1 + j_1))^\gamma \leq 1/M$  for some positive constant  $M$  and  $k \geq k_2$ , and hence by (6) we have  $p((k+1)\ell_2 + j_2)(\Delta_{\ell_1}u(k\ell_1 + \ell_2 + j_1))^\gamma \leq 1/M$ , so that

$$\frac{1}{(\Delta_{\ell_1}u(k\ell_1 + \ell_2 + j_1))^{\gamma-1}} \geq (Mp((k+1)\ell_2 + j_2))^{(\gamma-1)/\gamma} \quad (18)$$

then from (9), (16), (17) and (18) we have

$$\Delta_{\ell_2}z(k\ell_2 + j_2) =$$

$$\begin{aligned}
 &-\rho(k\ell_2 + j_2)q(k\ell_2 + j_2) + \frac{\Delta_{\ell_2}\rho(k\ell_2 + j_2)}{\rho((k+1)\ell_2 + j_2)}z((k+1)\ell_2 + j_2) \\
 &\frac{\beta\rho(k\ell_2 + j_2)M^{(\gamma-1)/\gamma}(z((k+1)\ell_2 + j_2))^2}{(p((k+1)\ell_2 + j_2))^2(p((k-\sigma)\ell_2 + j_2))^{1/\gamma}(\eta((k-\sigma+1)\ell_2 + j_2))^{1-\beta}} \quad (19)
 \end{aligned}$$

$$\begin{aligned}
 &\Delta_{\ell_2}z(k\ell_2 + j_2) \leq -\rho(k\ell_2 + j_2)q(k\ell_2 + j_2) + \\
 &\frac{(p((k-\sigma)\ell_2 + j_2))^{1/\gamma}\eta^{1-\beta}((k-\sigma+1)\ell_2 + j_2)^{1-\beta}(\Delta_{\ell_2}\rho(k\ell_2 + j_2))^2}{4\beta(M)^{(\gamma-1)/\gamma}\rho(k\ell_2 + j_2)} \\
 &\left[ \frac{\sqrt{\beta(M)^{(\gamma-1)/\gamma}\rho(k\ell_2 + j_2)z((k+1)\ell_2 + j_2)}}{\rho((k+1)\ell_2 + j_2)\sqrt{\eta((k-\sigma+1)\ell_2 + j_2)^{1-\beta}p((k-\sigma)\ell_2 + j_2)}} \right]
 \end{aligned}$$

$$\begin{aligned}
 &\left[ \frac{\sqrt{\eta^{1-\beta}((k-\sigma+1)\ell_2 + j_2)^{1-\beta}(p((k-\sigma)\ell_2 + j_2))^{1/\gamma}\Delta_{\ell_2}\rho(k\ell_2 + j_2)}}{2\sqrt{\beta(M)^{(\gamma-1)/\gamma}\rho(k\ell_2 + j_2)}} \right]^2 \\
 &< -[\rho(k\ell_2 + j_2)q(k\ell_2 + j_2) \\
 &\frac{(p((k-\sigma)\ell_2 + j_2))^{1/\gamma}\eta^{1-\beta}((k-\sigma+1)\ell_2 + j_2)^{1-\beta}(\Delta_{\ell_2}\rho(k\ell_2 + j_2))^2}{4\beta(M)^{(\gamma-1)/\gamma}\rho(k\ell_2 + j_2)}] \quad (20)
 \end{aligned}$$

Summing (20) from  $k_2$  to  $k$ , we obtain

$$\begin{aligned}
 & -z(k_2\ell_2 + j_2) < z((k+1)\ell_2 + j_2) - z(k_2\ell_2 + j_2) \\
 & < -\sum_{r=k_2}^k [\rho(r\ell_2 + j_2)q(r\ell_2 + j_2) \\
 & \frac{(p((r-\sigma)\ell_2 + j_2))^{1/\gamma}\eta^{1-\beta}((r-\sigma+1)\ell_2 + j_2)^{1-\beta}(\Delta_{\ell_2}\rho(r\ell_2 + j_2))^2}{4\beta(M)^{(\gamma-1)/\gamma}\rho(r\ell_2 + j_2)}]
 \end{aligned}$$

which yields

$$\begin{aligned}
 & \sum_{r=k_2}^k [\rho(r\ell_2 + j_2)q(r\ell_2 + j_2) \\
 & \frac{(p((r-\sigma)\ell_2 + j_2))^{1/\gamma}\eta^{1-\beta}((r-\sigma+1)\ell_2 + j_2)^{1-\beta}(\Delta_{\ell_2}\rho(r\ell_2 + j_2))^2}{4\beta(M)^{(\gamma-1)/\gamma}\rho(r\ell_2 + j_2)}] < c_1
 \end{aligned}$$

for all large  $k$ , and this is contrary to (5). The proof is complete.

**Theorem 2.2**

Assume that (3) holds. Let  $\rho(k\ell_2 + j_2)$  be a real valued function.

Also, we assume that there exists a double fuction

$$H(m, k\ell_2 + j_2); m \geq k \geq 0 \text{ such that}$$

$$(i) H(m, m) = 0 \text{ for } m \geq 0, (ii) H(m, k\ell_2 + j_2) > 0 \text{ for}$$

$$m > k\ell_2 + j_2 \geq 0,$$

$$(iii) \Delta_{2(\ell_2)}H(m, k\ell_2 + j_2) = H(m, (k+1)\ell_2 + j_2) - H(m, k\ell_2 + j_2) \leq 0$$

for  $m \geq k\ell_2 + j_2 \geq 0$ . If

$$\begin{aligned}
 & \limsup_{m \rightarrow \infty} \frac{1}{H(m, 0)} \sum_{k=k_0}^{m-1} [H(m, k\ell_2 + j_2)\rho(k\ell_2 + j_2)q(k\ell_2 + j_2) \\
 & \frac{(\rho((k+1)\ell_2 + j_2))^2}{\rho(k\ell_2 + j_2)} \\
 & \left( h(m, k\ell_2 + j_2) - \frac{\Delta_{\ell_2}\rho(k\ell_2 + j_2)}{\rho((k+1)\ell_2 + j_2)}\sqrt{H(m, k\ell_2 + j_2)} \right)^2] = \infty, \quad (21)
 \end{aligned}$$

$$\text{where } h(m, k\ell_2 + j_2) = -\frac{\Delta_{2(\ell_2)}H(m, k\ell_2 + j_2)}{\sqrt{H(m, k\ell_2 + j_2)}}, m \geq k \geq 0,$$

$$\rho(k\ell_2 + j_2) = \frac{\beta\rho(k\ell_2 + j_2)M^{(\gamma-1)/\gamma}}{(p((k-\sigma)\ell_2 + j_2))^{1/\gamma}\eta^{1-\beta}((k-\sigma+1)\ell_2 + j_2)^{1-\beta}}$$

Then every solution of equation (2) oscillates.

*Proof:* We proceed as in Theorem 2.1, we assume that equation (2)

has a nonoscillatory solution, say  $u((k-\sigma)\ell_2 + j_2) > 0$  for all  $k \geq k_0$ . From the proof of Theorem 2.1, we obtain (18) for all

$k \geq k_2$ . From (18) we have for  $k \geq k_2$ ,

$$\begin{aligned}
 & \rho(k\ell_2 + j_2)q(k\ell_2 + j_2) \leq -\Delta_{\ell_2}z(k\ell_2 + j_2) \\
 & + \frac{\Delta_{\ell_2}\rho(k\ell_2 + j_2)z((k+1)\ell_2 + j_2)}{\rho((k+1)\ell_2 + j_2)} - \frac{\rho(k\ell_2 + j_2)z^2((k+1)\ell_2 + j_2)}{(\rho((k+1)\ell_2 + j_2))^2}. \quad (22)
 \end{aligned}$$

Therefore, we have

$$\begin{aligned}
 & \sum_{k=n}^{m-1} H(m, k\ell_2 + j_2)\rho(k\ell_2 + j_2)q(k\ell_2 + j_2) \\
 & \leq -\sum_{k=n}^{m-1} H(m, k\ell_2 + j_2)\Delta_{\ell_2}z(k\ell_2 + j_2) \\
 & + \sum_{k=n}^{m-1} \frac{H(m, k\ell_2 + j_2)\Delta_{\ell_2}\rho(k\ell_2 + j_2)z((k+1)\ell_2 + j_2)}{\rho((k+1)\ell_2 + j_2)}
 \end{aligned}$$

$$-\sum_{k=n}^{m-1} \frac{H(m, k\ell_2 + j_2) \overline{\rho(k\ell_2 + j_2)} z^2((k+1)\ell_2 + j_2)}{(\rho((k+1)\ell_2 + j_2))^2}$$

which yields, after summing by parts the first term in the right hand side.

$$\begin{aligned} & \sum_{k=n}^{m-1} H(m, k\ell_2 + j_2) \rho(k\ell_2 + j_2) q(k\ell_2 + j_2) \leq H(m, k\ell_2 + j_2) z(k\ell_2 + j_2) \\ & + \sum_{k=n}^{m-1} z((k+1)\ell_2 + j_2) \Delta_{2(\ell_2)} H(m, k\ell_2 + j_2) \\ & + \sum_{k=n}^{m-1} H(m, k\ell_2 + j_2) \frac{\Delta_{\ell_2} \rho(k\ell_2 + j_2)}{\rho((k+1)\ell_2 + j_2)} z((k+1)\ell_2 + j_2) \\ & - \sum_{k=n}^{m-1} \frac{H(m, k\ell_2 + j_2) \overline{\rho(k\ell_2 + j_2)}}{(\rho((k+1)\ell_2 + j_2))^2} z^2((k+1)\ell_2 + j_2) \\ & = H(m, k\ell_2 + j_2) z(k\ell_2 + j_2) - \sum_{k=n}^{m-1} [z((k+1)\ell_2 + j_2) \\ & + \frac{\rho((k+1)\ell_2 + j_2)}{2\sqrt{H(m, k\ell_2 + j_2) \rho(k\ell_2 + j_2)}} \\ & \times \left( h(m, k\ell_2 + j_2) \sqrt{H(m, k\ell_2 + j_2)} - \frac{\Delta_{\ell_2} \rho(k\ell_2 + j_2) H(m, k\ell_2 + j_2)}{\rho((k+1)\ell_2 + j_2)} \right)^2] \\ & + \frac{1}{4} \sum_{k=n}^{m-1} \frac{(\rho((k+1)\ell_2 + j_2))^2}{\rho(k\ell_2 + j_2)} \\ & \times \left( h(m, k\ell_2 + j_2) - \frac{\Delta_{\ell_2} \rho(k\ell_2 + j_2)}{\rho((k+1)\ell_2 + j_2)} \sqrt{H(m, k\ell_2 + j_2)} \right)^2 \\ & \sum_{k=k_1}^{m-1} \frac{[H(m, k\ell_2 + j_2) \rho(k\ell_2 + j_2) q(k\ell_2 + j_2)]}{(\rho((k+1)\ell_2 + j_2))^2} \\ & \frac{1}{4\rho(k\ell_2 + j_2)} \\ & \times \left( h(m, k\ell_2 + j_2) - \frac{\Delta_{\ell_2} \rho(k\ell_2 + j_2) \sqrt{H(m, k\ell_2 + j_2)}}{\rho((k+1)\ell_2 + j_2)} \right)^2 \\ & < H(m, k_1\ell_2 + j_2) z(k_1\ell_2 + j_2) \leq H(m, 0) z(k_1\ell_2 + j_2) \\ & < H(m, 0) \left( z(k_1\ell_2 + j_2) + \sum_{k=0}^{k_1-1} \rho(k\ell_2 + j_2) q(k\ell_2 + j_2) \right) < \infty. \end{aligned}$$

Contradicts the assumption. Hence the proof.

## References

- [1] Agarwal RP, *Difference Equations and Inequalities*, Marcel Dekker, New York, (2000).
- [2] Benevatho Jaison A, Khadar Babu Sk (2016), Oscillation for generalized first order nonlinear difference equations. *Global Journal of Pure and Applied Mathematics* 12(1), 51-54.
- [3] Benevatho Jaison A, Khadar Babu Sk (2016), Kamenev-type oscillation criteria for second order generalized delay difference equations. *International Journal of Control Theory and Applications* 9(28), 463-469.
- [4] Benevatho Jaison A, Khadar Babu Sk (2016), Oscillation for generalized first order nonlinear a-difference equations. *International Journal of Pure and Applied Mathematics* 109(7), 67-74.
- [5] Benevatho Jaison A, Khadar Babu Sk (2017), Oscillation theorems for generalized second-order nonlinear delay difference equations. *International Journal of Pure and Applied Mathematics* 113(9), 84-92.
- [6] Benevatho Jaison A, Khadar Babu Sk (2017), Oscillation theorems for generalized second kind nonlinear delay difference equations. *International Journal of Pure and Applied Mathematics* 115(9), 25-36.
- [7] Benevatho Jaison A, Khadar Babu Sk (2017), Oscillatory behavior of generalized nonlinear difference equations. *Global Journal of Pure and Applied Mathematics* 13(1), 205-209.

- [8] Benevatho Jaison A, Khadar Babu Sk (2017), Oscillatory behavior of generalized Nonlinear difference equations. *Global Journal of Pure and Applied Mathematics* 13(2), 415-423.
- [9] Benevatho Jaison A, Khadar Babu Sk (2018), Kamenev-type oscillation criteria for generalized sublinear delay difference equations. *International Journal of Pure and Applied Mathematics* 118(10), 135-145.
- [10] Benevatho Jaison A, Khadar Babu Sk (2018), Oscillation for generalized second kind nonlinear delay a-difference equations. *International Journal of Pure and Applied Mathematics* 118(23), 507-515.
- [11] Chandrasekar V, Srimanju V (2016), Oscillation for generalized second order sublinear neutral delay alpha difference equations. *Global Journal of Pure and Applied Mathematics* 12(1), 55-59.
- [12] Chandrasekar V, Srimanju V (2016), Qualitative properties of discrete version of generalized kneser's and arzela-ascoli's theorems. *International Journal of Control Theory and Applications* 9(28), 549-554.
- [13] Hardy GH, Littlewood JE, Polya G, 2nd edn., Cambridge University Press, Cambridge, (1952), 39-40.
- [14] Maria Susai Manuel M, Britto Antony Xavier G, Thandapani E (2006), Theory of generalized difference operator and its applications. *Far East Journal of Mathematical Sciences* 20(2), 163-171.
- [15] Maria Susai Manuel M, Britto Antony Xavier G, Thandapani E (2006), Qualitative properties of solutions of certain class of difference equations. *Far East Journal of Mathematical Sciences* 23(3), 295-304.
- [16] Ronald E. Mickens, *Difference Equations*, Van Nostrand Reinhold Company, New York (1990).
- [17] Srimanju V, Khadar Babu Sk (2017), Oscillatory criteria for generalized second-order quasilinear neutral delay difference equations. *International Journal of Pure and Applied Mathematics* 113(9), 75-83.
- [18] Srimanju V, Khadar Babu Sk (2017), Oscillation of generalized quasilinear difference equations. *International Journal of Pure and Applied Mathematics* 115(9), 37-45.
- [19] Srimanju V, Khadar Babu Sk (2017), Oscillation criteria for generalized quasi-linear difference equations. *Global Journal of Pure and Applied Mathematics* 13(1), 210-216.
- [20] Srimanju V, Khadar Babu Sk (2017), Oscillation criteria for generalized second kind quasi-linear neutral a-difference equations. *Global Journal of Pure and Applied Mathematics* 13(2), 544-551.
- [21] Srimanju V, Khadar Babu Sk (2018), Oscillatory properties of third-order quasilinear generalized difference equations. *International Journal of Pure and Applied Mathematics* 118(10), 155-165.
- [22] Srimanju V, Khadar Babu Sk (2018), Oscillation of generalized quasilinear a-difference equations. *International Journal of Pure and Applied Mathematics* 118(23), 497-505.
- [23] Szafrański Z, Szmada B (1997), Oscillation theorems for some nonlinear difference equations. *Appl. Math. Comput.* 83 43-52.
- [24] Szmada B (1983), Oscillation criteria for second order non-linear difference equations. *Anal. Pol. Math.* 153, 225-235.