International Journal of Engineering & Technology, 7 (4.10) (2018) 366-370

International Journal of Engineering & Technology

Website: www.sciencepubco.com/index.php/IJET

Research paper



Self Centeredness of Total Graphs W.R.T. D-Distance

M.V. Ramanjaneyulu[1], P. L. N.Varma[2], D. Reddy Babu[3]

[1]*Dept. of S and H, KKR and KSR Inst. of Tech. and Sci.,Vinjanampadu,Guntur 522017, India.*
[2]*Dept. of S. and H. ,V.F.S.T.R. University,Vadlamudi, Guntur 522 217, India.*
[3] *Dept. of Mathematics GST, GITAM University,Doddaballapura-561203,Bengaluru North,Karnataka, India.*
**Corresponding Author E-Mail: reddybabu17@gmail.com , mutyam.09@gmail.com , plnvarma@gmail.com*

Abstract
In this article using *D*-distance, we study the self-centeredness of total graphs. We end with some open questions.

Keywords: *Total graph, D-distance, self-centered graphs*

1. Introduction

Given a graph $G=(V,E)$, the *total graph* of *G* is a graph whose vertex set consists of all the vertices and edges of *G* i.e., $V \cup E$ and two vertices are adjacent in the new graph if and only if they are adjacent in the graph *G* (see [1]). The total graph of *G* is denoted by $T(G)$ or G^T.

The concept of *D*-distance in a graph *G* was introduced by Reddy Babu and Varma in [3]. The *D*-distance, denoted by $d^D(u,v)$, between two vertices *u,v* of a connected graph *G* is defined as 0 if $u=v$ and $\max\left\{l_s^D(u,v)\right\}$, if $u \neq v$ where the maximum is taken over all $u-v$ paths *s* in *G* and $l^D(u,v)$ denotes the *D*-length of the paths which is given by $l_s{}^D(u,v)=l(s)$ + sum of the degrees of all vertices present in s. Using this distance we can define *D*-eccentricity of vertices and hence *D*-radius and *D*-diameter in a natural way.

A graph *G* is said to be *D-self centered* if *D*-radius and *D*- diameter of *G* are equal. In otherwords *G* is *D*- self centered if the *D*-eccentricities of all vertices are equal. The *D*-eccentricity of a vertex is denoted by $e^D(v)$ and radius, diameter are denoted by $r^D(G)$ and $dia^D(G)$ respectively.

In the present article we study the problem of a total graph being *D*-self centered. Throughout the article, we denote by x_{ij} the edge joining the vertices v_i and v_j in a graph *G*. All graphs we consider are finite and connected simple graphs.

2. Radius and Diameter of Some Total Graphs

In this section, we study the self – centeredness of total graph w.r.t. *D*-distance. We begin with a theorem.

Theorem 2.1
Consider the star graph $G=St_{n,1}$ with *n*+1 vertices. The *D*-radius and *D*-diameter of $T(St_{n,1})$ are $r^D(G)=2n+6$, $dia^D(G)=3n+2$ for $n \geq 4$. For $n=3, r^D(G)=11$ and $dia^D(G)=12$.

Proof: First we consider the case where $n \geq 4$. Consider the star graph with vertex set $\{v_0, v_1, v_2, \cdots v_n\}$ where v_0 is the central vertex and edge set is $\{x_{01}, x_{02}, \cdots x_{0,n-1}, x_{0n}\}$ where $x_{0,i}=v_0v_i$. The degree of each vertex is $d(v_i)=1$ for $i=1,2,\cdots,n$ and $d(v_0)=n$.

In the total graph of *G*, $T\left(St_{n,1}\right)=G^T$. The vertex set is $\begin{Bmatrix} v_0, v_1, v_2, \cdots, v_n, \\ x_{0,1}, x_{0,2}, \cdots, x_{0,n-1}, x_{0,n} \end{Bmatrix}$. In G^T, degree $d(v_i)=2$ for $i=1,2,\cdots,n$ and $d(v_0)=2n$ and $d(x_{0,i})=n+1$ for $i=1,2,\cdots,n$. Further we can see that the *D*-distances in G^T are as follows: for $1 \leq i, j \leq n$,

$$\max\left\{d^D(v_i,v_j)\right\}=2n+6,$$
$$\max\left\{d^D(v_0,v_i)\right\}=2n+3$$
$$\max\left\{d^D(v_i,x_{0,j})\right\}=2n+6,$$
$$\max\left\{d^D(v_i,x_{0,i})\right\}=n+4$$
$$\max\left\{d^D(x_{0,i},x_{0,j})\right\}=2n+3,$$
$$\max\left\{d^D(v_0,x_{0,i})\right\}=3n+2.$$

Copyright © 2018 Authors. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

From this it is clear that the eccentricities of vertices in G^T, are as follows:

$e^D(v_i) = 2n+6, e^D(v_0) = 3n+2,$

$e^D(x_{0,i}) = 3n+2 \text{ for } i = 1,2,\cdots,n.$

$r^D\{G^T\} = r^D\{T(St_{n,1})\} = 2n+6.$

Thus $dia^D\{G^T\} = dia^D\{T(St_{n,1})\} = 3n+2,$

$\text{for } n \geq 4$

Next we consider the case of star graph with 4 vertices $St_{3,1}$. The *D*-distance between vertices of $T(St_{3,1})$ are as follows:

Table 1: *D*-distances in $T(St_{3,1})$

Vertex	v_0	v_1	v_2	v_3	$x_{0,1}$	$x_{0,2}$	$x_{0,3}$	Ecc
v_0	0	9	9	9	11	11	11	11
v_1	9	0	12	12	7	12	12	12
v_2	9	12	0	12	12	7	12	12
v_3	9	12	12	0	12	12	7	12
$x_{0,1}$	11	7	12	12	0	9	9	12
$x_{0,2}$	11	12	7	12	9	0	9	12
$x_{0,3}$	11	12	12	7	9	9	0	12

From this table it is clear that

$r^D\{T(St_{3,1})\} = 11 \quad \text{and} \quad dia^D\{T(St_{3,1})\} = 12.$

Corollary 2.2

The total graph of $St_{4,1}$ *is D-self centered.*

Proof: From theorem 2.1 we get that for $n = 4$, $r^D\{T(St_{4,1})\} = 14 = dia^D\{T(St_{4,1}\}$,

Hence this graph *D*-self centered.

Remark 2.3:

From the above, it is clear that $St_{n,1}^T$ is not *D*-self centered except for $n = 4$. From the proposition 3.12 of [3], we can see that $St_{n,1}$ is also not *D*-self centered, because

$r^D\{St_{n,1}\} =$

$dia^D\{St_{n,1}\} - 2 = n+2.$

Theorem 2.4

The complete graph K_n $(n \geq 3)$ *and its total graph* $T(K_n)$, *both are D-self centered for* $n \geq 3$.

Proof: Let *G* be the complete graph K_n for $n \geq 3$. Then G^T denotes the total graph of K_n. We know that *G* has *n* vertices and $\frac{n(n-1)}{2}$ edges. Hence G^T has $\frac{n^2+n}{2}$ vertices and the number of edges in G^T is $\frac{n(n^2-1)}{2}$.

Let $\{v_1, v_2 \cdots v_n\}$ be the vertices of $G = K_n$. We can easily see that degree of each vertex in G^T is $2(n-1)$, i.e., $d(v_i) = d(x_{i,j}) = 2(n-1)$. Further, the *D*-distance between vertices in G^T are as follows: for $1 \leq i, j \leq n,$

$\max\{d^D(v_i, v_j)\} = 4n-3,$

$\max\{d^D(v_i, x_{i,k})\} = 4n-3,$

$\max\{d^D(v_i, x_{j,k})\} = 6n-4.$

Therefore the *D*-eccentricities of each vertex is

$e^D(v_i) = e^D(x_{j,k}) = 6n-4.$ Thus

$r^D\{T(K_n)\} = dia^D\{T(K_n)\} = 6n-4.$ Hence $T(K_n)$ is *D*-self centered.

Further from proposition 3.7 of [3] we have, for $n \geq 3$,

$r^D\{K_n\} = dia^D\{K_n\} = 2n-1.$ Hence K_n is *D*-self centered.

Next, we look at the path graph.

Theorem 2.5

Let P_n $(n \geq 3)$ *be the path graph with n vertices. Then we have*

$$r^D\{P_n^T\} = \begin{cases} \frac{5n+2}{2} & \text{if } n \text{ is even} \\ \frac{5n+1}{2} & \text{if } n \text{ is odd} \end{cases} \quad \textit{and}$$

$dia^D\{P_n^T\} = 5(n-1) \text{ for all } n.$

Proof: Consider the path graph P_n $(n \geq 3)$ with vertex set $\{v_1, v_2, \cdots, v_n\}$ and edge set $\{x_{1,2}, x_{2,3}, \cdots, x_{n-1,n}\}$, degree of each vertex is 2 except end vertices and $d(v_i) = d(v_n) = 1$. The total graph of P_n, P_n^T, is of order $(2n-1, 4n-5)$. The degrees of $v_2, v_3 \cdots v_{n-1}$ are 4, degrees of v_1 and v_n are 2, degrees of $x_{2,3}, x_{3,4}, \cdots, x_{n-2,n-1}$ are 4 and degrees of $x_{1,2}$ *and* $x_{n-1,n}$ are 3.

To compute the eccentricities of vertices in P_n^T, we consider the following two cases separately, namely, *n* is even and odd.

Case (i)

n is odd, say $n = 2m+1$.

In this case the *D*-distances between vertices are presented in the table 3.

From this, we can see that *D*-eccentricities are

$\{10m, 10m-2, 10m-7, \cdots, 5m+8,$

$5m+3, \cdots, 10m-2, 10m, 10m,$

$10m-4,10m-9,\cdots,10m-4,10m\}$. Thus the maximum *D*-eccentricity of P_n^T is 10*m* and minimum *D*-eccentricity is 5*m*+3. Hence the *D*-radius is 5*m*+3 and *D*-diameter is 10*m*.

Case (ii):

n is even say $n=2m$

In this case, the *D*-distances in P_n^T are as shown in table 4.

From this table , we can see that *D*-eccentricities of vertices in the graph P_n^T are

$$\left.\begin{array}{l}\{10m-5,10m-7,10m-12,\cdots,\\ 10m-7,10m-5,10m-5,\cdots,\\ 10m-9,10m-14,\cdots,5m+1,\\ \cdots,10m-8,10m-5\end{array}\right\}.$$ Thus the maximum *D*-eccentricity of P_n^T is $10m-5$ and minimum *D*-eccentricity of P_n^T is 5*m*+1. Hence the *D*-radius of P_n^T is 5*m*+1 and *D*-diameter of P_n^T is $10m-5$, if *n* is even.

Therefore $r^D\left\{P_n^T\right\}=\begin{cases}5m+1 & \text{if } n=2m\\ 5m+3 & \text{if } n=2m+1\end{cases}$

and

$$dia^D\left\{P_n^T\right\}=\begin{cases}10m-5 & \text{if } n=2m\\ 10m & \text{if } n=2m+1\end{cases}$$

In other words, we have

$$r^D\left\{P_n^T\right\}=\begin{cases}\dfrac{5n+2}{2} & \text{if } n \text{ is even}\\ \dfrac{5n+1}{2} & \text{if } n \text{ is odd}\end{cases}$$

and $dia^D\left\{P_n^T\right\}=5(n-1)$ for $n\geq 3$.

Proposition 2.6:

We have $r^D(P_2^T)=dia^D(P_2^T)=5.$

Proof: The vertex set in P_2^T is $\{v_1,v_2,x_{1,2}\}$ and degree of each vertex is 2. The *D*- distances in P_2^T are as follows :

Table 2: D-distances in P_2^T

Vertex	v_1	v_2	$x_{1,2}$	Ecc
v_1	0	5	5	5
v_2	5	0	5	5
$x_{1,2}$	5	5	0	5

From this table, it follows that the *D*-eccentricities are $e^D(v_1)=e^D(v_2)=e^D(x_{1,2})=5.$

Hence $r^D(P_2^T)=dia^D(P_2^T)=5$. Therefore P_2^T is *D*-self centered .

Remark 2.7

From the above it is clear that the total graph of path graph P_n is *D* -self centered only for *n* =2. It is not *D*-self centered for any $n\geq 3$. Also proposition 3.8 of [3] it follows that P_n for $n\geq 3$ is also not *D*-self centered and P_2 is D-self centered.

3. Conclusions

We begin this section with the example of a graph *G* for which total graph $T(G)$ is *D*-self centered but the graph *G* is not *D*-self centered.

Example 3.1

Consider the (4,5) graph *G* as shown below




fig 1

For this graph *G* , we have the *D*-eccentricities of vertices are as follows:

Table 5: D-distances in graph G(fig 1)

vertices	v_1	v_2	v_3	v_4	Ecc
v_1	0	6	7	6	7
v_2	6	0	6	9	9
v_3	7	6	0	6	7
v_4	6	9	6	0	9

From the table we can conclude that $r^D(G)=7$, $dia^D(G)=9$. Hence *G* is not *D*-self centered.

Next , consider the total graph $T(G)$ of *G* which is of order (9,23). In this the D-distances between vertices are as follows:



Total graph of G
fig 2

Table 6: D-distanecs in graph $T(G)$

T(G)	v_1	v_2	v_3	v_4	$x_{1,2}$	$x_{2,3}$	$x_{3,4}$	$x_{1,4}$	$x_{1,3}$	ecc
v_1	0	11	13	11	12	17	17	11	13	17
v_2	11	0	11	16	10	10	16	16	17	17

v_3	13	11	0	11	17	12	12	17	13	17
v_4	11	16	11	0	16	16	10	10	17	17
$x_{1,2}$	12	10	17	16	0	11	17	11	12	17
$x_{2,3}$	17	10	12	16	11	0	11	17	12	17
$x_{3,4}$	17	16	12	10	17	11	0	11	12	17
$x_{1,4}$	11	16	17	10	11	17	11	0	12	17
$x_{1,3}$	13	17	13	17	12	12	12	12	0	17

From this table, it follows that $e^D(v_i)=17$ for every vertex *v* of $T(G)$ and hence $T(G)$ is *D*- self centered .

Example 3.2
(see example 2.4 of [2]). Consider the (5,6) graph *G* as shown below. This graph *G* is *D*-self centered but its total graph $T(G)$ is not *D*-self centered.




fig 3

From the above examples and the results of section 2, we see that the following various cases are possible:
Case 1: A graph and its total graph both are *D*-self centered.
Case 2: A graph and its total graph both are not *D*-self centered.
Case 3: Among the graph and its total graph only one is *D*-self centered and the other is not.
Thus we conclude the article with the following open problems.

Open Problem 1.
Characterize all the graphs for which the total graph is D-self centered .

Open Problem 2.
Characterize all the graphs for which the graph and its total graph both are D-self centered .

Open Problem 3.
Characterize all the graphs for which its total graph is D-self centered implies the graph is also D-self centered

References

[1] Frank Harary (2001), Graph theory, Narosa Publishing House, New Delhi.
[2] Ramanjaneyulu M.V, Varma P.L.N, and Reddy Babu D (2017), Self Centered total graphs w.r.t. D-distance, Proceedings of the DST-SERB sponsored National Conference Essence of mathematics and Engineering Applications, KLEF , Vaddeswaram, 22-25 (ISBN No. 9788193063880)
[3] Reddy Babu D, Varma P.L.N. (2013)D-distance in graphs, Golden Research Thoughts, 2(9), 53-58.
[4] Reddy Babu D. and Varma P. L. N., (2017)D-Radius and D-diameter of some Families of graphs, Int. Jour. of Chem. Sci., 15(1), 1-9.

Table 3: D-distance in total graph of P_n (n odd)

	v_1	v_2	⋯	v_n	v_{n+1}	⋯	v_{2n+1}	$x_{1,2}$	$x_{2,3}$	⋯	$x_{2n-1,2n}$	$x_{2n,2n+1}$	Ecc
v_1	0	7	⋯	5*m*-3	5*m*+2	⋯	10*m*-3	6	11	⋯	10*m*-4	10*m*	10*m*
v_2	7	0	⋯	5*m*-6	5*m*-1	⋯	10*m*-3	8	9	⋯	10*m*-6	10*m*-2	10*m*-2
⋮	⋮	⋮	⋯	⋮	⋮	⋯	⋮	⋮	⋮	⋯	⋮	⋮	⋮
v_n	5*m*-3	5*m*-11	⋯	0	9	⋯	5*m*+7	5*m*-2	5*m*-6	⋯	5*m*+4	5*m*+8	5*m*+8
v_{n+1}	5*m*+2	5*m*-1	⋯	9	0	⋯	5*m*+2	5*m*+3	5*m*-1	⋯	5*m*-1	5*m*+3	5*m*+3
⋮	⋮	⋮	⋯	⋮	⋮	⋯	⋮	⋮	⋮	⋯	⋮	⋮	⋮
v_{2n+1}	10*m*	10*m*-3	⋯	5*m*+7	5*m*+2	⋯	0	10*m*	10*m*-4	⋯	11	6	10*m*
$x_{1,2}$	6	8	⋯	5*m*-2	5*m*+3	⋯	10*m*	0	8	⋯	10*m*-7	10*m*-3	10*m*
$x_{2,3}$	11	9	⋯	5*m*-6	5*m*-1	⋯	10*m*-4	8	0	⋯	10*m*-11	10*m*-7	10*m*-4
⋮	⋮	⋮	⋯	⋮	⋮	⋯	⋮	⋮	⋮	⋯	⋮	⋮	⋮
$x_{2n-1,2n}$	10*m*-4	10*m*-6	⋯	5*m*+4	5*m*-1	⋯	11	10*m*-7	10*m*-11	⋯	0	8	10*m*-4
$x_{2n,2n+1}$	10*m*	10*m*-2	⋯	5*m*+8	5*m*+3	⋯	6	10*m*-3	10*m*-7	⋯	8	0	10*m*

Table 4: D-distance in total graph of P_n (n even)

	v_1	v_2	⋯	v_{2n-1}	v_{2n}	$x_{1,2}$	$x_{2,3}$	⋯	$x_{n,n+1}$	⋯	$x_{2n-2,2n-1}$	$x_{2n-1,2n}$	Ecc
v_1	0	7	⋯	10m-8	10m-5	6	11	⋯	5m+1	⋯	10m-8	10m-5	10m-5
v_2	7	0	⋯	10m-11	10m-8	8	9	⋯	5m-1	⋯	10m-11	10m-7	10m-7
⋮	⋮	⋮	⋯	⋮	⋮	⋮	⋮	⋯	⋮	⋯	⋮	⋮	⋮
v_{2n-1}	10m-8	10m-11	⋯	0	6	10m-7	10m-11	⋯	5m-1	⋯	9	8	10m-7
v_{2n}	10m-5	10m-8	⋯	6	0	10m-5	10m-9	⋯	5m+1	⋯	11	6	10m-5
$x_{1,2}$	6	8	⋯	10m-7	10m-5	0	8	⋯	5m-2	⋯	10m-2	10m-8	10m-5
$x_{2,3}$	11	9	⋯	10m-11	10m-9	8	0	⋯	5m-6	⋯	10m-16	10m-12	10m-9
⋮	⋮	⋮	⋯	⋮	⋮	⋮	⋮	⋯	⋮	⋯	⋮	⋮	⋮
$x_{n,n+1}$	5m+1	5m-1	⋯	5m-1	5m+1	5m-2	5m-6	⋯	0	⋯	5m-6	5m-2	5m+1
⋮	⋮	⋮	⋯	⋮	⋮	⋮	⋮	⋯	⋮	⋯	⋮	⋮	⋮
$x_{2n-2,2n-1}$	10m-8	10m-11	⋯	9	11	10m-12	10m-16	⋯	5m-6	⋯	0	8	10m-8
$x_{2n-1,2n}$	10m-5	10m-7	⋯	8	6	10m-6	10m-12	⋯	5m-2	⋯	8	8	0