International Journal of Engineering & Technology, 7 (4.10) (2018) 610-612

International Journal of Engineering & Technology

Website: www.sciencepubco.com/index.php/IJET

Research paper



Numerical Solution of Time Fractional Gas Dynamics Equation

T.R.Ramesh Rao[1,*], R.Santosh Balaji[2]

[1]*B.S.Abdur Rahman Crescent Institute of Science and Technology, Chennai, India 600048.*
[2]*Rajalakshmi Engineering College, Chennai, India 602105.*
[*]*Corresponding author E-mail:rameshrao@crescent.education*

Abstract

In this paper, we propose new technique for solving time fractional gas dynamics equation using Laplace-Adomian Decomposition method coupled with fractional complex transform. It is found that the tested examples reveal the present method is more reliable and does not require strong assumptions.

Keywords: *Gas Dynamics equation, Laplace-Adomian decomposition method, Fractional derivative, Fractional complex transform.*

1. Introduction

In this paper, we consider the time fractional nonlinear gas dynamics equation:

$$\frac{\partial^\alpha u}{\partial t^\alpha} = -\mathrm{u}u_x + \mathrm{ku}(1-\mathrm{u}) + \mathrm{g}(x,t) \qquad \mathrm{t} > 0, \qquad 0 < \alpha \leq 1 \quad (1)$$

Subject to the initial condition

$$u(x,0) = f(x) \quad (2)$$

Where $\mathrm{g}(x,t)$, the source term and k is suitable constant. The different types of gas dynamics equations in mathematical physics have been handled by Evans et al [1] and Elizarova [2]. Numerical solution of nonlinear gas dynamics equations was studied by many researchers. Das and Kumar [4] have applied differential transform method (DTM) to solve the nonlinear gas dynamics equations. Later, the same type of problems was solved by applying fractional homotopy analysis transform method by Rashidi et al [5]. In 2013, Jagdev Singh et al [19] have given the numerical solution of (1) by homotopy perturbation method coupled with Sumudu transform. Aminikanth et al [23] have solved the same type of problem by new homotopy perturbation method via Laplace transform. Recently, Tamsir et al [6] solved the similar type of problems by fractional reduced differential transform method (FRDTM).

For the past two decades, many researchers have paid attention in studying the coupling technique i.e., combination of two analytic methods to solve nonlinear problems The Laplace Adomian-Decomposition method is one such technique introduced by Khuri[16] used to solve the class of nonlinear differential equations. Later many researchers have handled this technique to investigate many nonlinear problems. For example, Wazwaz [13] used Adomian decomposition method coupled with Laplace transform technique for handling nonlinear Volterra integro-differential equations. The similar numerical technique was investigated by Susmita Paul et al [12] for solving Lokta-Volterra Prey Predator model which is dynamic in nature. Ongun [10] and Nurettin Dogan[11] proposed this technique for solving the mathematical model for HIV infection of $CD4^+$Cells. Later the same method has been applied for solving nonlinear fractional diffusion wave equation by Jafari [14] and nonlinear partial differential equation by Arun kumar et al [15].

2. Brief Analysis of Complex Fractional Transform

2.1. Jumaries Fractional Derivatives

Jumaries fractional derivative is a modified Riemann-Liouvilles derivative of order $'\alpha'$ defined as

$$D_t^\alpha f(t) = \begin{cases} \frac{1}{\Gamma(-\alpha)}\int_0^t (t-\zeta)^{-\alpha-1}[f(\zeta)-f(0)]d\zeta & \alpha < 0 \\ \frac{1}{\Gamma(-\alpha)}\frac{d}{dt}\int_0^t (t-\zeta)^{-\alpha-1}[f(\zeta)-f(0)]d\zeta, & 0 < \alpha < 1 \\ \left[f^{(\alpha-m)}(t)\right]^{(m)}, & m \leq \alpha \leq m+1, \quad m \geq 1 \end{cases} \quad (3)$$

where $f: R \to R$ is a continuous function.
We list some important properties for the modified Riemann-Liouvilles derivatives as follows:

$$(i) D_t^\alpha(c) = 0, \alpha > 0, c \text{ is a constant} \quad (4)$$

$$(ii) D_t^\alpha[cf(t)] = cD_t^\alpha f(t), \alpha > 0 \quad (5)$$

$$(iii) D_t^\alpha t^\beta = \frac{\Gamma(1+\beta)}{\Gamma(1+\beta-\alpha)} t^{\beta-\alpha}, \ \beta > \alpha > 0 \quad (6)$$

$$(iv) D_t^\alpha[f(t)g(t)] = [D_t^\alpha f(t)]g(t) + f(t)[D_t^\alpha g(t)] \quad (7)$$

$$(v) D_t^\alpha\left[f\left(h(t)\right)\right] = f_h'\left(h(t)\right) D_t^\alpha h(t) \quad (8)$$

2.2 Fractional Complex Transform Method

The fractional complex transform is the simplest approach in fractional calculus which converts the fractional differential equation

Copyright © 2018 Authors. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

into integer order differential equation making the solution procedure extremely simple [19, 20, 21, 22].The fractional complex transform was
first proposed by Zeng Biao Li et al [20] and is defined as

$$T = \frac{at^{\alpha}}{\Gamma 1+\alpha} \tag{9}$$

$$\mathrm{X} = \frac{bx^{\beta}}{\Gamma 1+\beta} \tag{10}$$

$$Y = \frac{cy^{\gamma}}{\Gamma 1+\gamma} \tag{11}$$

$$Z = \frac{dz^{\delta}}{\Gamma 1+\delta} \tag{12}$$

Where a, b, c, d are unknown constants and $0 \leq \alpha, \beta, \gamma, \delta \leq 1$

3. Basic Idea of Laplace-Adomian Decomposition Method

In this section, we discuss the general procedure for numerical treatment of initial value problem of fractional gas dynamics equation (1)
Applying the complex transformation [31], we get the following nonlinear differential equation:

$$\frac{\partial u}{\partial T} = -\mathrm{u}u_x + \mathrm{ku}(1-\mathrm{u}) + \mathrm{g}(x,T) \;\; \mathrm{T} > 0, 0 < \alpha \leq 1 \tag{13}$$

Applying Laplace transform on both sides of (13) and using differential property, we have

$$L(u) = \frac{1}{s}L(ku^2 - uu_x) - \frac{k}{s}L(u) + \frac{1}{s}L[g(x,T) + f(x)] \tag{14}$$

According to Adomian decomposition method, the solution u (x, T) can be expanded as an infinite series

$$u(x,,T) = \sum_{n=0}^{\infty} u_n(x,T) \tag{15}$$

Where u_n for $n \geq 0$ can be computed recursively and $ku^2 - uu_x$ is decomposed as

$$ku^2 - uu_x = \sum_{n=0}^{\infty} A_n \tag{16}$$

where $A_n,\ n = 0,1,2 \ldots$ are Adomian polynomials of $u_0, u_1, u_2, \ldots$ defined by

$$A_n = \frac{1}{n!}\frac{d^n}{d\lambda^n}\left[k\left(\sum_{i=0}^{\infty}\lambda^i u_i\right)^2 - \left(\sum_{i=0}^{\infty}\lambda^i u_i\right)\left(\sum_{i=0}^{\infty}\lambda^i u_{i_x}\right)\right]_{\lambda=0} \tag{17}$$

The first few Adomian polynomials are given by

$$A_0 = ku_0^2 - u_0 u_{0_x} \tag{18}$$

$$A_1 = 2ku_0u_1 - u_0u_{1_x} - u_1u_{0_x} \tag{19}$$

$$A_2 = 2ku_0u_2 + ku_1^2 - u_0u_{2_x} - u_1u_{1_x} - u_2u_{0_x} \tag{20}$$

$$A_3 = 2ku_0u_3 + 2ku_1u_2 - u_0u_{3_x} - u_1u_{2_x} - u_2u_{1_x} - u_3u_{0_x} \tag{21}$$

And so on.
Now, applying inverse Laplace transform to both sides of (14) and using (15) and (16), we have

$$\sum_{n=0}^{\infty} u_n = L^{-1}\left[\frac{1}{s}L\left(\sum_{n=0}^{\infty} A_n\right)\right] - kL^{-1}\left[\frac{1}{s}L\left(\sum_{n=0}^{\infty} u_n\right)\right] + F(x,T) \tag{22}$$

Where $F(x,T) = L^{-1}\left[\frac{1}{s}L[g(x,T) + f(x)]\right]$
The Adomian Decomposition presents the recursive relation

$$u_0 = F(x,T) \tag{23}$$

$$u_{n+1} = L^{-1}\left[\frac{1}{s}L(A_n)\right] - kL^{-1}\left[\frac{1}{s}L(u_n)\right] n = 0,1,2, \ldots \tag{24}$$

Utilizing the Adomian polynomials (18) - (21) in to the recursive relation (24), we get

$$u_1 = L^{-1}\left[\frac{1}{s}L(A_0)\right] - kL^{-1}\left[\frac{1}{s}L(u_0)\right] \tag{25}$$

$$u_2 = L^{-1}\left[\frac{1}{s}L(A_1)\right] - kL^{-1}\left[\frac{1}{s}L(u_1)\right] \tag{26}$$

$$u_3 = L^{-1}\left[\frac{1}{s}L(A_2)\right] - kL^{-1}\left[\frac{1}{s}L(u_2)\right] \tag{27}$$

4. Numerical Applications

Example 4.1. Consider the time fractional gas dynamics Eqn. (1) with $g(x,t) = 0, k = 1$ and $u(x,0) = f(x) = e^{-x} = u_0$
we utilizing the eqns.(23) and (25) – (27), we can obtain the first few components of series solution:

$$u_0 = e^{-x} \tag{28}$$

$$u_1 = Te^{-x} \tag{29}$$

$$u_2 = \frac{T^2}{2!}e^{-x} \tag{30}$$

$$u_3 = \frac{T^3}{3!}e^{-x} \tag{31}$$

Substituting the above components in to the Eqn. (15), we have

$$u(x,T) = e^{-x} + e^{-x}\left(\frac{t^{\alpha}}{\Gamma 1+\alpha}\right) + \frac{e^{-x}}{2}\left(\frac{t^{\alpha}}{\Gamma 1+\alpha}\right)^2 + \frac{e^{-x}}{6}\left(\frac{t^{\alpha}}{\Gamma 1+\alpha}\right)^3 + \cdots \tag{32}$$

That converges to the exact solution
$u(x,t) = e^{-x+t}$ when α = 1
which is the same solution as obtained by DTM[4], FHATM[5], HPTM[7], HPSTM[8], HAM [9], RDTM[17] and LTNHPM [23] .
Example 4.2. Consider the nonhomogeneous fractional gas dynamics equation (1) with $g(x,t) = -e^{t-x}$, $k = 1$ and $u(x,0) = f(x) = 1 - e^{-x}$
Utilizing the eqns. (23) and (25) – (27), we can obtain the first few components of series solution:

$$u_0 = 1 - e^{-x+T} \tag{33}$$
$$u_1 = u_2 = u_3 = \cdots = 0$$

Substituting the above components in to the Eqn. (15), we have

$$u(x,t) = 1 - e^{-x+\left(\frac{t^{\alpha}}{\Gamma 1+\alpha}\right)} \tag{34}$$

that converges to the exact solution
$u(x,t) = e^{-x+t}$ when α = 1
which is exactly the same as the result obtained by
DTM [4], FRDTM [6], RDTM[17]and LTNHPM [23].
Example 4.3. Consider the nonlinear fractional gas dynamics equation (1) with $k = loga$
$g(x,t) = 0$and $u(x,0) = f(x) = a^{-x}$
By utilizing the Eqns. (13) and (15) – (18), we can obtain the first few components of series solution given by

$$u_0 = a^{-x} \tag{35}$$

$$u_1 = Ta^{-x}loga \tag{36}$$

$$u_2 = \frac{a^{-x}(T\ loga)^2}{2!} \tag{37}$$

$$u_3 = \frac{a^{-x}(T\ loga)^3}{3!} \tag{38}$$

And so on
Finally, substituting the above components in to the Eqn. (15), we Have

$$u(x,T) = a^{-x} + a^{-x}loga\left(\frac{t^\alpha}{\Gamma 1+\alpha}\right) + \frac{a^{-x}}{2!}(loga)^2\left(\frac{t^\alpha}{\Gamma 1+\alpha}\right)^2 + \frac{a^{-x}}{3!}(loga)^3\left(\frac{t^\alpha}{\Gamma 1+\alpha}\right)^3 + \cdots \tag{39}$$

that converges to the exact solution
$u(x,t) = e^{-x+t}$ when $\alpha = 1$
which is exactly the same as the result obtained by
FHATM [5], FRDTM [6], HPTM [7] and RDTM[17].

5. Conclusion

In this paper, Laplace-Adomian decomposition method has been successfully applied to solve time fractional gas dynamics equation via fractional complex transform. The present technique shows great potential to handle fractional nonlinear differential in simple manner, does not require linearization, discretization and small perturbation.

References

[1] Evans DJ & Bulut H (2002), A new approach to the gas dynamics equation: An application of the decomposition method, *International Journal of Computer Mathematics* **79**, 817-822.

[2] Elizarova TG (2009), Quasi-gas dynamics equations, ISBN: 978-3-642-00291-5, Springer Verlag.

[3] Aminikhah H & Ali Jamalian (2013), Numerical Approximation for Nonlinear Gas Dynamic Equation, *International Journal of Partial Differential Equations*, http://dx.doi.org/10.1155/2013/846749

[4] Das S & Kumar R (2011), Approximate analytical solutions of fractional gas dynamics equations, *Applied Mathematics and Computation* **217**, 9905-9915.

[5] Sunil Kumar, Mohammad Mehdi Rashidi, Deepak Kumar & Navid Freidoonimehr (2014), New analytical method for gas dynamics equation arising in shock fronts, *Computer Physics Communications*, http://dx.doi.org/10.1016/j.cpc.2014.03.025.

[6] Mohammed Tamsir & Vineet K. Srivastava (2016), Revisiting the approximate analytical solution of fractional-order gas dynamics equation, *Alexandria Engineering Journal* **55**, 867-874.

[7] Sunil Kumar, HuseyinKocak & AhmetYıldırım (2012), A Fractional Model of Gas Dynamics Equations and its Analytical Approximate Solution Using Laplace Transform, *Verlag der Zeitschrift fur Naturforschung* **67**, 389-396.

[8] Jagadev Singh, Devendra Kumar & Kilicman A (2013), Homotopy Perturbation Method for Fractional Gas Dynamics Equation Using Sumudu Transform, *Abstract and Applied Analysis*, http://dx.doi.org/10.1155/2013/934060

[9] Jafari H, Chun C, Seifi S & Saeidy M (2009), Analytical solution for nonlinear gas dynamics equation by homotopy analysis method, *Appl.Appl.Math*..**4**, 149-154.

[10] Ongun MY(2011), The Laplace Adomian decomposition method for solving a model for HIV infection of $CD4^+$ T cells, *Mathematical and Computer Modeling* **53**,597-603.

[11] NurettinDogan (2012), Numerical treatment of the model for HIV infection of CD4+ Tcells by using multistep Laplace Adomian Decomposition method, *Discrete Dynamics in nature and Society*, http://dx.doi.org/10.1155/2012/976352

[12] Susmita Paul, Sankar Prasad Mondal & Paritosh Battacharya (2016), Numerical solution of Lokta-Volterra Prey Predator Model Runge-Kutta Felhberg method and Laplace Adomian Decomposition method, *Alexandria Engineering Journal* **55**, 613-617.

[13] Abdul-MajidWazwaz (2010), The combined Laplace transform-Adomian Decomposition method for handling Volterraintegro-differential equations, *Applied Mathematics and Computation* **216**, 1304-1309.

[14] Jafari M, Khalique CM & Nazari M (2011), Application of the Laplace decomposition method method for solving linear and nonlinear fractional diffusion-wave equations, *Applied Mathematics Letters* **24**, 1799-1805.

[15] Arunkumar & Ram Dayal Pankaj (2012), Laplace Decomposition method to study wave solutions of coupled nonlinear partial differential equation, *Computational Mathematics*, Article ID 423469.

[16] Khuri SA (2001), A Laplace decomposition algorithm applied to class of nonlinear differential equations, *J. Math. Appl*. **4**, 141-155.

[17] Ramesh Rao T R (2017), Numerical solution for time fractional gas dynamics equations through reduced differential transform method coupled with fractional complex transform, *International Journal of Pure and Applied Mathematics* **117**, 111-118.

[18] Ji-Huan He, Elagan SK & Li ZB (2012), Geometrical explanation of the fractional complex transform and derivative chain rule for fractional calculus, *Physics Letters A* **376**, 257-259.

[19] Guy Jumarie (2007), Fractional partial differential equations and modified Riemann-Liuville derivatives new methods for solution, *J.Appl.Math.& Comp*. **24**, 31-48.

[20] Zeng Biao Li & Ji-Juan He (2010), Fractional complex transform for fractional differential equation, *Mathematical and Computational Applications* **15**, 970-973.

[21] Saha Ray S & Sahoo S (2015), A Novel Analytical Method with fractional complex transform for new exact solutions of time-fractional fifth-order Swada-Kotera Equation, *Reports on Mathematical Physics* **75**, 65-72.

[22] Zeng Biao Li & Ji-Juan He (2010), Fractional complex transform for fractional differential equation, *Mathematical and Computational Applications* **15**, 970-973.

[23] Hossein Aminihkah & Ali Jamalian (2013), Numerical Approximation for Nonlinear Gas Dynamics Equations, *International Journal of Partial differential Equations*, http://dx.doi.org/10.1155/2013/846749.