International Journal of Engineering & Technology, 7 (4) (2018) 2925-2927

International Journal of Engineering & Technology

Website: www.sciencepubco.com/index.php/IJET
doi: 10.14419/ijet.v7i4.18041
Research paper

Inverse majority vertex covering number of a graph

Paulraj Jayasimman I [1] *, Dhivya T [1], Joseline Manora J [2]

[1] *AMET Deemed to be University, Kanathur, Chennai*
[2] *TBML College, Porayar, Nagappattinam, Tamilnadu, India*
**Corresponding author E-mail: ipjayasimman@gmail.com*

Abstract

A set of vertices$S$, which covers atleast half of the edges is a Majority vertex cover of G. The majority vertex covering number $\alpha_M(G)$ of G is the minimum number in a Majority vertex cover. In this paper, new parameter has been introduced Inverse majority vertex covering number of a graph G with respect to Majority vertex covering set. Also majority vertex covering number obtained for classic graphs and Cartesian product graph.

Keywords: *Majority Vertex Covering Number; Inverse Majority Vertex Set; Inverse Majority Vertex Covering Number of A Graph.*

1. Introduction

Definition 1.1 [4]:

A subset S of V(G) is said to be a majority dominating set (MDS) if at least half of the vertices of V(G) are either in S or adjacent to elements of S. A majority dominating set S is minimal if no proper subset of S is a majority dominating set. The minimum cardinality of a minimal majority dominating set is called majority domination number, denoted by $\gamma_M(G)$.

Definition 1.2: *[5] A vertex set v is called majority dominating vertex if the degree of v, if $d(v) \geq \left\lceil \frac{p}{2} \right\rceil - 1$. For $S \subseteq V(G)$, a vertex $v \in S$ is called an enclave of S if $N[v] \subseteq S$ and $v \in S$ is an isolate of S if $N(v) \subseteq V - S$.*

Definition 1.3: *[10] Let D be minimum vertex cover of G. If $V - D$ contains a vertex cover D' of$G$, then $D'$is called an inverse vertex cover with respect to D. The inverse vertex covering number $\alpha_0^{-1}(G)$ of G is the minimum cardinality of an inverse vertex cover of G.*

Definition 1.4: *[6] A set of vertices S, which covers at least half of the edges, is a majority vertex cover of$G$. The majority vertex covering number $\alpha_M(G)$ of G is the minimum number of vertices in a majority vertex cover.*

2. Inverse majority covering number of a graph

Definition 2.1: *A set D_M be the majority vertex cover of G. If $V - D_M$ contains a majority vertex cover D_M^{-1} of G then D_M^{-1} is called a inverse majority vertex cover with respect to D_M. The inverse majority vertex covering number $\alpha_M^{-1}(G)$ of G is the minimum vertex covering number of G.*

Proposition 2.2

For the graph $G = W_p$ with$p \geq 3$, $\alpha_M^{-1}(G) = \left\lceil \frac{p-1}{3} \right\rceil$.

Proof
Let $G = W_p = K_1 + C_{p-1}$ with $p \geq 3$ and$q = 2(p-1)$. The vertex set $V(G) = \{u, v_1, v_2, \ldots, v_{p-1}\}$. The majority vertex covering number $\alpha_M^{-1}(G) = 1$ since$d(u) = p - 1$. Let D_M be majority vertex coving set. The inverse majority vertex covering set

$$D_M^{-1}(G) = \left\{v_1, v_2, v_3, \ldots v_{\left\lceil \frac{p-1}{3} \right\rceil}\right\} \Rightarrow \langle N[V - D_M]\rangle = C_p$$

Then $\left|D_M^{-1}\right| = \left\lceil \frac{p-1}{3} \right\rceil = \left\lceil \frac{2(q-1)}{3} \right\rceil \geq \left\lceil \frac{q}{2} \right\rceil$

Therefore$\alpha_M^{-1}(G) = \left|D_M^{-1}\right| = \left\lceil \frac{p-1}{3} \right\rceil$.

Proposition 2.3:

For the fan graph $G = F_p$ with$\alpha_M^{-1}(G) = \left\lceil \frac{p-1}{3} \right\rceil$.

Theorem 2.4: *If the graph $G = S_{2m+1}$ is the spider graph then $\alpha_M^{-1}(G) = \left\lceil \frac{m}{2} \right\rceil$.*

Theorem 2.5: *For the friendship graph $G = F_{1,2n}$ then $\alpha_M^{-1}(G) = \left\lceil \frac{p}{2} \right\rceil$.*

Proposition 2.6:
If the graph G is complete bipartite graph $G = K_{m,n}$with $m, n \geq 2$ then

$$n_M^{-1}(G) = \begin{cases} \left\lceil \frac{m}{2} \right\rceil, \text{if } m \leq n \text{ and } m = \text{even and } m - n = 1 \\ \left\lceil \frac{m}{2} \right\rceil + 1, \text{if } m \leq n \text{ and } m = \text{odd} \end{cases}$$

with $m, n \geq 2$.

Copyright © 2018 Pauraj Jayasimman I et. al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

3. Inverse Majority vertex covering number for Grid graph

Definition 3.1: *[3] Let G and H be any two connected graphs with the vertex set* $\{u_1, u_2, u_3, \dots, u_n\}$ *and* $\{v_1, v_2, v_3, \dots, v_n\}$ *respectively. The Cartesian products graph* $K = G \times H$ *has* $V(K) = V(G) \times V(H)$ *and vertices* (u_1, v_1) *and* (u_2, v_2) *in* $V(K)$ *are adjacent if and only if either* $u_1 = u_2$ *and* v_1v_2 *in* $V(H)$ *or* $v_1 = v_2$ *and* u_1u_2 *in* $E(G)$*. The grid graph is* $P_i \times P_j$ *and the cylinder is* $C_i \times P_j$ *for* $i, j \geq 3$.

Theorem 3.2: *If the graph* $G = P_2 \times P_j$ *is grid graph with* $j \geq 3$ *then* $\alpha_M^{-1}(G) = \left\lfloor \frac{j+1}{2} \right\rfloor$.

Proof

Consider the graph $G = P_2 \times P_j$ is ladder graph with vertex set $\{v_{11}, v_{12}, v_{13}, \dots, v_{1j}\}$ and the $\{v_{21}, v_{22}, v_{23}, \dots, v_{2j}\}$, $\Delta(G) = 3$, $d(v_{i1}) = d(v_{ij}) = 2$ and $p = 2j$, $q = 2(j-1) + j$.

Case (i) $j =$ even Let the majority vertex covering set is $D_M^{-1} = \{v_{13}, v_{15}, v_{17}, \dots, v_{1j-1}, v_{22}\}$with $d(v_i, v_j) = 2$

$\Rightarrow |D_M^{-1}| = \left\lceil \frac{j-2}{2} \right\rceil + 1$and $|\langle D_M^{-1} \rangle| = 3\left(\left\lceil \frac{j-2}{2} \right\rceil + 1\right) > \left\lceil \frac{2(j-1)+j}{2} \right\rceil = \left\lceil \frac{q}{2} \right\rceil$.

Therefore, $|D_M^{-1}| \leq \alpha_M^{-1}(G)\alpha_M^{-1}(G)$.

Suppose $D_M^{-1} = \{v_{13}, v_{15}, v_{17}, \dots, v_{1j-1}\} \Rightarrow |D_M^{-1}| = \left\lceil \frac{p-2}{2} \right\rceil$

and $|\langle D_M^{-1} \rangle| = 3\left(\left\lceil \frac{j-2}{2} \right\rceil\right) < \left\lceil \frac{2(j-1)+j}{2} \right\rceil = \left\lceil \frac{q}{2} \right\rceil$.

Therefore, $|D_M^{-1}| \geq \alpha_M^{-1}(G)$. Hence, $\alpha_M^{-1}(G) = \left\lceil \frac{j-2}{2} \right\rceil + 1 = \left\lfloor \frac{j+1}{2} \right\rfloor$.

Case (ii) $j =$ odd

If $j = 3$ then $D_M^{-1} = \{v_{11}, v_{22}\} \Rightarrow |D_M^{-1}| = \left\lceil \frac{j-2}{2} \right\rceil + 1$ and $|\langle D_M^{-1} \rangle| = 3\left(\left\lceil \frac{j-2}{2} \right\rceil + 1\right) > \left\lceil \frac{q}{2} \right\rceil$.


If $j > 3$ then $D_M^{-1} = \{v_{13}, v_{15}, v_{17}, \dots, v_{1j-2}, v_{22}, v_{24}\} \Rightarrow |D_M^{-1}| = \left\lceil \frac{j-3}{2} \right\rceil + 2$ and $|\langle D_M^{-1} \rangle| = 3\left(\left\lceil \frac{j-3}{2} \right\rceil + 2\right) > \left\lceil \frac{2(j-1)+j}{2} \right\rceil = \left\lceil \frac{q}{2} \right\rceil$. Therefore, $|D_M^{-1}| \leq \alpha_M^{-1}(G)$.

Suppose

$D_M^{-1} = \{v_{13}, v_{15}, v_{17}, \dots, v_{1j-2}, v_{22}\} \Rightarrow |D_M^{-1}| = \left\lceil \frac{j-3}{2} \right\rceil + 1$

and $|\langle D_M^{-1} \rangle| = 3\left(\left\lceil \frac{j-3}{2} \right\rceil + 1\right) < \left\lceil \frac{2(j-1)+j}{2} \right\rceil = \left\lceil \frac{q}{2} \right\rceil$.

Therefore, $|D_M^{-1}| \geq \alpha_M^{-1}(G)$. Hence, $\alpha_M^{-1}(G) = \left\lceil \frac{j-3}{2} \right\rceil + 2 = \left\lfloor \frac{j+1}{2} \right\rfloor$.



Theorem 3.3: *If the graph* $G = P_3 \times P_j$ *with* $j \geq 4$ *then*$\alpha_M^{-1}(G) = \left\lfloor \frac{3(j-2)}{4} \right\rfloor$.

Proof

Let $G = P_3 \times P_j$ be the grid graph with $V(G) = \{v_{11}, v_{12}, v_{13}, \dots, v_{1j}, v_{21}, v_{22}, \dots, v_{2j}, v_{31}, v_{32}, \dots v_{3j}\}$ then $p = 3j$ and $= 3(2j-1)$, $\Delta(G) = 4$. $V_1(G) = \{v_{11}, v_{12}, v_{13}, \dots, v_{1j}\}$, $V_2(G) = \{v_{21}, v_{22}, v_{23}, \dots, v_{2j}\}$, $V_3(G) = \{v_{31}, v_{32}, v_{33}, \dots, v_{3j}\}$ I, II and III row vertices respectively. $d(v_{2j}) = 4$ and $d(v_{1j}) = d(v_{3j}) = 3$..

Case (i) $j = even$

Let $D_M^{-1}(G) = \{v_{23}, v_{25}, v_{27}, \dots v_{2j-1}, v_{12}, v_{14}, \dots, v_{1j-2}\}$. Then $|D_M^{-1}| = \left\lceil \frac{j-2}{2} \right\rceil + \left\lceil \frac{j}{4} \right\rceil = \left\lceil \frac{3(j-2)}{4} \right\rceil \Rightarrow |\langle D_M^{-1} \rangle| = 4\left\lceil \frac{j-2}{2} \right\rceil + 3\left\lceil \frac{j}{4} \right\rceil > \left\lceil \frac{3(2j-1)}{2} \right\rceil = \left\lceil \frac{q}{2} \right\rceil$. Since $d(v_{2j}) = 4$ and $d(v_{1j}) = d(v_{3j}) = 3$. Therefore $|D_M^{-1}| \leq \alpha_M^{-1}(G)$. If $|D_M^{-1}| = \left\lceil \frac{j-2}{2} \right\rceil + \left\lceil \frac{j}{4} \right\rceil - 1$ then $|\langle D_M^{-1} \rangle| = 4\left\lceil \frac{j-2}{2} \right\rceil + 2\left\lceil \frac{j}{4} \right\rceil < \left\lceil \frac{3(2j-1)}{2} \right\rceil = \left\lceil \frac{q}{2} \right\rceil$. Therefore $|D_M^{-1}| > \alpha_M^{-1}(G)$. Hence $|D_M^{-1}| = \alpha_M^{-1}(G) = \left\lceil \frac{3(j-2)}{4} \right\rceil$.

Case (ii) $j = odd$

Choose set

$D_M^{-1}(G) = \{v_{23}, v_{25}, v_{27}, \dots, v_{2j-2}\} \Rightarrow |D_M^{-1}| = \left\lceil \frac{j-3}{2} \right\rceil + \left\lfloor \frac{j}{4} \right\rfloor = \left\lceil \frac{j-2}{2} \right\rceil + \left\lceil \frac{j}{4} \right\rceil = \left\lceil \frac{3(j-2)}{4} \right\rceil \Rightarrow |\langle D_M^{-1} \rangle| = 4\left\lceil \frac{j-3}{2} \right\rceil + 3\left\lceil \frac{j}{4} \right\rceil > \left\lceil \frac{3(2j-1)}{2} \right\rceil = \left\lceil \frac{q}{2} \right\rceil$.

Therefore $|D_M^{-1}| \leq \alpha_M^{-1}(G)$. Hence $\alpha_M^{-1} = \left\lfloor \frac{3(j-2)}{4} \right\rfloor$.

Corollary 3.4:

For the Grid graph $G = P_4 \times P_j$ with $j \geq 4$. Then $\alpha_M^{-1} = \left\lfloor \frac{p}{8} \right\rfloor$.

3.5. Majority vertex covering number for Cylinder graph

Theorem 3.5.1: *For a Cylinder graph* $= C_3 \times P_j, \alpha_M^{-1}(C_3 \times P_j) = \left\lceil \frac{3(2j-1)}{8} \right\rceil$.


Proof : Consider the graph $G = C_3 \times P_j$ with $j \geq 4$ and $V(G)$ be the vertex set with cardinality $3j$, $E(G)$ be edge set with $|E(G)| = 3j + 3(j-1)$. Let $V_1(G)$, $V_2(G)$ and $V_3(G) \subseteq V(G)$ be the I, II and III row vertices respectively. $d(v_{ij}) = 4$, $v_{ij} \in V_1(G), V_3(G)$, $ij \neq 12, 1j, 21, 2j, 31, 3j$ and $d(v_{ij}) = 3$.

If j is even then the inverse majority vertex covering$D_M^{-1}(G) = \{v_{13}, v_{15}, v_{17}, \dots, v_{32}, v_{34}, \dots v_{ij}\}$, $v_{ij} \in V_1(G), V_3(G) \Rightarrow |D_M^{-1}(G)| = \left\lceil \frac{3(2j-1)}{8} \right\rceil, |\langle N[D_M^{-1}] \rangle| = 4\left\lceil \frac{j-2}{2} \right\rceil + 4\left\lceil \frac{j-2}{2} \right\rceil = 8\left\lceil \frac{j-2}{2} \right\rceil = 4\left\lceil \frac{3(2j-1)}{8} \right\rceil$, $4\left\lceil \frac{3(2j-1)}{8} \right\rceil \geq \left\lceil \frac{3(2j-1)}{2} \right\rceil \geq \left\lceil \frac{q}{2} \right\rceil$. Therefore $|D_M^{-1}(G)| = \left\lceil \frac{3(2j-1)}{8} \right\rceil \leq \alpha_M^{-1}(C_3 \times P_j)$. Suppose $|D_M^{-1}(G)| = \left\lceil \frac{3(2j-1)}{8} \right\rceil - 1 \Rightarrow 4\left\lceil \frac{3(2j-1)}{8} \right\rceil - 1 < \left\lceil \frac{3(2j-1)}{2} \right\rceil - 1 < \left\lceil \frac{q}{2} \right\rceil$. Therefore $|D_M^{-1}(G)| = \left\lceil \frac{3(2j-1)}{8} \right\rceil > \alpha_M^{-1}(C_3 \times P_j)$. Hence $|D_M^{-1}(G)| = \alpha_M^{-1}(C_3 \times P_j) = \left\lceil \frac{3(2j-1)}{8} \right\rceil$. If $j = odd$ then $|D_M^{-1}(G)| = \left\lceil \frac{3(2j-1)}{8} \right\rceil \Rightarrow |\langle N[D_M^{-1}] \rangle| = 4\left\lceil \frac{j-3}{2} \right\rceil + 4\left\lceil \frac{j-2}{2} \right\rceil = 4\left\lceil \frac{3(2j-1)}{8} \right\rceil \geq \left\lceil \frac{3(2j-1)}{2} \right\rceil \geq \left\lceil \frac{q}{2} \right\rceil$.

Therefore $\alpha_M^{-1}(C_3 \times P_j) = \left\lceil \frac{3(2j-1)}{8} \right\rceil$.

3.5.2. Example

Consider the graph $G = C_3 \times P_7$ with $p = 3(7) = 21, q = 3(7) + 3(6) = 39$ then $\left\lceil \frac{q}{2} \right\rceil = \left\lceil \frac{39}{2} \right\rceil = 20$.



$D_M^{-1}(G) = \{v_{13}, v_{15}, v_{22}, v_{24}, v_{26}\}$ is a majority vertex covering set of G

$|D_M^{-1}| = \alpha_M^{-1} = \left\lceil \frac{3(2(7)-1)}{2} \right\rceil$=20.

4. Bounds on Inverse majority vertex covering number

Proposition 4.1:
For any graph G without isolated vertices then

$\alpha_M(G) \leq \alpha_M^{-1}(G)$

This bound is sharp for $K_{m,m}$ with$m \geq 2$.

Proposition 4.2:
For any connected graph $\boldsymbol{G}$ then

$\gamma_M(G) \leq \alpha_M^{-1}(G)$

This bound holds for$G = C_{2n}$.

Proposition 4.3
For any connected graph G,

$\alpha_M(G) + \alpha_M^{-1}(G) = p - 2.$

Proposition 4.4:

For any graph $\boldsymbol{G}$ without isolated vertices ,

$\gamma_M(G) \leq \alpha_M(G) \leq \alpha_M^{-1}(G).$

Proposition 4.5:
For any connected graph $\boldsymbol{G}$ then,

$\alpha_M^{-1}(G) \leq p - \gamma_M(G)$

Observation 4.6:

i) If the graph $G = P_i \times P_j$ then $\alpha_M(G) = \alpha_M^{-1}(G)$
ii) If the graph $G = C_3 \times P_j$ then $\alpha_M(G) = \alpha_M^{-1}(G)$

5. Conclusion

The researcher has introduced the new parameter inverse majority vertex covering of a graph $\boldsymbol{G}$. This number defined $\boldsymbol{\alpha_M^{-1}(G)}$ and determined for some classes of graph and also obtained the bounds. This generalized for Cartesian product graphs

References

[1] E.J. Cockayane and S.T. Hedetniemi, "Towards a theory of domination in graphs", Networks. Seven, pp. 247-261, (1977). https://doi.org/10.1002/net.3230070305.
[2] Frank Harary, Graph theory, Addision-Wesley, Reading, Mass., (1972).
[3] T.W Haynes, S.T Hedetniemi and P.J Slater, Fundamentals of domination in graphs, Marces Dekker. Inc, New york, (1998).
[4] J. Joseline Manora and V. Swaminathan, "Majority Dominating sets", J A R J; Vol.3, No.2, pp.75-82, (2006).
[5] J. Joseline Manora and V. Swaminathan, "Results on Majority Dominating sets", Scientia Magna, Northwest University, Xitan, P.R China, Vol. 7, No. 3, pp. 53-58, (2011).
[6] Joseline Manora .J and Swaminathan .V, Majority neighborhood number of a graph-published in Scientia Magna, Dept. of Mathematics, Northwest Universtiy, Xitan, P.R China – Vol (6), N0.2, 20-25(2010).
[7] Joseline Masnora J and Paulraj Jayasimman. I, Independent Majority Neighborhood Number of a graph, International Journal of Applied Computational Science & Mathematics. Volume 4, Number 1 (2014), pp. 103-112. https://doi.org/10.5121/ijcsa.2014.4110.
[8] Joseline Masnora J and Paulraj Jayasimman. Neighborhood sets polynomial of a graph, International Journal of Applied Mathematical Sciences, ISSN 0973-0176 Volume 6, Number 1 (2013), pp. 91-97.
[9] Kulli.V.R and Kattimani, The Inverse Neighbourhood Number of a graph, South. East.Asian.J. Math. & Math. Sc. Vol.6 No.3 (2008), pp. 23-28.
[10] V.R.Kulli,S.C, Sigarkanthi, Inverse domination in graphs, National Academic Science Letter, Vol.14,1991,pp 473-475.
[11] V.R.Kulli, Inverse vertex covering number of a graph, Journal of Discrete Mathematical Sciences & Cryptography, Vol.15 (2012), No.6, pp.389-393.
[12] E. Sampathkumar and H.B. Walikar, "The connected domination number of a graph", Jour, Math, Phy, Sci. Vol.13 No.6, pp. 607-613 (1979).