



# Determination of Number of New Elements for Sequence of Fuzzy Topographic Topological Mapping

Azrul Azim Mohd Yunus<sup>1\*</sup>, Tahir Ahmad<sup>2</sup>

<sup>1</sup>Faculty of Science and Technology, Universiti Sains Islam Malaysia (USIM), 71800 Nilai, Negeri Sembilan, Malaysia

<sup>2</sup>Ibnu Sina Institute of Fundamental Studies, Universiti Teknologi Malaysia, 81310 UTM Johor Bahru, Johor, Malaysia

\*Corresponding author E-mail: azrulazim@usim.edu.my

## Abstract

Problem involving neuro magnetic inverse problem can be solved with Fuzzy Topographic Topological Mapping (FTTM). FTTM is a model consists of four components and connected by three algorithms. FTTM version 1 and version 2 were designed to present 3D view of an unbounded single current and bounded multicurrent sources, respectively. Several definitions related to sequence of FTTM were introduced by Suhana and the feature, namely the cube of FTTM are developed. In this paper, sequence of FTTM namely  $FTTM_n$  are discussed. Consequently, some theorems are proved to describe the number of the new elements produced from the  $FTTM_n$ . Besides that, the number of new elements produced from combination of sequence of  $FTTM_n$  can be written as a combination from cubes of  $FTTM_n$ .

**Keywords:** Fuzzy Topographic Topological Mapping; Sequence; Cubes of  $FTTM_n$ .

## 1. Introduction

Fuzzy Topographic Topological Mapping (FTTM) is a model designed to solve neuro magnetic inverse problem. There are four components and connected by three algorithms in FTTM. Detail of FTTM can be found in [1]. In general, there are two version of FTTM namely FTTM version 1 and FTTM version 2. Each were designed to present 3D view of an unbounded single current and bounded multicurrent sources, respectively. In 2008, Suhana introduced some definitions on sequence of FTTM. In this paper, the vertices of FTTM are extended. The geometrical features for some finite vertices of FTTM; namely  $FK_n$  are investigated. One of the geometrical features produced from the combination of sequences of FTTM are cubes of  $FTTM_n$  [1]. From cubes of FTTM, one can determine the number of new elements produced from the combination of FTTM.

In general term, FTTM is a 4-tuple of topological spaces homeomorphic to each other [2]. The representation for FTTM are given by the equation

$$FTTM = \{(M, B, F, T) : M \cong B \cong F \cong T\}. \quad (1)$$

Sequential elements of FTTM [1] are present exactly as shown in Figure 1.

FTTM version 1 was developed to present a 3-D view of an unbounded single current source [3, 4] in one angle observation (upper of a head model). There are three algorithms, which link between four components as shown in Figure 2. FTTM version 2 was developed to present a 3-D view of a bounded multi current source [5] in 4 angles of observation (upper, left, right and back of a head model). It contains three algorithms, where all four components of FTTM connected to the three algorithms (refer Figure 3).




Fig. 1: Sequence of  $FTTM_n$




Fig. 2: FTTM version 1

### 1.1. Cube of $FTTM_n$

A cube is a combination of two, three or more FTTM in  $FTTM_n$  (see Figure 4). A cube of FTTM is denoted by  $FTTM_{k/n}$  where k is a combination of k number of terms in  $FTTM_n$ .




Fig. 3: FTTM version 2

Figure 4 shows some combination and the number of cube for some  $FTTM_n$ . In general, a cube of  $FTTM_n$  is defined as follows.

**Definition 1.1:** Sequence of  $FTTM_{k/n}$ , namely  $FTTM_{k/1}$ ,  $FTTM_{k/2}$ ,  $FTTM_{k/3}$ , ... is given recursively by equation

$$FK_{k/n} = \frac{n(n-1)(n-2)(n-3)(n-4)\dots(n-k-1)}{k!}, n \geq 1, k \geq 2. \quad (2)$$

### 1.2. FTTM as Pascal's Triangle

Jamaian [6] found that there exists a relation on the sequences of FTTM and Pascal's Triangle. Jamaian showed that the sequence of  $FTTM_{2/n}$ ,  $FTTM_{3/n}$ , and  $FTTM_{4/n}$  are presented in the third, the fourth, and the fifth main diagonals of Pascal's Triangle respectively as shown in Figure 5.

## 2. Relation between Sequence of $FTTM_n$ and cube of $FTTM_n$

The determination of numbers of elements produced from the extended  $FTTM_n$  proved to be time consuming. For example, one has to list all 36 elements of  $FTTM_4$  in order to find the number of elements. Alternatively, using the conjecture introduced by Jamaian which was later proved by Sayed and Ahmad, the number new elements of  $FTTM_n$  is given by  $n^4 - n$  [7]. In addition, the elements of  $FTTM_n$  can be written as the linear combination of the sequence of cubes of  $FTTM_n$ .

Type \ n	1	2	3	4	5	6
Segment						
$F2_n$	0	1	3	6	10	15
Triangular						
$F3_n$	0	0	1	4	10	20
Square						
$F4_n$	0	0	0	1	5	15
Pentagonal						
$F5_n$	0	0	0	0	1	6
$\binom{n}{5}$	0	0	0	0	1	6

Fig. 4: Cubes of  $FK_n$




Fig. 5: FTTM in Pascal's Triangle

A sequence for  $FTTM_n$  can be written as a combination of the cubes of  $FTTM_n$ . Therefore, a new theorem on elements for sequence of  $FTTM_n$  is deduced. Generally, elements of  $FTTM_n$  are given by

$$FK_n = C_1 \binom{n}{2} + C_2 \binom{n}{3} + C_3 \binom{n}{4} + C_4 \binom{n}{5} + \dots + C_p \binom{n}{k}; k = K. \quad (3)$$

with  $k$  being the number of components i.e. the vertices of  $FTTM_n$ , and  $C_1, C_2, C_3, C_4, \dots, C_p$  are the coefficients for each combination. Finding the coefficient  $C_1, C_2, C_3, C_4, \dots, C_p$  is not trivial.

**Remark:** Since  $FK_n$  contains  $FK_{2/n}, FK_{3/n}, FK_{4/n}, \dots, FK_{k/n}$ , then it is obvious that there could only be a cube with a combination of  $k$  in  $FK_n$ . For combination of  $K + 1$   $FK$  in  $FK_n$ , i.e.  $FK_{K+1/n}$ , this implies that there are  $K + 1$  components in  $FK$  such that  $\{T_1, T_2, T_3, T_4, \dots, T_k, T_{k+1}\}$ . It leads to contradiction since  $FK$  only has  $K$  components which is  $\{T_1, T_2, T_3, T_4, \dots, T_K\}$ . Therefore a cube of  $FK$  can be formed from a combination of  $2, 3, 4, \dots, k$  in  $FK_n$ ; i.e.  $FK_{2/n}, FK_{3/n}, FTTM_{4/n}, FK_{k/n}$ .

From the remark, FTTM was presented as a combination of cubes namely

$$FTTM_n = C_1 \binom{n}{2} + C_2 \binom{n}{3} + C_3 \binom{n}{4} = n^4 - n. \quad (4)$$

## 3. Results and Discussion

The coefficients for  $FTTM_n$  can be found as given in the following theorem.

**Theorem 3.1** The number of elements produced from sequence of  $FTTM_n$  is

$$FTTM_n = 14 \binom{n}{2} + 36 \binom{n}{3} + 24 \binom{n}{4} = n^4 - n. \quad (5)$$

### Proof of Theorem 3.1

$F4_n$  can be written as a system of simultaneous equations given in the following.

$$\begin{aligned}
 x_1 \binom{n}{2} + x_2 \binom{n}{3} + x_3 \binom{n}{4} &= n^4 - n \\
 x_1 \binom{n}{2} + x_2 \binom{n}{3} &= n^4 - n \\
 x_1 \binom{n}{2} &= n^4 - n
 \end{aligned} \tag{6}$$

or

$$\begin{bmatrix} x_1 & x_2 & x_3 \\ 0 & x_1 & x_2 \\ 0 & 0 & x_1 \end{bmatrix} \begin{bmatrix} \binom{n}{2} \\ \binom{n}{3} \\ \binom{n}{4} \end{bmatrix} = \begin{bmatrix} n^4 - n \\ n^4 - n \\ n^4 - n \end{bmatrix}. \tag{7}$$

For  $n = 2$ , in (7) leads to

$$\begin{bmatrix} x_1 & x_2 & x_3 \\ 0 & x_1 & x_2 \\ 0 & 0 & x_1 \end{bmatrix} \begin{bmatrix} \binom{2}{2} \\ \binom{2}{3} \\ \binom{2}{4} \end{bmatrix} = \begin{bmatrix} 14 \\ 14 \\ 14 \end{bmatrix} \tag{8}$$

which imply  $x_1 = 14$ . Next, for  $n = 3$ , in (7) will yield

$$\begin{bmatrix} x_1 & x_2 & x_3 \\ 0 & x_1 & x_2 \\ 0 & 0 & x_1 \end{bmatrix} \begin{bmatrix} \binom{3}{2} \\ \binom{3}{3} \\ \binom{3}{4} \end{bmatrix} = \begin{bmatrix} 78 \\ 78 \\ 78 \end{bmatrix}. \tag{9}$$

Solving the above equation will resulting  $x_2 = 36$ . Similarly, letting  $n = 4$  will give  $x_3 = 24$ . Therefore,

$$FTTM_n = 14 \binom{n}{2} + 36 \binom{n}{3} + 24 \binom{n}{4}.$$

## 4. Conclusion

Homeomorphism between components of  $FTTM_n$  gives many new elements. One can list all the new elements of  $FTTM_n$ . However, it proved to be time consuming. Alternatively, using the conjecture introduced by Jamaian which was later proved by Sayed and Ahmad, the number of new elements of  $FTTM_n$  is given by  $n^4 - n$ . In addition, the elements of  $FTTM_n$  can be written as the linear combination of the sequence of cubes of  $FTTM_n$ . The method to find the coefficient for the sequence of cubes of  $FTTM_n$  is discussed in this paper. An alternative method to find the coefficient for  $FTTM_n$  is also presented.

## Acknowledgement

The authors would like to thank Universiti Sains Islam Malaysia for the financial support receive from short term grant (PPP/USG/0216/FST/30/17216).

## References

- [1] Jamaian S S, Ahmad T, Talib J. Generalized Finite Sequence of Fuzzy Topographic Topological Mapping. *Journal of Mathematics and Statistics*, 2010, 6(2):151–156.
- [2] Ahmad T, Abdullah J, Zakaria F, Mustapha F, Thakaran J, Zabidi-Hussin A. Nonlinear Dynamic Modelling of Brain Storm of Epilepsy Sufferers. *Proceedings of the 7th International Neurotrauma Symposium*, 2004.
- [3] Zakaria F. Algoritma Penyelesaian Masalah Songsang Arus Tunggal Tak Terbatas MEG. Master thesis, Universiti Teknologi Malaysia, 2002.
- [4] Yun L L. Homeomorfisma  $S^2$  Antara  $E^3$  Melalui Struktur Permukaan Riemann Serta Deduksi Teknik Pembuktiannya Bagi Homeomorfisma Pemetaan Topologi Topografi Kabur. Master thesis, Universiti Teknologi Malaysia, 2001.
- [5] Rahman W E Z W A, Ahmad T, Ahmad R S. Simulating the Neuronal Current Sources in the Brain. *Proceedings of the BIOMED*, 2002.
- [6] Jamaian S S. Generalized Fuzzy Topographic Topological Mapping. Master thesis, Universiti Teknologi Malaysia, 2008.
- [7] Ahmad T, Ahmad R S, Yun L L, Zakaria F, Rahman W E Z W A. Homeomorphisms of Fuzzy Topographic Topological Mapping. *Matematika*, 2005, 21(1):35–42.