International Journal of Engineering & Technology, 7 (4.10) (2018) 793-796

International Journal of Engineering & Technology

Website: www.sciencepubco.com/index.php/IJET

Research paper



Oscillation of Generalized Second-Order Quasi Linear Difference Equations

V.Srimanju[1]*, Sk.Khadar Babu[2], V.Chandrasekar[3]

[1,2]Vellore Institute of Technology, Vellore – 632 014, Tamil Nadu, India
[3]Thiruvalluvar University College of Arts and Science, Thennangur - 604 408, Tamil Nadu, India
**Corresponding author E-mail: srimanjushc@gmail.com*

Abstract

Authorsipresent sufficienticonditions for theioscillation of the generalizediperturbed quasilinearidifferenceiequation $\Delta_\ell\left(a((x-1)\ell+i)\left|\Delta_\ell v((x-1)\ell+i)\right|^{\gamma-1}\Delta_\ell v((x-1)\ell+i)\right)+F(x\ell+i,v(x\ell+i))=G(x\ell+i,v(x\ell+i),\Delta_\ell v(x\ell+i))$ where $0<\gamma<1$, $x\in[0,\infty)$ and $i=x-\left[\frac{x}{\ell}\right]\ell$. Examplesiillustrates the importanceiof our results are alsoiincluded.

Keywords: *Generalizedidifference equation; Oscillation; iQuasilinear,*

1. Introduction

Difference equations represent captivating mathematical field, has rich field of the applications in such diverse disciplines as population dynamics, operations research, ecology, economics, biology etc. For thelbackgroundlof differencelequations and its applicationslin diverselfields withlexamples, see [1]. The study of difference equations is based on the operator Δ defined as $\Delta u(x)=u(x+1)-u(x), x\in[0,\infty)$.

Thoughlmany authors [1],[16] have discussed the definition of Δ as

$$\Delta u(x)=u(x+\ell)-u(x),\quad \ell\in(0,\infty), \tag{1}$$

no notable progress have been taken on this line. Butlin [13] the authors took up the definition of Δ as given in (1), and given many important results and applications. They labeled the operator Δ defined by (1) as Δ_ℓ and its inversely Δ_ℓ^{-1}, many interesting results in number theory were obtained. Qualitativelproperties like rotatory, expanding, lshrinking, spiral and web like were established by extending theory ofl Δ_ℓ to complex function, for the solutions of difference equations involving Δ_ℓ in [2-15, 17-21].

In the sequel, in this paper we will be considered the generalized perturbed quasi linear difference equation for $x\in[0,\infty)$

$$\Delta_\ell\left(a((x-1)\ell+i)\left|\Delta_\ell v((x-1)\ell+i)\right|^{\gamma-1}\Delta_\ell v((x-1)\ell+i)\right)$$

$$+F(x\ell+i,v(x\ell+i))=G(x\ell+i,v(x\ell+i),\Delta_\ell v(x\ell+i)) \tag{2}$$

where $0<\gamma<1$, $a(x\ell+i)$ is an eventually positive real valued function, land Δ_ℓ is the generalized forward difference operator defined as $\Delta_\ell v(x\ell+i)=v((x+1)\ell+i)-v(x\ell+i)$.

By alsolution of (2), we mean a nontrivial real valued function $v(x\ell+i)$ satisfying (2) for $x\in[0,\infty)$. A solution $v(x\ell+i)$ is said to be oscillatory if it is neither eventually positive nor negative, and non oscillatory otherwise.

2. Main Results

In this paper we assume that there exist real valued functions $q(x\ell+i)$, $p(x\ell+i)$ andla function $f:R\to R$ such that

(i). $vf(v)>0$ for all $v\neq 0$;

(ii). $f(v)-f(w)=g(v,w)(v-w)$ for $v,w\neq 0$, where g is a nonnegativelfunction; land

(iii). $\frac{F(x\ell+i,v\ell+i)}{f(v\ell+i)}\geq q(x\ell+i)$,

$\frac{G(x\ell+i,v\ell+j,w\ell+i)}{f(v\ell+i)}\leq p(x\ell+i)$ for $v,w\neq 0$.

Thelfollowing conditionslare usedlthroughout thislpaper:

$$\sum\frac{1}{a^{1/\gamma}((x-1)\ell+i)}=\infty, \tag{3}$$

$$\int_\theta^\infty\frac{dx}{f(x)^{1/\gamma}}<\infty,\ \int_{-\theta}^{-\infty}\frac{dx}{f(x)^{1/\gamma}}<\infty \text{ forall } \theta>0 \tag{4}$$

$$\liminf_{x\to\infty}\sum_{r=x_0}^{x}(q(r\ell+i)-p(r\ell+i))\geq 0 \text{ for all large } x_0, \tag{5}$$

$$\int_0^\theta\frac{dx}{f(x)^{1/\gamma}}<\infty,\ \int_0^{-\theta}\frac{dx}{f(x)^{1/\gamma}}<\infty \text{ for all } \theta>0, \tag{6}$$

Copyright © 2018 Authors. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

$$\sum\left[\frac{N}{a(r\ell+i)}-\frac{1}{a(r\ell+i)}\sum_{s=x_0}^{r}(q(s\ell+i)-p(s\ell+i))\right]=-\infty \qquad (7)$$

forlevery constantl*N*,

$$\limsup_{x\to\infty}\sum_{r=x_0}^{x}(q(r\ell+i)-p(r\ell+i))=\infty \text{ for allllarge } x_0, \qquad (8)$$

$$\limsup_{x\to\infty}\sum_{r=x_0}^{x}r(q(r\ell+i)-p(r\ell+i))=\infty \text{ forlall large } x_0, \qquad (9)$$

$$\sum(q(x\ell+i)-p(x\ell+i))R(x\ell+i,x_0\ell+i)=\infty \text{ where}$$

$$R(x\ell+i,x_0\ell+i)=\sum_{r=x_0}^{x}\frac{1}{a((r-1)\ell+i)}, \qquad (10)$$

$$\sum\frac{1}{(a(n-1)\ell+i)}<\infty, \qquad (11)$$

$$\sum(q(x\ell+i)-p(x\ell+i))T(x\ell+i,x_0\ell+i)=\infty \text{ where} \qquad (12)$$

$$T(x\ell+i,x_0\ell+i)=R((x-1)\ell+i,x_0\ell+i)=\sum_{r=x_0}^{x-1}\frac{1}{a((r-1)\ell+i)},$$

$$\sum\frac{1}{a((x-1)\ell+i)}=\infty, \qquad (13)$$

$$\frac{a(x\ell+i)}{a((x-1)\ell+i)}\le 1 \text{ for } x\ge 1. \qquad (14)$$

Theorem 1 *Supposal* $\gamma\ge 1$ *and (5)-(7) hold. Then, all solutions of (2) are oscillatory.*

Proof. Suppose that $v(x\ell+i)$ is alnonoscillatory solution of (2), say, $v(x\ell+i)>0$ for $k\ge k_0\ge 1$. Since (5) holds, $\Delta_\ell v(x\ell+i)$ does not oscillate. Welbegin withlthe followinglidentity

$$\Delta_\ell\left[\frac{a((x-1)\ell+i)\,|\Delta_\ell v((x-1)\ell+i)|^{\gamma-1}\Delta_\ell v((x-1)\ell+i)}{f(v((x-1)\ell+i))}\right]$$
$$=\frac{G(x\ell+i,v(x\ell+i),\Delta f(v(x\ell+i)))}{f(v(x\ell+i))}-\frac{F(x\ell+i,v(x\ell+i))}{f(v(x\ell+i))}$$
$$-\frac{a((x-1)\ell+i)g(v(x\ell+i),v((x-1)\ell+i))(\Delta_\ell v((x-1)\ell+i))^2}{f(v((x-1)\ell+i))\,f(v(x\ell+i))}$$
$$\times|\Delta_\ell v((x-1)\ell+i)|^{\gamma-1} \qquad (15)$$

whichlimplies

$$\Delta_\ell\left[\frac{a((x-1)\ell+i)\,|\Delta_\ell v((x-1)\ell+i)|^{\gamma-1}\Delta_\ell v((x-1)\ell+i)}{f(v((x-1)\ell+i))}\right]$$
$$\le p(x\ell+i)-q(x\ell+i). \qquad (16)$$

Case 1. Suppose that $\Delta_\ell v(x\ell+i)\ge 0$ for $x\ge x_1\ge x_0$. Summing (16) from (x_1+1) to x gives

$$\frac{a(x\ell+i)|\Delta_\ell v(x\ell+i)|^{\gamma-1}\Delta_\ell v(x\ell+i)}{f(v(x\ell+i))}$$
$$\le\frac{a(x_1\ell+i)|\Delta_\ell v(x_1\ell+i)|^{\gamma-1}\Delta_\ell v(x_1\ell+i)}{f(v(x_1\ell+i))}$$
$$-\sum_{r=x_1+1}^{x}(q(r\ell+i)-p(r\ell+i))$$

$$\frac{|\Delta_\ell v(x\ell+i)|^{\gamma-1}\Delta_\ell v(x\ell+i)}{f(v(x\ell+i))}\le\frac{N}{a(x\ell+i)}$$

$$-\frac{1}{a(x\ell+i)}\sum_{r=x_1+1}^{x}(q(r\ell+i)-p(r\ell+i)) \qquad (17)$$

where $X=a(x_1\ell+i)\,|\Delta_\ell v(x_1\ell+i)|^{\gamma-1}\Delta_\ell v(x_1\ell+i)/f(v(x_1\ell+i))$. Again we sum (17) from (k_1+1) to k, to obtain

$$\sum_{r=x_1+1}^{x}\frac{|\Delta_\ell v(r\ell+i)|^{\gamma-1}\Delta_\ell v(r\ell+i)}{f(v(r\ell+i))} \qquad (18)$$
$$\le\sum_{s=x_1+1}^{x}\left[\frac{N}{a(r\ell+i)}-\frac{1}{a(r\ell+i)}\sum_{s=x_1+1}^{r}(q(s\ell+i)-p(s\ell+i))\right].$$

By (7), the right side of (18) tends to $-\infty$ as $x\to\infty$ where as the left side is nonnegative.

Case 2. Suppose that $\Delta_\ell v(x\ell+i)<0$ for $x\ge x_1\ge x_0$. Then, lfrom (18) welfind

$$-\sum_{r=x_1+1}^{x}\left[\frac{N}{a(r\ell+i)}-\frac{1}{a(r\ell+i)}\sum_{s=x_1+1}^{r}(q(s\ell+i)-p(s\ell+i))\right]$$
$$\le\sum_{r=x_1+1}^{x}\frac{|\Delta_\ell v(r\ell+i)|^{\gamma}}{f(v(r\ell+i))} \qquad (19)$$

$$\le\left[\sum_{r=x_1+1}^{x}\frac{|\Delta_\ell v(r\ell+i)|}{f(v(r\ell+i))^{1/\gamma}}\right]^{\gamma}$$
$$\le\left[\int_{v(x+1)}^{v(x_1+1)}\frac{du}{f(u)^{1/\gamma}}\right]^{\gamma}$$
$$\le\left[\int_{0}^{v(x_1+1)}\frac{du}{f(u)^{1/\gamma}}\right]^{\gamma}. \qquad (20)$$

By (7), the left side of (20) tends to ∞ asl $x\to\infty$ where as the right side is finite by (6).

Theorem 2 *Suppose* $a((x-1)\ell+i)\equiv 1$, $\gamma\ge 1$ l*and (4), (5), (9) hold. Then all solutions of (2) are oscillatory.*

Proof. Assume that $v(x\ell+i)$ is a nonsocial atory solution of (2), say, $v(x\ell+i)>0$ for $x\ge x_0\ge 1$. Since (5) holds, we see that $\Delta_\ell v(x\ell+i)$ does not oscillate.

Welbegin withlthe followinglidentity

$$\Delta_\ell\left[\frac{(x\ell+i)\,|\Delta_\ell v((x-1)\ell+i)|^{\gamma-1}\Delta_\ell v((x-1)\ell+i)}{f(v(x\ell+i))}\right]$$
$$=\frac{(x\ell+i)G(x\ell+i,v(x\ell+i),\Delta f(v(x\ell+i)))}{f(v(x\ell+i))}$$
$$-\frac{(x\ell+i)F(x\ell+i,v(x\ell+i))}{f(v(x\ell+i))}+\frac{|\Delta_\ell v(x\ell+i)|^{\gamma-1}\Delta_\ell v(x\ell+i)}{f(v((x+1)\ell+i))}$$
$$-\frac{(x\ell+i)g(v((x+1)\ell+i),v(x\ell+i))(\Delta_\ell v(x\ell+i))^2}{f(v(x\ell+i))}$$
$$\times\frac{|\Delta_\ell v(x\ell+i)|^{\gamma-1}}{f(v((x+1)\ell+i))}$$

Which give rise to

$$\Delta_\ell\left[\frac{(x\ell+i)\,|\Delta_\ell v((x-1)\ell+i)|^{\gamma-1}\Delta_\ell v((x-1)\ell+i)}{f(v(x\ell+i))}\right]$$

$$\le (x\ell+i)(p(x\ell+i)-q(x\ell+i))+\frac{|\Delta_\ell v(x\ell+i)|^{\gamma-1}\Delta_\ell v(x\ell+i)}{f(v((x+1)\ell+i))}. \quad (21)$$

Case 1. Suppose that $\Delta_\ell v(x\ell+i)\ge 0$ for $x\ge x_1\ge x_0$. Summing (21) from (x_1+1) to x gives

$$\sum_{r=x_1+1}^{x}(r\ell+i)(q(r\ell+i)-p(r\ell+i)) \le \frac{((x_1+1)\ell+i)(\Delta_\ell v(x_1\ell+i))^\gamma}{f(v((x_1+1)\ell+i))}$$
$$-\frac{((x+1)\ell+i)(\Delta_\ell v(x\ell+i))^\gamma}{f(v((x+1)\ell+i))}+\sum_{r=x_1+1}^{x}\frac{(\Delta_\ell v(r\ell+i))^\gamma}{f(v((r+1)\ell+i))}$$
$$\le \frac{((x_1+1)\ell+i)(\Delta_\ell v(x_1\ell+i))^\gamma}{f(v((x_1+1)\ell+i))}+\sum_{r=x_1+1}^{x}\frac{(\Delta_\ell v(r_1\ell+i))^\gamma}{f(v((r+1)\ell+i))}$$
$$\le \frac{((x_1+1)\ell+i)(\Delta_\ell v(x_1\ell+i))^\gamma}{f(v((x_1+1)\ell+i))}+\left[\sum_{r=x_1+1}^{x}\frac{\Delta_\ell v(r\ell+i)}{f(v((r+1)\ell+i))^{1/\gamma}}\right]^\gamma$$
$$\le \frac{((x_1+1)\ell+i)(\Delta_\ell v(x_1\ell+i))^\gamma}{f(v((x_1+1)\ell+i))}+\left[\int_{v(x_1+1)}^{v(x+1)}\frac{du}{f(u)^{1/\gamma}}\right]^\gamma. \quad (22)$$

By (9), the left side of (22) tends to ∞ as $x\to\infty$ whereas the right side is finite by (4).

Case 2. Suppose that $\Delta_\ell v(x\ell+i)<0$ for $k\ge k_1\ge k_0$. Condition (21) implies the existence of an integer $x_2\ge x_1$ such that

$$\sum_{r=x_1+1}^{x}(r\ell+i)(q(r\ell+i)-p(r\ell+i))\ge 0,\ x\ge x_2+1. \quad (23)$$

Multiplying l (2) by $x\ell+i$ land using l (iii), welobtain

$$(x\ell+i)(|\Delta_\ell v((x-1)\ell+i)|^{\gamma-1}\Delta_\ell v((x-1)\ell+i))$$
$$\le (x\ell+i)f(v(x\ell+i))(p(x\ell+i)-q(x\ell+i))$$

Which on summing by parts from (x_2+1) to x provides

$$((x+1)\ell+i)|\Delta_\ell v(x\ell+i)|^{\gamma-1}\Delta_\ell v(x\ell+i)$$
$$\le ((x_2+1)\ell+i)|\Delta_\ell v(x_2\ell+i)|^{\gamma-1}\Delta_\ell v(x_2\ell+i)$$
$$+\sum_{r=x_1+1}^{x}|\Delta_\ell v(r\ell+i)|^{\gamma-1}\Delta_\ell v(r\ell+i)$$
$$-\sum_{r=x_1+1}^{x}(r\ell+i)(q(r\ell+i)-p(r\ell+i))$$
$$=((x_2+1)\ell+i)|\Delta_\ell v(x_2\ell+i)|^{\gamma-1}\Delta_\ell v(x_2\ell+i)$$
$$+\sum_{r=x_1+1}^{x}|\Delta_\ell v(r\ell+i)|^{\gamma-1}\Delta_\ell v(r\ell+i)-f(v((x+1)\ell+i))$$
$$\times\sum_{r=x_1+1}^{x}(r\ell+i)(q(r\ell+i)-p(r\ell+i))$$
$$+\sum_{r=x_1+1}^{x}\Delta_\ell f(v(r\ell+i))\left[\sum_{s=x_1+1}^{r}(s\ell+i)(q(s\ell+i)-p(s\ell+i))\right]$$
$$=((x_2+1)\ell+i)|\Delta_\ell v(x_2\ell+i)|^{\gamma-1}\Delta_\ell v(x_2\ell+i)$$
$$+\sum_{r=x_1+1}^{x}|\Delta_\ell v(r\ell+i)|^{\gamma-1}\Delta_\ell v(r\ell+i)$$
$$-f(v((x+1)\ell+i))\sum_{r=x_1+1}^{x}(r\ell+i)(q(r\ell+i)-p(r\ell+i))$$
$$+\sum_{r=x_1+1}^{x}g(v((r+1)\ell+i),v(r\ell+i))\Delta_\ell v(r\ell+i)$$
$$\times\left[\sum_{s=x_1+1}^{r}(s\ell+i)(q(s\ell+i)-p(s\ell+i))\right]$$
$$\le ((x_2+1)\ell+i)|\Delta_\ell v(x_2\ell+i)|^{\gamma-1}\Delta_\ell v(x_2\ell+i)$$

Where we have also used (23) in the last inequality. It followslthat

$$\Delta_\ell v(x\ell+i)\le\frac{-((x_2+1)\ell+i)^{1/\gamma}|\Delta_\ell v(x_2\ell+i)|}{((x+1)\ell+i)^{1/\gamma}}, \quad (24)$$

For $x\ge x_2+1$. Once again we sum (24) from (x_2+1) to x to get

$$v((x+1)\ell+i)\le v((x_2+1)\ell+i)$$
$$-((x_2+1)\ell+i)^{1/\gamma}|\Delta_\ell v(x_2\ell+i)|\sum_{r=x_2+1}^{x}\frac{1}{((r+1)\ell+i)^{1/\gamma}}. \quad (25)$$

The right side of (25) tends to $-\infty$ as $x\to\infty$, this contradicts the assumption that $v(x\ell+i)$ is eventually positive.

References

[1] Agarwal RP, *Difference Equations and Inequalities*, Marcel Dekker, New York (2000).

[2] Benevatho Jaison A, Khadar Babu Sk (2016), Oscillation for generalized first order nonlinear difference equations. *Global Journal of Pure and Applied Mathematics* 12(1), 51-54.

[3] Benevatho Jaison A, Khadar Babu Sk (2016), Kamenev-type oscillation criteria for second order generalized delay difference equations. *International Journal of Control Theory and Applications* 9(28), 463-469.

[4] Benevatho Jaison A, Khadar Babu Sk (2016), Oscillation for generalized first order nonlinear a-difference equations. *International Journal of Pure and Applied Mathematics* 109(7), 67-74.

[5] Benevatho Jaison A, Khadar Babu Sk (2017), Oscillation theorems for generalized second-order nonlinear delay difference equations. *International Journal of Pure and Applied Mathematics* 113(9), 84-92.

[6] Benevatho Jaison A, Khadar Babu Sk (2017), Oscillation theorems for generalized second kind nonlinear delay difference equations. *International Journal of Pure and Applied Mathematics* 115(9), 25-36.

[7] Benevatho Jaison A, Khadar Babu Sk (2017), Oscillatory behavior of generalized nonlinear difference equations. *Global Journal of Pure and Applied Mathematics* 13(1), 205-209.

[8] Benevatho Jaison A, Khadar Babu Sk (2017), Oscillatory behavior of generalized Nonlinear difference equations. *Global Journal of Pure and Applied Mathematics* 13(2), 415-423.

[9] Benevatho Jaison A, Khadar Babu Sk (2018), Kamenev-type oscillation criteria for generalized sublinear delay difference equations. *International Journal of Pure and Applied Mathematics* 118(10), 135-145.

[10] Benevatho Jaison A, Khadar Babu Sk (2018), Oscillation for generalized second kind nonlinear delay a-difference equations. *International Journal of Pure and Applied Mathematics* 118(23), 507-515.

[11] Chandrasekar V, Srimanju V (2016), Oscillation for generalized second order sublinear neutral delay alpha difference equations. *Global Journal of Pure and Applied Mathematics* 12(1), 55-59.

[12] Chandrasekar V, Srimanju V (2016), Qualitative properties of discrete version of generalized kneser's and arzela-ascoli's theorems. *International Journal of Control Theory and Applications* 9(28), 549-554.

[13] Maria Susai Manuel M, Britto Antony Xavier G, Thandapani E (2006), Theory of generalized difference operator and its applications. *Far East Journal of Mathematical Sciences* 20(2), 163-171.

[14] Maria Susai Manuel M, Chandrasekar V, Britto Antony Xavier G (2007), Generalized bernoulli polynomials through weighted pochhammer symbols. *Journal of Modern Methods in Numerical Mathematics* 26(3), 321-333.

[15] Maria Susai Manuel M, George Maria Selvam A, Britto Antony Xavier (2006), Rotatory and boundedness of solutions of certain class of difference equations. *International Journal of Pure and Applied Mathematics* 33(3), 333-343.

[16] Ronald E. Mickens, *Difference Equations*, Van Nostrand Reinhold Company, New York (1990).

[17] Srimanju V, Khadar Babu Sk (2017), Oscillatory criteria for generalized second-order quasilinear neutral delay difference equations. *International Journal of Pure and Applied Mathematics* 113(9), 75-83.
[18] Srimanju V, Khadar Babu Sk (2017), Oscillation of generalized quasilinear difference equations. *International Journal of Pure and Applied Mathematics* 115(9), 37-45.
[19] Srimanju V, Khadar Babu Sk (2017), Oscillation criteria for generalized quasi-linear difference equations. *Global Journal of Pure and Applied Mathematics* 13(1), 210-216.
[20] Srimanju V, Khadar Babu Sk (2017), Oscillation criteria for generalized second kind quasi-linear neutral a-difference equations. *Global Journal of Pure and Applied Mathematics* 13(2), 544-551.
[21] Srimanju V, Khadar Babu Sk (2018), Oscillatory properties of third-order quasilinear generalized difference equations. *International Journal of Pure and Applied Mathematics* 118(10), 155-165.
[22] Srimanju V, Khadar Babu Sk (2018), Oscillation of generalized quasilinear a-difference equations. *International Journal of Pure and Applied Mathematics* 118(23), 497-505.