International Journal of Engineering & Technology, 7 (4.10) (2018) 1001-1006

International Journal of Engineering & Technology

Website: www.sciencepubco.com/index.php/IJET

Research paper

G, $\overline{G}$ Properties – from $\overline{G}$, G

A. Elakkiya[1], M. Yamuna[2]*

[1]*Vellore Institute of Technology, Vellore*
[2] *Vellore Institute of Technology, Vellore*
**Corresponding author E-mail: myamuna@vit.ac.in*

Abstract

In this paper weprovide a method of determining the chromatic polynomial of $\overline{G}$ without actual construction of $\overline{G}$. A planar graph characterization of graphs whose domatic partition is $\frac{n}{\gamma(G)}$ using $\overline{G}$ properties is established and provide a MATLAB program for identifying just excellent graphs.

Keywords:*Chromatic Polynomial; Complement Graph; Just Excellent; Planar.*

1. Introduction

In [1] M. Yamuna et.al have determined the chromatic polynomial of G^* without actual construction In [2], M. Yamuna et al has introduced just excellent graph . M. Yamuna et.al have proved that just excellent graph is not a domatic subdivision stable graph[3]. In [4], [5],M. Yamuna et al introduced Non domination subdivision stable graphs (NDSS) and characterized planarity of complement of NDSS graphs.. In [6],[7], M. Yamuna et al introduced γ - uniquely colorable graphs and also provided the constructive characterization of γ - uniquely colorable trees and characterized planarity of complement of γ - uniquely colorable graphs.This paper targets to determine properties of $\overline{G}$using properties of G without constructing $\overline{G}$.

2. Terminology

We consider only simple graph with n and m vertices and edges. The graph H is said to be the complement of G if any two vertices arenotadjacent in G, then in H they are adjacent.G**.** uvis obtained by contracting (u, v), where u ⊥v. Results related to graph theory we refer to [9].
D is a dominating set if every vertex of V – D is adjacent to some vertex of D. Minimum cardinality of D, is said to be a minimum dominating set (MDS). The cardinality of any MDS for G is said to be domination number of G, represented by $\gamma(G)$. Results related to domination we refer to [10].
We know that in a complete graph, the maximum number of edges is $(n\,(n-1))/2$. Also $E(G) \cup E(\overline{G}) = (n\,(n-1))/2$. If n = 100, then $G \cup \overline{G}$ has 4950 edges. If G has say thousand edges $\overline{G}$ has 3950 edges. It is obvious that construction $\overline{G}$ is very difficult in such cases. As n increases, this complexity invariably increases. When properties of $\overline{G}$ are required in such cases it is better to determine them from G without drawing $\overline{G}$.

3. Result and discussion

This section gives a recursive technique for finding the chromatic polynomial of $\overline{G}$ from G **.**
This results and discussions is for modifying the existing technique of determining the chromatic polynomial ($P_n(\lambda)$) for any graph, so that $P_n(\lambda)$ of $\overline{G}$ can be determined from G. Throughout this section for any edge e = (u, v) in G the corresponding edge in $\overline{G}$ is denoted by $\overline{e}$ = ($\overline{u}$, $\overline{v}$).

3.1. Modified G – { e }

We know that no edges in G implies an edge in $\overline{G}$. We use this to modify the operation G – { e }. Comparing G and$\overline{G}$, picking two non – adjacent vertices in $\overline{G}$ is equivalent to choosing adjacent vertices in G.
For example consider the graph G in Fig.1 and its complement $\overline{G}$ **.**




Fig.1

1 not adjacent to 2 in $\overline{G}$. Add an edge between 1 and 2 in $\overline{G}$ to generate graph G_1, that is $G_1 = \overline{G} \cup \{ (1\,2) \}$.
We aim to generate the graph G_1 from graph G. Since it is just an additional edge available in G_1, removal of this edge in G would satisfy our requirement. Also we observe that $\overline{G - \{ (1\,2) \}} = G_1$. So we conclude that adding an edge between non adjacent vertices in $\overline{G}$ is equivalent to removing the edge between same vertex pair in G.

Copyright © 2018 Authors. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

3.2. Type – I Operation

Remove edge e = (u, v) in G. Let $G_1 = G - \{ e \}$. By applying Type – I Operation we create a new graph G_1,so that $\overline{G}_1 = \overline{G} + \{\overline{e}\}$.

3.3. Modified G.uv




Fig.2

Merging vertices 1 and 2 of $\overline{G}$, we obtain graph $\overline{G}$.12. We observe that vertices1and2 are together not adjacent to the vertices 3, 5 respectively in $\overline{G}$. The same is true in $\overline{G}$.12 also.

Vertices 4, 6 are adjacent to atleast one of vertex 1 or 2. So in $\overline{G}$.12 vertex 12 is adjacent to the vertex 4 and 6. We have to decide a suitable operation in G which would generate a graph G_2, so that complement of G_2 is G.uv for any two non adjacent vertices u, v in $\overline{G}$. From the above observation, we can decide that for any two adjacent vertices e = (u, v) in G generate a graph G.uv as follows

1.If u and v are collectively ⊥to vertices $v_1, v_2, \ldots v_p$, then in G draw an edge between vertex pairs (uv, v_1), (uv, v_2), … (uv, v_p) respectively.

2.If $u_1, u_2, \ldots u_q$are vertices adjacent to atleast one of u or v then there is no edge between vertex pairs (uv, u_1), (uv, u_2), … (uv, u_q) respectively.

By applying these two operations, we observe that in the complement of $\overline{G}$.uv there exist no edge between uv and vertices $v_1, v_2, \ldots v_p$. Also there exist edges between uv and vertices $u_1, u_2, \ldots u_q$.

3.4. Type – II Operation

Let e = (u, v). Let $X = \{ v_1, v_2, \ldots v_p \} \subseteq V (G)$ such that u and v are both adjacent to $v_1, v_2, \ldots v_p$. Let $Y = \{ u_1, u_2, \ldots u_q \} \subseteq V (G)$ such that $u_1, u_2, \ldots u_q$ are adjacent to atleast one of u or v.

Generate G.uv by applying the following operations

1.Draw edge between (uv, v_1), (uv, v_2), … (uv, v_p) .


2.There is no edge between (uv, u_1), (uv, u_2), … (uv, u_q).

3.5. Complete Graph Identification


By recursive applying Type I, II operations on G, we reach a graph H such that H is a null graph. This means that $\overline{H}$ is a complete graph. Let G be any graph. Apply modified G – { e } and G. uv on H to generate G_1, G_2 , that is G_1 is the graph obtained by applying Type I operation on G and G_2 is the graph obtained by applying Type II operation on G. Continue this procedure recursively on G_1 and G_2 and on the remaining graphs generated until we cannot continue any further. This meanst hat we now have a sequence of graphs $S_1, S_2, \ldots, S_k$ such that each S_i , i = 1, 2, …, k are null graphs, so that $\overline{S}_i$, i = 1, 2, …, k is a complete graph.

- 1). Chromatic polynomial of $\overline{S}_i = \prod_{k=0}^{n-1}(\lambda - k)$, if S_i is a null graph with n vertices. So $P_n(\lambda)(S_i) = \sum_{i=1}^{k} P_n(\lambda)(\overline{S_i})$. This procedure is summarized in hierarchy.

In Fig. 3 $P_n(\lambda)(\overline{S_i}) = \lambda(\lambda - 1)(\lambda - 2)(\lambda - 3)(\lambda - 4) + \lambda(\lambda - 1)(\lambda - 2)(\lambda - 3) + \lambda(\lambda - 1)(\lambda - 2)(\lambda - 3) + \lambda(\lambda - 1)(\lambda - 2)(\lambda - 3) + \lambda(\lambda - 1)(\lambda - 2) + \lambda(\lambda - 1)(\lambda - 2)(\lambda - 3) + \lambda(\lambda - 1)(\lambda - 2) + \lambda(\lambda - 1)(\lambda - 2) + \lambda(\lambda$



Hierarchy. 1.






Fig.3

From modified G – e and G.uv , we understand that for every pair of adjacent vertices (non – adjacent in $\overline{G}$)

- $(\overline{G - e}) = \overline{G} + \bar{e}$ and
- $(\overline{G \bullet uv}) = \overline{G}.uv$

that is the recursive process is determined in such a way that at every stage the number of non – adjacent vertices in G =the number of adjacent vertices in $\overline{G}$. The numberof operations involved in determining the chromatic polynomial of $\overline{G}$ fromG is equal to the number of operations involved in determining the same from $\overline{G}$, since the operations are repeated for every adjacent pair in G

4. Planarity of $\overline{G}$ - from G

Planar graphs are often characterized by the following famous theorems that uses Kuratowski's graphs.

R1. G is planar iff G does not contain K_5 or $K_{3,3}$as a subgraph or any graph homeomorphic to K_5 or $K_{3,3}$ as a subgraph.

R2. G is planar iff it does not have a subgraph contractible to K_5 or $K_{3,3}$ [6].

R_3. G is outer planar iff it has no subgraph homeomorphic to K_4 or $K_{2,3}$except $K_4 - x$ [6].

In this section, we shall prove that a just excellent graph is non – planar, when $\gamma(G) \geq 4$ using R_1 and R_2. If $\gamma(G) = 1$, $\overline{G}$ is disconnected, so we consider the cases only when $1 < \gamma(G) \leq k$, for some integer k.

Theorem 1

Let G be a just excellent graph. Let $D = \{u_1, u_2, \ldots u_m\}$,$m \geq 3$ be a γ - set for G. Let $X_1 = pn(u_1, D) = \{a_1, a_2, \ldots, a_{k1}\}$, $X_2 = pn(u_2, D) = \{b_1, b_2, \ldots, b_{k2}\}$, . . . , $X_m = pn(u_m, D) = \{m_1, m_2, \ldots, mk_m\}$. Then

A.$\langle pn(u_i, D)\rangle$ is not complete, for all i = 1, 2, . . . , m.

B.$\langle X_i, X_j\rangle$ is not complete, $\forall\ i \neq j$.

C. Assume that condition B is satisfied. Then for some $z_k \in X_k$, it is not possible that (y_j, z_k) is the only non – adjacent vertex pair in (X_j, X_k) such that $y_j \in X_j$, $z_k \in X_k$.

Proof

A.If possible assume that $\langle pn(u_1, D)\rangle$ is complete. Then $D' = D - \{u_1\} \cup \{a_i\}$ is a γ - set for G. D, D′ are γ - sets for G containing $\{u_2, u_3, \ldots u_m\}$, a contradiction, $\Rightarrow \langle pn(u_i, D)\rangle$ is not complete, for all i = 1, 2, . . . , m.

B. If $\exists$ some X_i, X_j, $i \neq j$, $\ni \langle X_i, X_j\rangle$ is complete, then $D' = D - \{u_i\} - \{u_j\} \cup \{x_i\} \cup \{y_j\}$, where $x_i \in pn(u_i, D)$, $y_j \in pn(u_j, D)$ is a γ - set for G. D, D′ are two distinct γ - sets for G containing $\{u_1, u_2, \ldots u_{i-1}, u_{i+1}, \ldots, u_{j-1}, u_{j+1}, \ldots u_m\}$, a contradiction to our assumption that G is just excellent, $\Rightarrow \langle X_i, X_j\rangle$ is not complete, $\forall\ i \neq j$.

C.By part B, we know that there exist some $x_i \in pn(u_i, D)$, $y_j \in pn(u_j, D)$ such that x_i not adjacent to y_j. If possible, assume that in (X_j, X_k) , $y_j$not adjacent to $z_k$is the only non – adjacent vertex pair that is, $\forall$vertex in $X_j - \{y_j\}$ is $\perp$ to every vertex in X_k. $D' = D - \{u_j\} - \{u_k\} \cup \{y_a\} \cup \{z_b\}$, $y_a \in X_j$, $z_b \in X_k$, $a \neq j$, $b \neq k$ are γ - sets for G that is, D, D′ are two distinct γ - sets for G containing u_i, a contradiction. Hence statement C is true

Theorem 2

Let G be a just excellent graph, $\ni \gamma(G) = 3$. Then $\overline{G}$ is non – outer planar.

Proof

Let $\gamma(G) = 3 = \{u_1, u_2, u_3\}$. Let $X_1 = pn(u_1, D) = \{a_1, a_2, \ldots, a_{k1}\}$, $X_2 = pn(u_2, D) = \{b_1, b_2, \ldots, b_{k2}\}$, $X_3 = pn(u_3, D) = \{c_1, c_2, \ldots, c_{k3}\}$. We prove that $\overline{G}$ is non – planar by showing that $\overline{G}$contains $K_{2,3}$ as a subgraph. Let V_1 and V_2 be the vertex set of $K_{2,3}$in$\overline{G}$. Choose $V_2 = \{u_1, u_2, u_3\}$.




Fig.4

By part B of Theorem 1, we know that there exist
i. a_i, b_j such that a_i not adjacent to b_j,
ii b_k, c_l such that b_k not adjacent to c_l, $k \neq j$,
This is possible, because pn $(u_i, D) \geq 2$, for all $u_i \in D$. a_i not adjacent to b_j, b_k not adjacent to c_l, in $\overline{G}$. Contract these edges in $\overline{G}$. Let $V_1 = \{ a_ib_j, b_kc_l \}$. $a_i \in X_1$, $b_j \in X_2$, a_i not adjacent to $\{u_2, u_3\}$ and b_j not adjacent to u_1 in G. This means that a_i adjacent to $\{u_2, u_3\}$ and b_j adjacent to u_1 in $\overline{G}$. That is vertex $a_ib_j \perp$ to $\{ u_1, u_2, u_3 \}$ in $\overline{G}$. Similarly vertex $b_kc_l \perp$ to $\{ u_1, u_2, u_3 \}$ in $\overline{G}$. $\langle u_1, u_2, u_3, a_ib_j, b_kc_l \rangle$ is $K_{2,3}$ in $\overline{G}$, that is $\overline{G}$ is non – outer planar.




Fig.5

Theorem 3
Let Gbe a just excellentgraph, ∋ γ (G) = 4. Then$\overline{G}$ is non – planar.

Proof
Let γ (G) = 4 = D = $\{u_1, u_2, u_3\}$. Let X_1 = pn $(u_1, D) = \{a_1, a_2, \ldots, a_{k1}\}$, X_2 = pn $(u_2, D) = \{b_1, b_2, \ldots, b_{k2}\}$, X_3 = pn $(u_3, D) = \{c_1, c_2, \ldots, c_{k3}\}$, X_4 = pn $(u_4, D) = \{d_1, d_2, \ldots, d_{k4}\}$. We prove that $\overline{G}$ is non – planar by showing that $\overline{G}$ contains $K_{3,3}$ as a subgraph. Let V_1 and V_2 be the vertex set of $K_{3,3}$ in $\overline{G}$.
Choose some $b_i \in X_2$, $a_i \in X_1$. b_i not adjacent to u_1, a_i not adjacent to u_4 in G, implies u_1 adjacent to b_i, u_4 adjacent a_i in $\overline{G}$. Choose $V_1 = \{u_1b_i, u_2, u_3\}$, $V_2 = \{u_4a_i, d_i, d_j\}$ where $d_i, d_j \in X_4$.




Fig.6

i. u_1 not adjacent to d_i, d_j and u_4 not adjacent to b_i.
ii. u_2 not adjacent to d_i, d_j, a_i.
iii. u_3 not adjacent to d_i, d_j, a_i in G.




Fig.7

This implies, these vertices are adjacent in $\overline{G}$. Hence $\langle V_1, V_2 \rangle$ is $K_{3,3}$ implies $\overline{G}$ is non – planar.

Note
- Generalizing Theorem 3, we conclude that, if G is just excellent graph ∋ γ (G) ≥ 4, then $\overline{G}$ is non – planar.
- When γ (G) = 2, G just excellent, $\overline{G}$may be planar or non – planar.

Example
In Figure 8 G,$\overline{G}$ are planar, G is just excellent.




Fig.8

Figure 9 G is just excellent, $\overline{G}$ - {(6, 4)}, {(7, 4)} is a subgraph of$\overline{G}$




Fig. 9

Contracting (4, 5), (6, 3), (7, 2), we see that ⟨1, 63, 45, 72, 8⟩ is K_5 and hence $\overline{G}$is non planar.
If G is just excellent graph ∋

i. γ (G) = 2, then $\overline{G}$ may be planar or non – planar.
ii. γ (G) = 3, then $\overline{G}$ is non – outer planar.
iii. γ (G) ≥ 4, then $\overline{G}$ is non – planar.

5. Matrix Representation for Just Excellent Graphs

Let G be any graph with n – vertices. Let A represents the adjacency matrix of G. Let N denote a n ×n matrix, where

$$N = [n_{ij}]_{n \times n} = \begin{cases} 1 & \text{if } i = j \\ a_{ij} & \text{the } (i,j)^{th} \text{ entry in the adjacency matrix} \end{cases}$$

Let $x = \langle x(v_1), x(v_2), \ldots, x(v_n) \rangle^T$ be a { 0, 1 } vectors . We

know that, if x represents the any dominating set, then Nx≥ 1, that is in a resulting matrix Nx, all the entry values are non zero.




Fig.10

$$\begin{matrix} & \mathbf{N} & & & & & \mathbf{x} & & \mathbf{Nx} \end{matrix}$$
$$\begin{bmatrix} 1 & 0 & 1 & 1 & 1 \\ 0 & 1 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 1 \\ 2 \\ 1 \end{bmatrix} \geq \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$

That is { v_1, v_4} is a dominating set for G.
We know that, a graph G is just excellent if every vertex is contained in a unique γ - set of G.
If k does not divide n, then G is not just excellent. If k divides n, then we proceed as follows.

Consider the non – zero columns of NV. Let Xbe the set of all vectors ∋$Nx_i \geq 1$, i.e., $Ns \geq 1$. Let $|s| = q$, since p represents all possible permutation, there exist two vectors x_i, x_j, $i \neq j$ such that both x_i and x_j have non – zero entry in corresponding position, this mean that a vertex is contained in more than one γ - set. If $| q | = | p |$, then we mean to tell that every x_i is a γ - set for G. So atleast one vertex is contained in more than one γ - set, implies G is not just excellent.

If $q/n \neq k$, then R1 is not satisfied implies G is not just excellent.

If $q/n = k$, then add every row of matrix of order $n \times 1$. If every entry of the resulting column matrix is one, then every vertex is contained in exactly one γ - set implies G is just excellent else if atleast one entry of this matrix is not equal to one implies G is not just excellent.

If G is just excellent ∋ $\gamma(G) = 3$, then $\overline{G}$is non – outer planar. If G is just excellent ∋ $\gamma(G) \geq 4$, then $\overline{G}$is non – planar.

Example




Fig.11

For the graph in Fig. 11, $\gamma(G) = 2$. We consider all possible subsets with 2 vertice sand label them as $\{ S_1, S_2, S_3, \ldots, S_{15} \} =$ $\{\{ v_1, v_2\}, \{ v_1, v_3\}, \{ v_1, v_4\}, \{ v_1, v_5\}, \{ v_1, v_6 \}, \{ v_2, v_3 \}, \{ v_2, v_4 \}, \{ v_2, v_5\}, \{ v_2, v_6 \}, \{ v_3, v_4 \}, \{ v_3, v_5 \}, \{ v_3, v_6 \}, \{ v_4, v_5 \}, \{ v_4, v_6 \}, \{ v_5, v_6 \} \}$.

$$NV = \begin{pmatrix} 1&1&0&0&0&1 \\ 1&1&1&0&0&0 \\ 0&1&1&1&0&0 \\ 0&0&1&1&1&0 \\ 0&0&0&1&1&1 \\ 1&0&0&0&1&1 \end{pmatrix} \begin{pmatrix} 0&0&0&0&0&0&0&0&0&1&1&1&1&1 \\ 0&0&0&0&0&0&1&1&1&1&0&0&0&0&1 \\ 0&0&0&1&1&1&0&0&0&1&0&0&0&1&0 \\ 0&1&1&0&0&1&0&0&1&0&0&0&1&0&0 \\ 1&0&1&0&1&0&0&1&0&0&0&1&0&0&0 \\ 1&1&0&1&0&0&1&0&0&0&1&0&0&0&0 \end{pmatrix}$$

$$\begin{pmatrix} 1&1&0&1&0&0&2&1&1&1&2&1&1&1&2 \\ 0&0&0&1&1&1&1&1&1&2&1&1&1&2&2 \\ 0&1&1&1&1&2&1&1&2&2&0&0&1&1&1 \\ 1&1&2&1&2&2&0&1&1&1&0&1&1&1&0 \\ 2&2&2&1&1&1&1&1&1&0&1&1&1&0&0 \\ 2&1&1&1&1&0&1&1&0&0&2&2&1&1&1 \end{pmatrix}$$


$$S = \begin{pmatrix} 1&0&0 \\ 0&1&0 \\ 0&0&1 \\ 1&0&0 \\ 0&1&0 \\ 0&0&1 \end{pmatrix} \quad z = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}$$

where z is the matrix obtained by adding each rows of S. Since all the entries in the Z matrix is 1, the graph seen in Fig.10 is just excellent.

.

MAT LAB program for just excellent graphs

Snapshot – 1 provides the output for the graph in Fig. 11.



Snapshot - 1

6. Conclusion

The difficulty in constructing $\overline{G}$ particularly, when G has large number of vertices and less number of edges resulted in experimenting if $\overline{G}$ property can be determined without constructing $\overline{G}$. Is it possible to achieve this with the same number of graph operations originally used in $\overline{G}$? In this paper, we have devised a technique of determining the chromatic polynomial of $\overline{G}$ without actual construction of $\overline{G}$, but with the same number of operations used in $\overline{G}$. This result paved way for new method of approaching graph problems. Deciding $\overline{G}$ properties without constructing $\overline{G}$. As an output, non – planarity of $\overline{G}$ is determined from properties of G without knowing the actual structure of $\overline{G}$. We further plan to extend this idea and establish properties of $\overline{G}$ without constructing $\overline{G}$.

References

[1] Yamuna, M., &Karthika, K, " G^*properties – without G^*construction", *WSEAS Transaction on Mathematics,*Vol*15*, (2016), pp312 – 314.

[2] Yamuna, M., &Sridharan, N ,"Just excellent graphs", *International Journal of Engineering Science Advanced Computing and Bio-Technology*, Vol1 (3),(2010).pp.129-136.

[3] Yamuna, M., &Karthika, K. "Domatic subdivision stable and just excellent graphs". *International Journal of Applied Sciences and Innovation*,Vol 2, (2015), pp.56-60.

[4] Yamuna, M., Elakkiya, A., "Non domination subdivision stable graphs", *IOP Conf. Series: Materials Science and Engineering*. Vol 263, (2017).

[5] Yamuna, M., Elakkiya, A, "Planar graph characterization of NDSS graphs", *IOP Conf. Series: Materials Science and Engineering* ,Vol 263 ,(2017).

[6] M. Yamuna, A. Elakkiya, "γ - Uniquely colorable graphs*", IOPConf. Series: Materials Science and Engineering* , Vol.263 ,(2017).

[7] M. Yamuna, A. Elakkiya," Planar graph characterization of γUniquely colorable graphs", *IOP Conf. Series: Materials Science and Engineering* ,Vol263, (2017).

[8] Harary, F,*Graph Theory*, Addison Wesley, Narosa Publishing House, (2001).

[9] Haynes, T.W., Hedetniemi, S. T & Slater, P. J. *Fundamentals of domination in graphs*, New York, Marcel Dekker, (1998).