International Journal of Engineering & Technology, 7 (4.10) (2018) 1007-1011

International Journal of Engineering & Technology

Website: www.sciencepubco.com/index.php/IJET

Research paper



K[th] Root Transformation for a subclass of Log-Sigmoid Analytic Function based on Quasi-Subordination

M.Hari Priya[1]*, R. Bharavi Sharma[2], V.Suman Kumar[3]

[1]*Department of Mathematics, Kakatiya University, Warangal, Telangana-506009. INDIA*
[2]*Department of Mathematics, Kakatiya University, Warangal, Telangana-506009, INDIA.*
[3]*Department of Mathematics, TSMS, Karimnagar, Telangana, INDIA, 505481.*
**Corresponding author E-mail: maroju.hari@gmail.com*

Abstract

In the present investigation, using the concept of quasi-subordination, two subclasses of analytic functions have been introduced. The coefficient inequalities, the Fekete-Szego inequality, upper bounds for k[th] root transformation were studied. This study is extended to function f^{-1} and for $\frac{z}{f(z)}$.

Keywords: *Analytic function, Starlike function, Convex function, Quasi-subordination, Log-Sigmoid function, k^{th} root transformation*

1. Introduction

The study of Sigmoid functions plays a vital role in geometric function theory. These functions are usual in function analysis related to univalent function theory and sigmoid function is differential every where which is represented by truncated power series and is usually written as

$$h(z)=\frac{1}{1+e^{-z}} \tag{1}$$

This function is monotonically increasing and one-to-one. Let the family of analytic functions in the open disc Δ by A which takes the form

$$f(z)=z+\sum_{n=2}^{\infty}a_n z^n \tag{2}$$

are normalized by $f(0)=0$ and $f'(0)=1$. Thus without loss of generality an univalent analytic function can be written in the form of equation (2) denoted by S. Let f and g be two analytic functions in Δ. The function f is said to be subordinate to g, if there exists a Schwarz function w such that $f(z)=g(w(z))$. It is denoted by $f \prec g$. Furthermore, if g is univalent in Δ then the following equivalence satisfies as $f(0)=g(0)$ and $f(\Delta)\subset g(\Delta)$.

The concept of quasi subordination was introduced by Robertson [17]. Let f and g be two analytic functions in Δ. The function f is said to be quasi-subordinate to g, if there exists φ and w are two analytic functions which satisfies as $|w(z)|<1$ such that $f(z)=\varphi(z)g(w(z))$. It is written as Δ. If $\varphi(z)=1$, then $f(z)=g(w(z))$ so that $f \prec g$ in Δ. If $\varphi(z)=z$, then $f(z)=zg(w(z))$ and it is said that f is majorized by g and written by $f \square g$ in Δ. Both subordination and majorization are generalized by quasi subordination.

The problem of finding the maximum value of the coefficient functional $|a_3-\mu a_2^2|$, where μ is a real or complex parameter for the class of univalent analytic functions. This work is initiated by Fekete-Szego coefficient functional [6]. The k^{th} root transformation for a univalent analytic function f of the form in a equation (1.2) is denoted by $F(z)$ and is given by

$$F(z)=\left[f\left(z^k\right)\right]^{\frac{1}{k}}=z+\sum_{n=1}^{\infty}b_{nk+1}z^{nk+1}\left(\forall\, k\in N\right) \tag{3}$$

This transformation was studied by Ali et al.[1]. Several authors ([2], [10], [11], [21], [23], [24]), have studied the coefficient inequalities corresponding to the k^{th} root transformation for the function f in some subclasses of univalent and multivalent analytic functions. Olatunji et al. [15] introduced and studied the coefficient inequalities for the function f in the class $G_q^{\lambda}(\Phi,s,b)$, $G^{\lambda}(\Phi,s,b)$ and the class involving both quasi subordination and majorization. Murugusundaramoorthy and Janani [9] have studied the Fekete-Szego inequality for the univalent λ-pseudo starlike function in the space of sigmoid functions denoted by $L_{\beta}^{\lambda}(\Phi)$.

$$\mathrm{Re}\left(\frac{z(f'(z))^{\lambda}}{f(z)}\right)>\beta$$

Copyright © 2018 Authors. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Hamzat Jamiu Olusegun [7] have studied the coefficient inequalities for the function f in the Bazilevic subclass $B(\alpha,n,\gamma,\Phi)$. Several authors ([5], [7], [9], [13], [14], [22]), have introduced and studied the coefficient inequalities in the space of modified sigmoid and sigmoid functions.

Motivated by the above work, using the concept of quasi subordination two new subclasses of analytic functions denoted by $S_q^*(f,f')(\Phi)$ and $S_q^*(f,f',g)(\Phi)$ in the space of modified sigmoid function has been introduced. The coefficient inequalities, Fekete-Szegö inequality, upper bounds for k^{th} root transformation for the function f in these classes are obtained. This study have been extended for f^{-1} and for $\frac{z}{f(z)}$.

2. Preliminaries

Definition 1: The class of functions f analytic in the unit disc Δ, normalized by $f(0)=f'(0)-1=0$ and satisfies the condition

$$(1-\alpha)\left(\frac{zf'(z)}{f(z)}\right)^{\beta}+\alpha\left(\frac{(zf'(z))'}{f(z)}\right)^{1-\beta}\prec_q \Phi(z)$$

This class is denoted by $S_q^*(f,f')(\Phi)$.

Definition 2: The class of functions f analytic in the unit disc Δ, normalized by $f(0)=f'(0)-1=0$ and satisfies the condition

$$(1-\alpha)\left(\frac{z(f*g)'(z)}{(f*g)f(z)}\right)^{\beta}+\alpha\left(\frac{(z(f*g)'(z))'}{(f*g)(z)}\right)^{1-\beta}\prec_q \Phi(z)$$

This class is denoted by $S_q^*(f,f',g)(\Phi)$. For $g(z)=\frac{z}{1-z}$, we have $(f*g)=f$.

Now we recall the following Lemmas:

Lemma 3. [4] Let $p(z)=1+c_1z+c_2z^2+c_3z^3+.....$ be an analytic function in Δ satisfying $p(0)=1$ and $\mathrm{Re}\{p(z)\}>0$, for all $z\in\Delta$, then $|c_n|\le 2\ \forall n\ge 1$. The class of all such functions with positive real part is denoted by P.

Lemma 4. [8] Let $w(z)=w_1z+w_2z^2+w_3z^3+.....$, then $|w_1|\le 1$, $|w_2-tw_1^2|\le 1+(|t|-1)|w_1^2|\le \max\{1,|t|\}$, where $t\in C$

Lemma 5. Let h be a sigmoid function and

$$\Phi(z)=2h(z)=1+\sum_{m=1}^{\infty}\frac{(-1)^m}{2^m}\left(\sum_{n=1}^{\infty}\frac{(-1)^n}{n!}z^n\right)^n$$ then $|\Phi_{n,m}(z)|<2$ and $\Phi(z)\in P,\ |z|<1$ where $\Phi(z)$ is a modified sigmoid function.

Lemma 6. If $\Phi(z)\in P,\ |z|<1$ and it is starlike then f is a normalized univalent function of the form (1.2). Taking $m=1$, Joseph et al.[5] remarked the following:

Remark : Let $\Phi(z)=1+\sum_{n=1}^{\infty}C_nz^n$ where $C_n=\frac{-1(-1)^n}{2n!}$ then $|C_n|\le 1$, n=1,2,3,….. this result is sharp for each n

3. Coefficient estimates for the transforms of the function $f\in S_q^*(f,f',g)(\Phi)$

Theorem 7. If $f\in S_q^*(f,f',g)(\Phi)$, then

$$|b_{k+1}|\le\frac{1}{2kg_2(\beta-3\alpha\beta+2\alpha)} \tag{4}$$

$$|b_{2k+1}|\le\frac{1}{2kg_3(\beta+3\alpha-4\alpha\beta)}\left\{\frac{1}{2}+\max\left[1,|t_1|\right]\right\} \tag{5}$$

Where

$$t_1=\frac{1}{2}+\frac{1}{4(\beta+2\alpha-3\alpha\beta)^2}[g_2^2(5\beta^2+3\beta+3\alpha\beta-\alpha\beta^2-4)\ \frac{g_3(k-1)(\beta+3\alpha-4\alpha\beta)}{kg_2^2}] \tag{6}$$

Proof. If $f\in S_q^*(f,f',g)(\Phi)$, then there exists a Schawrz function w with $w(0)=0$ and $|w(z)|\le 1$ such that

$$(1-\alpha)\left(\frac{z(f*g)'(z)}{(f*g)f(z)}\right)^{\beta}+\alpha\left(\frac{(z(f*g)'(z))'}{(f*g)(z)}\right)^{1-\beta}=\psi(z)\left[\Phi(w(z))-1\right] \tag{7}$$

where the function $\Phi(z)$ is a modified sigmoid function defined as

$$\Phi(z)=1+\frac{z}{2}-\frac{z^3}{24}+\frac{z^5}{240}-\frac{z^6}{64}+\frac{779}{20160}z^7-..... \tag{8}$$

The analytic function, $\psi(z)$ in Δ, of the form

$$\psi(z)=c_0+c_1z+c_2z^2+c_3z^3+..... \tag{9}$$

Define a function $P(z)$ such that

$$w(z)=\frac{P(z)-1}{P(z)+1}\ \text{for some}\ P\in P \tag{10}$$

From (8) and (10), by applying $w(z)$ on the right hand side of the (7) after simplification, we get

$$\psi(z)\left[\Phi(w(z))-1\right]=\frac{p_1c_0}{4}z+\left(\frac{p_1c_1}{4}+\frac{p_2c_0}{4}-\frac{p_1^2c_0}{4}\right)z^2+\left(p_3c_0-p_1p_2c_0+\frac{5p_1^3c_0}{24}+\frac{p_1c_2}{4}+\frac{c_1}{4}\left(p_2-\left[p_2-\frac{p_1^2}{2}\right]\right)\right)z^3+..... \tag{11}$$

By Simple computation on the left hand side of (7), we have

$$(1-\alpha)\left(\frac{z(f*g)'(z)}{(f*g)f(z)}\right)^{\beta}+\alpha\left(\frac{(z(f*g)'(z))'}{(f*g)(z)}\right)^{1-\beta}=(1-\alpha)[1+\beta a_2g_2z+\left(2\beta a_3g_3+\frac{\beta(\beta-1)}{2}a_2^2g_2^2\right)z^2.....]+\alpha[1+2(1-\beta)a_2z+\left(2(1-\beta)\left(3a_3g_3-(\beta+2)a_2^2g_2^2\right)\right)z^2.....] \tag{12}$$

By applying k[th] root transformation and comparing the coefficients of z and z^2 from (7), (11) and (12), we get

$$b_{k+1}=\frac{p_1c_0}{4kg_2(\beta+2\alpha-3\alpha\beta)} \tag{13}$$

$$b_{2k+1}=\frac{1}{2kg_3(\beta+3\alpha-4\alpha\beta)}\left\{\frac{p_1c_1}{4}+\frac{c_0}{4}[p_2-t_1p_1^2]\right\} \tag{14}$$

where t_1 is given by (6). Upon simplification on the right hand side of equation (13) by applying Lemma 3 one can obtain the result as in (4). Similarly by taking modulus and applying Lemma 4 one can obtain the result as in (5). The sharpness of the result can be obtained by the rotations of the functions $w(z)=z$ and $w(z)=z^2$.

4. Fekete-Szegö coefficient function for $f\in S_q^*(f,f'g)(\Phi)$

Theorem 8. If $f\in S_q^*(f,f'g)(\Phi)$, then

$$\left|b_{2k+1}-\mu\, b_{k+1}^2\right|\le\frac{1}{2kg_3(\beta+3\alpha-4\alpha\beta)}\max\ \{1,\ |t_2|\} \tag{15}$$

where

$$t_2=\frac{1}{2}+\frac{1}{4(\beta+2\alpha-3\alpha\beta)^2}[g_2^2(5\beta^2+3\beta+3\alpha\beta-\alpha\beta^2-4)\ \frac{g_3\left[(k-1)+2\mu\right](\beta+3\alpha-4\alpha\beta)}{kg_2^2}] \tag{16}$$

The result is sharp.

Proof. From relations (13) and (14), consider

$$\left[b_{2k+1}-\mu b_{k+1}^2\right]=\frac{1}{2kg_3(\beta+3\alpha-4\alpha\beta)}\left\{\frac{p_1y}{4}+\frac{c_0}{4}\left[p_2-t_2p_1^2\right]\right\} \tag{17}$$

Since $\psi(z)$ given by (9) is analytic and bounded in Δ, therefore on using [12], we have for some $y(|y|\le1)$: such that $|c_0|\le1$ and $c_1=(1-c_0^2)y$. On substituting the value of c_1 in (17), we get

$$\left[b_{2k+1}-\mu b_{k+1}^2\right]=\frac{1}{2kg_3(\beta+3\alpha-4\alpha\beta)}\left\{\frac{p_1y}{4}+\frac{c_0}{4}\left[p_2-t_2p_1^2\right]-\frac{p_1c_0^2y}{4}\right\} \tag{18}$$

If $c_0=0$ in (18), one can be seen that

$$\left[b_{2k+1}-\mu b_{k+1}^2\right]=\frac{p_1c_1}{8kg_3(\beta+3\alpha-4\alpha\beta)} \tag{19}$$

Taking modulus on both sides and by applying Lemma 4 on the right hand side of (19), one can obtain the result

$$\left|b_{2k+1}-\mu b_{k+1}^2\right|\le\frac{1}{4kg_3(\beta+3\alpha-4\alpha\beta)} \tag{20}$$

If $c_0\ne0$ in (18), then suppose that

$$F(c_0):=\left\{\frac{p_1y}{4}+\frac{c_0}{4}\left[p_2-t_2p_1^2\right]-\frac{p_1c_0^2y}{4}\right\} \tag{21}$$

Which is a polynomial in c_0 and hence analytic in $|c_0|\le1$, and maximum value is attained at $c_0=e^{i\theta}$,

$$\max_{0\le\theta\le2\pi}\left|F\left(e^{i\theta}\right)\right|=\left|F(1)\right| \tag{22}$$

Here $$\left|b_{2k+1}-\mu b_{k+1}^2\right|\le\frac{1}{2kg_3(\beta+3\alpha-4\alpha\beta)}\left|p_2-t_2p_1^2\right| \tag{23}$$

By applying the Lemma 3 on the right hand side of (23), one can obtain the result as in (15).

If $g(z)=\frac{z}{1-z}$, then $(f*g)=f(z)$, the above result can be reduced to the following corollary.

Corollary 9. If $f\in S_q^*(f,f')(\Phi)$, then

$$\left|b_{2k+1}-\mu\, b_{k+1}^2\right|\le\frac{1}{2k(\beta+3\alpha-4\alpha\beta)}\max\ \{1,\ |t_3|\} \tag{24}$$

Where

$$t_3=\frac{1}{2}+\frac{1}{4(\beta+2\alpha-3\alpha\beta)^2}[(5\beta^2+3\beta+3\alpha\beta-\alpha\beta^2-4)\ \frac{\left[(k-1)+2\mu\right](\beta+3\alpha-4\alpha\beta)}{k}] \tag{25}$$

The result is sharp.

5. Coefficient inequality for the function $\frac{z}{f(z)}$

Theorem 10. If $f\in S_q^*(f,f')(\Phi)$ and $G(z)=\frac{z}{f(z)}$**,** then for any real number μ, we have

$$\left|q_2-\mu\, q_1^2\right|\le\frac{1}{2(\beta+3\alpha-4\alpha\beta)}\max\{1,|t_4|\} \tag{26}$$

Where

$$t_4=\frac{1}{2}+\frac{1}{4(\beta+2\alpha-3\alpha\beta)^2}\left\{5\beta^2+5\beta-5\alpha\beta+6\alpha-\alpha\beta^2-4-2\mu\right\} \tag{27}$$

The result is sharp.

Proof.

As $f\in S_q^*(f,f')(\Phi)$ and

$$G(z)=\frac{z}{f(z)}=1+\sum_{n=1}^{\infty}q_nz^n \tag{28}$$

By a simple computation we get

$$\frac{z}{f(z)}=1-a_2z+\left(a_2^2-a_3\right)z^2-..... \tag{29}$$

Upon equating the coefficients of z and z^2, from relations (27) and (28), we have

$$q_1=-a_2 \tag{30}$$

$$q_2 = -a_3 + a_2^2 \tag{31}$$

From equations (13), (14), (29) and (30), it can be obtained

$$q_1 = -\frac{p_1 c_0}{4(\beta + 2\alpha - 3\alpha\beta)} \tag{31}$$

$$q_2 = -\frac{1}{2(\beta + 3\alpha - 4\alpha\beta)}\{\frac{p_1 c_1}{4} + \frac{c_0}{4}(p_2 - p_1^2[\frac{1}{2} + \frac{c_0}{4(\beta + 2\alpha - 3\alpha\beta)^2}(5\beta^2 + 5\beta - 5\alpha\beta + 6\alpha - \alpha\beta^2 - 4)\}} \tag{32}$$

For any complex number μ, consider

$$q_2 - \mu\, q_1^2 = -\frac{1}{2(\beta + 3\alpha - 4\alpha\beta)}\{\frac{p_1 c_1}{4} + \frac{c_0}{4}(p_2 - t_4 p_1^2)\} \tag{33}$$

Where t_4 is given by (27). By applying similar computation as in Theorem 8, one can obtain the result as in (24).

6. Coefficient Inequalities for the function f^{-1}

Theorem 11. If $f^{-1}(w) = w + \sum_{n=2}^{\infty} d_n w^n$ is the inverse of $f \in S_q^*(f, f')(\Phi)$, then for any complex parameter μ, we have

$$\left|d_3 - \mu d_2^2\right| \le \frac{1}{2(\beta + 3\alpha - 4\alpha\beta)} \max\ \{1, |t_5|\} \tag{34}$$

Where

$$t_3 = \frac{1}{2} + \frac{c_0}{4(\beta + 2\alpha - 3\alpha\beta)^2}\{(5\beta^2 + 5\beta - 5\alpha\beta + 6\alpha - \alpha\beta^2 - 4 - 2\mu)\} \tag{35}$$

Proof.

As $f^{-1}(w) = w + \sum_{n=2}^{\infty} d_n w^n$ (36)

is the inverse function of f, it can be seen that

$$f^{-1}(f(z)) = f(f^{-1}(z)) = z \tag{37}$$

From equations (36) and (37), it can be reduced to

$$z + (a_2 + d_2)z^2 + (a_3 + 2a_2 d_2 + d_3)z^3 + \ldots\ldots\ldots = z \tag{38}$$

By comparing the coefficients of z and z^2, from relation (38), it can be seen that

$$d_2 = -\frac{p_1 c_0}{4(\beta + 2\alpha - 3\alpha\beta)} \tag{39}$$

$$d_3 = -\frac{1}{2(\beta + 3\alpha - 4\alpha\beta)}\{\frac{p_1 c_1}{4} + \frac{c_0}{4}(p_2 - p_1^2[\frac{1}{2} + \frac{c_0}{4(\beta + 2\alpha - 3\alpha\beta)^2}(5\beta^2 + 7\beta - 13\alpha\beta + 12\alpha - \alpha\beta^2 - 4)\}} \tag{40}$$

For any complex number μ, we have

$$d_3 - \mu\, d_2^2 = -\frac{1}{2(\beta + 3\alpha - 4\alpha\beta)}\{\frac{p_1 c_1}{4} + \frac{c_0}{4}(p_2 - t_5 p_1^2)\} \tag{41}$$

Where t_5 is given by (36). By applying similar computation as in Theorem 8 one can obtain the result as in equation (34).

Acknowledgements

1. The authors are very much thankful to Prof. G. Murugusundaramoorthy for his valuable suggestions and comments which helped in writing this article.
2. The work in this paper is partially supported by DST/FIST grant no: SR/FST/MSI-101/2014, dated 14-01-2016.

References

[1] Ali,R.M., Lee.S.K., Ravichandran. and Supramaniam,S ., "The Fekete –Szegö coefficient functional for transforms of analytic functions", Bulletin of the Iranian Mathematical Society, Vol. 35 No. 2, (2009), pp: 119-142.
[2] Annamalia.S, Ramachandran.C and Murugusundaramoorthy, "Fekete- Szegö coefficient for the Janowski α -spirallike function in open unit disc", International Journal of Mathematical Analysis, Vol.8, No.19, (2012), pp: 913-918.
[3] Babalola.K.O (@013), "On $\lambda-$ pseudo starlike functions", Journal of Classical Analysis, Vol.3, No.2, pp: 137-147.
[4] Duren.P.l, "Univalent Functions", Grundlehren der Mathematischen Wissenchaften, Vol.259, (1983) , Springer, New York.
[5] Faipe-Joseph.O.A, Oladipo.A.T. Ezeafulukwe.U.A. (2013), Modified sigmoid function in univalent function theory, International Journal of Mathematical Sciences and Engineering Application, Vol.7,No.7, pp:313-317.
[6] Fekete.M and Szego.G), "Eine Bermerkung uber ungerade schlichte function", Journal of London Mathematical Society, Vol.8, (1933), pp:85-89.
[7] Hamzat Jamiu Olusegun , "Coefficient bounds for bazilevic functions associated with modified sigmoid functions, Asian Research Journal of Mathematics, Vol.5, No.3, (2017), pp: 1-10.
[8] Keogh.F.R and Merkes.E.P., "A coefficient inequality for certain classes of analytic functions", Proceedings of the American Mathematical Society, Vol.20, (1969), pp:8-12.
[9] Murugusundarmoorthy.G and Janani.T, "Sigmoid function in the space of univalent λ -pseudo starlike functions", International Journal of Pure and Applied Mathematics, Vol.101, No.1, (2015), pp : 33-41.
[10] Nanjundan Magesh, Jagadeesan Yamini, "Fekete-Szego inequalities associated with k-th root transformation based on quasi-subordination", Annales Universitatis Paedagogicae Cracoviensis Studia Mathematica, Vol.16, (2017), pp :7-15.
[11] Najla.M, Alarifi and Maisarah Haji Mohd, "Best bounds of the second Hankel determinants of the kth root transform of analytic functions", AIP Conference Proceedings, Vol.1682, No.2, (2015).
[12] Nehari.Z, "Conformal Mapping", McGraw-Hill, New York, NY,USA, 1st Edition, .(1952).
[13] Olatunji.A.O, Gbolagde. M, Anake.T and Fadipe-Joseph. "Sigmoid function in the space of univalent function of Bazelevic type", Scientia Magna, Vol.97, No.3, (2013), pp :43-51.
[14] Olantunji.A.O, "Sigmoid function in the space of univalent $\lambda-$ pseudo starlike function with Sakaguchi type functions", Journal of Progressive Research in Mathematics (JPRM), Vol.7, No.4, pp: 1164-1172.
[15] Olantunji.S.O., Dansu.E.J and Abidemi.A. , "On a Sakaguchi type class of analytic functions associated with quasi-subordination in the space of modified sigmoid functions", Electronic Journal of Mathematical Analysis and Application, Vol.5, No.1, (2017), pp :97-105.
[16] Pommernke.C.(1966), On the coefficients and Hankel determinants of univalent functions, Journal of London Mathematical Society, Vol.41, pp: 111-122.
[17] Robertson.M.S., "Quasi-Subordination and coefficient conjecture", Bulletin of the American Mathematical Society, Vol.76, .(1970) pp: 1-9.

[18] Sharma.R.B, and Hari Priya.M, "On a class of $\alpha-$ convex functions subordinate to a shell shaped region", The Journal of Analysis, Vol.25, (2016), pp: 93-105.
[19] Singh.R.(1973), " On Bazilevic functions", Proceedings of the American Mathematical Society, Vol.28, pp: 261-271.
[20] Sharma.R.B. and Tam Reddy.T and Saroja.K, "A new subclass of Meromorphic functions with positive coefficients", Indian Journal of Pure and Applied Mathematics, Vol.44, No.1, .(2013).
[21] Sharma.R.B. and Tam Reddy.T and Hari Priya.M., "Kth root transformations for a subclass of p-valent functions", Bulletin of Calcutta Mathematical Society, Vol.107, No.4, .(2015), pp: 291-304.
[22] Sunday Olufemi Olantunji and Emmanuel Jesuyon Dansu, "Coefficient estimates for Bazilevic Ma-Minda functions in the space of sigmoid functions", Malaysian Journal of Mathematik, Vol.4, No.3, (2016), pp: 505-512.
[23] Vamshee Krishna.D, Venkateshwarlu.B, and Ram Reddy.T, "Coefficient inequality for transforms of reciprocal of bounded turning functions", Annales Univ.Sci. Budapest., Sect.Comp., Vol.44, (2015), pp: 69-77.
[24] Vamshee Krishna.D, Venkateshwarlu.B, Hankel determinant for transforms of pre-starlike functions of order alpha, South-East Asian Bulletin of Mathematics, Vol.40, (2016), pp: 131-140.