International Journal of Engineering & Technology, 7 (4.10) (2018) 880-882

International Journal of Engineering & Technology

Website: www.sciencepubco.com/index.php/IJET

Research paper



On Sαrw-Homeomorphism in Topological Spaces

Basavaraj M. Ittanagi [1] and Mohan V[2*]

[1]Department of Mathematics,
Siddaganga Institute of Technology, Tumakuru-03,
Affiliated to VTU, Belagavi, Karnataka state, India.
[2]Department of Mathematics,
Gopalan College of Engineering and Management, Bangalore-48,
Affiliated to VTU, Belagavi, Karnataka state, India.
Corresponding author E-mail: vengatachalam.mohan@gmail.com

Abstract

New class of homeomorphisms named as sαrw-homeomorphism and sαrw*-homeomorphism are explored & elaborated. Few basic properties are inspected. Their relations with some existing homeomorphisms in topological spaces are studied.

Keywords: *homeomorphism; sαrw-homeomorphism and sαrw*-homeomorphism.*

1. Introduction

The concept of generalized homeomorphism was introduced and studied in the year 1991 by Balachandran et all [2]. N. Nagaveni [7] introduced and studied rwg homeomorphism in topological spaces. In the year 2002, M Sheik John [10] introduced and studied w-homeomorphism in topological space and rgα-homeomorphisms, rps-homeomorphisms and gs and sg homeomorphisms have been introduced and studied by A. Vadivel [11], T. Shyla Isac Mary [8], H. Maki [6] respectively. This paper aims to explore and elaborate sαrw-homeomorphism, sαrw*-homeomorphism and its relation with some existing homeomorphisms in topological spaces. Few of its properties are investigated.

2. Preliminaries

Without considering any separation axioms, a topological space Y or (Y,τ) and Z or (Z,σ) is assumed. For B ⊆ Y, where Y is a topological space. Closure of B- cl (B), interior of B- int (B), and complement of B - Y-B or B^C

Definition 2.2: g: (Y, τ) → (Z, σ) is a map.
i) If $g^{-1}(F)$ is r-closed in Y ∀ closed F of Z then it is called as regular-continuous (in brief r- continuous) [1].
ii) If $g^{-1}(F)$ is r- closed in Y ∀ closed F of Z, then it is called as completely–continuous [1].
iii) If $g^{-1}(F)$ is g–closed in Y ∀ closed F of Z, then it is called as g– Continuous [2].
iv) If $g^{-1}(F)$ is sαrw–closed in Y ∀ closed F of Z, then it is called as sαrw–continuous [4].
v) If $g^{-1}(F)$ is closed set in Y ∀ sαrw–closed F in Z, then it is called as strongly sαrw–continuous [4].
vi) If $g^{-1}(F)$ is αrw–closed in Y ∀ closed F of Z, then it is called as αrw– continuous [12].

Definition 2.5: g: (Y, τ) → (Z, σ) is map
i) If g (F) is g-open in (Z, σ) ∀ open F of (Y, τ), then it is called as g-open map [2].
ii) If g (F) is gpr-open in (Z, σ) ∀ open set F of (Y, τ), then it is called as gpr-open map [5].
iii) If g (F) is open in (Z, σ) ∀ regular open set F of (Y, τ), then it is called as regular open map [9].
iv) If g (F) is rwg-open in (Z, σ) ∀ open set F of (Y, τ), then it is called as rwg-open map [7].
v) If g (F) is wg-open in (Z, σ) ∀ open set F of (Y, τ), then it is called as wg-open map [7].
vi) If g (F) is w-open in (Z, σ) ∀ open set F of (Y, τ), then it is called as w-open map [10].

Definition 2.6: g: (Y, τ) → (Z, σ) is a map.
i) If g is open and continuous then it is called as homeomorphism
ii) If g is g-continuous and g-open then it is called as g- homeomorphism [2].
iii) If g is rgα-continuous and rgα-open then it is called as rgα-homeomorphism [11].
iv) If g is rwg-continuous and rwg-open then it is called as rwg-homeomorphism [7].
v) If g is sg-continuous and sg-open then it is called as sg-homeomorphism [6].
vi) If g is gs-continuous and gs-open then it is called as gs- homeomorphism [6].
vii) If g is rps-open and rps-continuous then it is called as rps-homeomorphism [11].

Results 2.7 [3]:
i. In Y, each closed set (resp regular-closed, α-closed)is sαrw closed .
ii. In Y, each sαrw-closed set is sg-closed.
iii. In Y, each sαrw-closed set is gsp-closed (resp rps-closed, gs closed , gspr-closed) .

Copyright © 2018 Authors. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

3. sαrw-homeomorphism in topological Space

Definition 3.1: g: (Y, τ) → (Z, σ) is a bijective map and if g is sαrw-continuous as well as sαrw-open then we call it sαrw-homeomorphism.

Theorem 3.2: Each homeomorphism is sαrw-homeomorphism.
Proof: g: (Y, τ) → (Z, σ) is a homeomorphism. g is bijective, continuous and open because each continuous map is sαrw-continuous as well as each open map is sαrw-open. Thus f is sαrw-homeomorphism.

Remark 3.3: Each sαrw-homeomorphism need not homeomorphism.

Example 3.4: Let Y=Z={1, 2 ,3}, τ ={ϕ,{1},{2},{1, 2},Y} and σ ={ϕ,{1},{2},{1, 2},{1, 3}, Z }.g: Y→Z is defined as g (1)=2 , g(2)=3,g(3)=3,then g is sαrw-homeomorphism but not homeomorphism as closed set F= {2} in Z, $g^{-1}(F)$ ={1} in Y which isn't closed in Y.

Theorem 3.5: Each α-homeomorphism is sαrw-homeomorphism.

Proof: Let g: (Y, τ) → (Z, σ) be homeomorphism. g is bijective, continuous and open map. Because each α-continuous map is sαrw-continuous and each α-open map is sαrw-open, Thus g is sαrw-homeomorphism.

Remark 3.6: Each sαrw- homeomorphism need not be α- homeomorphism.

Example 3.7: From example 3.4 g is sαrw-homeomorphism however isn't α-homeomorphism as closed set F= {2} in Z, then $g^{-1}(F)$ = {1} in X which is n't α-closed in Y.

Theorem 3.8: Each sαrw–homeomorphism is gs-homeomorphism.
Proof: Let g: (Y, τ) → (Z, σ) be sαrw-homeomorphism. g is sαrw-continuous and sαrw-open map, which implies g is gs continuous and gs open map. Thus g is gs-homeomorphism.

Remark 3.9: Each gs –homeomorphism need not be sαrw - homeomorphism

Example 3.10: Let Y={1,2,3}, Y={1,2,3,4}, τ ={ϕ,{1},{2,3}, {1,2,3},Y} and σ ={ϕ,{1},{2},{1, 2},{1, 3 },Z}. g: X→Y be defined as g(1)=2 , g(2)= 3 , g(3)=1, then g is gs–homeomorphism but it isn't sαrw-homeomorphism, as a closed set F= {3} in Z and $g^{-1}(F)$ = {2} in Y which are not sαrw-closed in Y.

Theorem 3.11: Each sαrw–homeomorphism is sg-homeomorphism.

Proof: Let g: (Y, τ) → (Z, σ) be sαrw-homeomorphism. Then g is sαrw-continuous and sαrw-open map. This implies that g is sg continuous and sg open map. Thus g is sg-homeomorphism.

Remark 3.12: Each sg–homeomorphism need not be sαrw - homeomorphism.

Example 3.13: From example 3.10, g is sg–homeomorphism. But not sαrw-homeomorphism as closed set F= {3} in Z, then $g^{-1}(F)$ = {2} in Y which isn't sαrw-closed in Y.

Theorem 3.14: Each sαrw–homeomorphism is rps-homeomorphism.

Proof: Let g: (Y, τ) → (Z, σ) be sαrw-homeomorphism. Then g is sαrw-continuous and sαrw-open map. This implies that g is rps continuous and rps open map. Thus g is rps-homeomorphism.

Remark 3.15: Each rps–homeomorphism need not be sαrw - homeomorphism.

Example 3.16: From example 3.10, g is rps–homeomorphism, but not sαrw-homeomorphism since closed set F= {3} in Z, then $g^{-1}(F)$ = {2} in Y that is not sαrw-closed in Y.

Theorem 3.17: Each sαrw–homeomorphism is gsp-homeomorphism.

Proof: Assume that g: (Y, τ) → (Z, σ) is sαrw-homeomorphism, then g is sαrw-continuous and sαrw-open map. It shows that g is gsp continuous and gsp open map. Thus g is gsp-homeomorphism.

Remark 3.18: Each gsp–homeomorphism need not be sαrw-homeomorphism.

Example 3.19: From example 3.10, g is gsp–homeomorphism, but not sαrw homeomorphism as closed set F= {3} in Z, then $g^{-1}(F)$ = {2} in Y that isn't sαrw-closed in Y.

Theorem 3.20: Each sαrw–homeomorphism is gspr-homeomorphism.

Proof: Assume g: (Y, τ) → (Z, σ) be sαrw-homeomorphism, then g is sαrw-continuous and sαrw-open map. This implies that g is gspr continuous and gspr open map. Thus g is gspr-homeomorphism.

Remark 3.21: Each gspr–homeomorphism need not sαrw - homeomorphism.

Example 3.22: From example 3.10, g is gspr–homeomorphism. But not sαrw-homeomorphism as closed set F= {3} in Z, then $g^{-1}(F)$ = {2} in Y which isn't sαrw-closed in Y.

Remark 3.23: The example below illustates sαrw-homeomorphisms is independent of pre–homeomorphism, β–homeomorphism, gp–homeomorphism, gpr–homeomorphism, swg–homeomorphism, rwg–homeomorphism, wg–homeomorphism, gprw–homeomorphism, rgw–homeomorphism, pgpr–homeomorphism.

Example 3.24: Assume Y={1, 2 ,3, 4}, Z ={1, 2, 3}, τ = { ϕ,{1},{2, 3},{1, 2, 3},Y} and σ ={ϕ,{1},{2},{1 ,2},{1, 3},Z}. g: X→Y be defined as g(1)=2 , g(2)= 3 , g(3)=1, then g is pre–homeomorphism, β–homeomorphism, gp–homeomorphism, gpr–homeomorphism, swg–homeomorphism, rwg–homeomorphism , wg–homeomorphism, gprw–homeomorphism, rgw-homeomorphism, pgpr– homeomorphism, but not sαrw-homeomorphism as closed set F= {3} in Z, then $g^{-1}(3)$ = {2} in Y that isn't sαrw-closed in Y.

Theorem 3.25: Assume that g: Y→Z be bijective and sαrw continuous . Then the below facts are true.

a. g is sαrw-open.
b. g is sαrw-homeomorphism
c. g is sαrw-closed.

Proof:

1. g is sαrw-open.⇒ g is sαrw-homeomorphism
g: Y→Z is bijective, sαrw-continuous map and sαrw-open map. Thus g is a sαrw-homeomorphism.
2. g is sαrw-homeomorphism ⇒ g is sαrw-closed.
g: Y→Z is sαrw-homeomorphism. Let G be any closed in Y and hence G^{c} is closed in Y. Then g (G^{c}) is open in Z. But $g(G^{c})=g(G)^{c}$. Thus g is a sαrw-closed.
3. g is sαrw-closed.⇒ g is sαrw-open.

g: Y→Z is bijective, sαrw-continuous and sαrw-closed map. Let G be any open in Y and hence G^c is closed set in Y. Then $g(G^c)$ is closed in Z. But $g(G^c)=g(G)^c$. Thus g is sαrw-open.

Remark 3.26: Ccomposition of 2 sαrw-homeomorphisms is need not be a sαrw-homeomorphism .it is illustrated below

Example 3.27: Assume Y=Z=M= {1, 2, 3}, τ = {ϕ, {1}, {2},{1, 2}, {1, 3},Y} , σ = {ϕ, {1}, {2}, {1, 2},Z} and η = {ϕ, {1}, {2, 3},M}.We define g: (Y, τ) → (Z, σ) by g (1) =1, g (2) =3, g(3) =2 and h: (Y, σ) →(M, η) be the identity mapping. Both g and h are sαrw-homeomorphism but their composition (hog): (Y, τ)→(M, η) isn't a sαrw-homeomorphism as the closed set F={1} in (M, η), but $(hog)^{-1}(F)$={1} isn't sαrw-closed in Y.

Definition 3.28: g: Y→Z is bijective map. Then it is sαrw*-homeomorphism if g and g^{-1} are sαrw-irresolute. Where sαrw*-h(Y,τ) represents the set of all sαrw-homeomorphisms of Y on to itself.

Theorem 3.29: Each sαrw*-homeomorphism is a sαrw-homeomorphism.

Proof: Let g: Y→Z be a sαrw*-homeomorphism. Then g is bijective, g and $g^{-1}$are sαrw-irresolute. Because each sαrw-irresolute map is sαrw-continuous then g and g^{-1} are sαrw-continuous. Also g^{-1}: Z→Y is a sαrw-continuous map is equivalent to sαrw-open map. Thus g is bijective, sαrw-continuous and sαrw-open map. Hence g is a sαrw-homeomorphism.

Each sαrw-homeomorphism need not be sαrw*-homeomorphism.

Example 3.30: Assume Y=Z={1 ,2 ,3}, τ ={ϕ,{1},{2},{1,2},Y} and σ ={ϕ,{1},{2},{1, 2},{1,3},Z}.Let g: Y→Z defined by g(1)=2, g(2)= 3 , g(3)=3, then g is sαrw–homeomorphism but not sαrw*-homeomorphism, because g isn't sαrw-irresolute.

Theorem 3.31: Each sαrw*-homeomorphism is a sg-homeomorphism.

Proof: The proof follows from the facts that each sαrw*-homeomorphism is a sαrw-homeomorphism and each sαrw-homeomorphism is a sg-homeomorphism.
Each sg-homeomorphism need not be sαrw*-homeomorphism.

Example 3.32: Assume Y= {1, 2, 3}, Z= {1, 2, 3, 4}, τ = {ϕ, {1}, {2,3}, Y} and σ = {ϕ, {1}, {2}, {1, 2}, {1, 3}, {1, 2, 3}, Z}. Let g: Y→Z defined by g(1)=2 , g(2)= 3 , g(3)=1, then g is sg–homeomorphism. But not sαrw-homeomorphism as closed set F= {3} in Z, then $g^{-1}(F)$ = {2} in Y which is not sαrw-closed set in Y.

Theorem3.33: Each sαrw*-homeomorphism is a gs-homeomorphism (resp gsp-homeomorphism, rps-homeomorphism, gspr-homeomorphism).
Proof: we know that each sαrw*-homeomorphism is a sαrw-homeomorphism and each sαrw-homeomorphism is a gs-homeomorphism (resp gsp-homeomorphism, rps-homeomorphism, gspr-homeomorphism)
Each gs-homeomorphism is a sαrw*-homeomorphism

Example 3.34: From example 3.27, g is gs-homeomorphism (resp) gsp-homeomorphism, rps-homeomorphism, gspr homeomorphism), but not sαrw-homeomorphism, as closed set F= {3} in Z, then, $f^{-1}(F)$ = {2} in Y which isn't sαrw-closed set in Y.

Remark 3.35: From discussions and facts already known ,relation between sαrw-homeomorphism with some existing homeomorphisms in topological space is shown in following figure.




Figure-1

Theorem 3.36: The composition of two sαrw*-homeomorphisms is also a sαrw*-homeomorphism.

Proof: Assume g: X→Y and h: Y→Z is two sαrw*-homeomorphisms. To prove hog: X→Z is sαrw*-homeomorphism. Let V be any closed set in Z. Because h is sαrw-irresolute, h(V) is sαrw-closed in Y. Because g is sαrw-irresolute, $g^{-1}(h^{-1}(V))$ is sαrw-closed in X. But $g^{-1}(h^{-1}(V)) = (hog)^{-1}(V)$. Thus hog is sαrw-irresolute. Let W be any closed in X. Because g is sαrw-irresolute, g (W) is sαrw-closed in Y and also h (g (W)) is sαrw-closed in Z. But (hog) (W) =h(g(W)). Thus (hog) is sαrw-irresolute. Hence (hog) is sαrw*-homeomorphism.

Theorem 3.37: Under the composition of maps, sαrw*-h (Y, τ) is group.

Proof:* be binary operation,*:sαrw*h (Y, τ) (x) sαrw*h (Y, τ) → sαrw*h (Y, τ) defined by f * g = g o f ∀ f, g ∈ sαrw*h (Y, τ). We have (gof) ∈ sαrw*h(X, τ). Also the associative property holds in composition of maps. The identity mapping I: (Y, τ) → (Y, τ) which ∈ sαrw*h (Y, τ), acts as the identity element. For the inverse element, if f ∈ sαrw*h (Y, τ) then f^{-1} ∈ sαrw*h(Y, τ) such that $fof^{-1}= f^{-1}of=I$. Thus, under the composition of maps, the set sαrw*-h(Y, τ) is a group.

References

[1] Arya S P, Gupta R [1974], On strongly continuous functions, Kyungpook Math.J 14,131-143.
[2] Balachandran K, Sundaram P, Maki H [1991], On Generalized homeomorphisms in Topological Spaces, Fukuoka University, Ed., part III.40, 13-21.
[3] Basavaraj M Ittanagi and Mohan V [2017], on semi α-regular weakly closed set in Topological spaces, Int J. of Math Archive 8(7), 197-204.
[4] Basavaraj M. Ittanagi and Mohan V , On sαrw continuous and sαrw irresolute Maps in Topological Spaces (Accepted paper in JUSPS)
[5] Gnanambal Y [1997], on generalized pre regular closed sets in topological spaces, Indian J Pure. Appl. Math. 28(3), 351-360.
[6] Maki H, K. Balachandran, R.Devi [1982], Semi-generalized homeomorphisms and generalized semi homeomorphisms in Topological Spaces, Indian J.of Closed Maps, J. Karnatk Univ. Sci. 27 , 82-88.
[7] Nagaveni N [1999],Studies on Generalizations of Homeomorphisms in Topological Spaces, Ph.D. Thesis, Bharathiar University,Coimbatore,
[8] Shyla Isac Mary T, P. Thangavelu [2013], rps-homeomorphisms in Topological Spaces, Asian Journal of Current Engineering and Maths. 2(1), 74-76.
[9] Stone M [1937], Application of the theory of Boolean rings to general topology, Trans. Amer. Math. Soc. 41,374- 481.
[10] Sundaram P, Sheik John M [2000], On w-closed sets in topology, Acta Ciencia Indica 4,389-39.
[11] Vadivel A, K. Vairamanickam [2010], rgα-Homeomorphisms in Topological Spaces, Int. Journal of Math. Analysis. 4, 881-890.
[12] Wali R.S., Mandalgeri P.S [2014], on αrw–Continuous and αrw–Irresolute Maps in Topological Spaces 10(6), 14-24.