International Journal of Engineering &Technology, 7 (4) (2018) 6947-6950

International Journal of Engineering & Technology

Website: www.sciencepubco.com/index.php/IJET
doi: 10.14419/ijet.v7i4.21509
Research paper

On gaussian diophantine quadruples

S.Vidhyalakshmi[1], M.A.Gopalan[1], S. AarthyThangam[2] *

[1]Professor, Department of Mathematics, Shrimati Indira Gandhi College, Trichy-620 002, Tamil Nadu, India
[2]Research Scholar, Department of Mathematics, Shrimati Indira Gandhi College, Trichy-620 002, Tamil Nadu, India
**Corresponding author E-mail:aarthythangam@gmail.com*

Abstract

This paper concerns with the problem of constructing gaussian diophantine quadruples with the property that the product of any two distinct gaussian integers added with 1 and 4 in turn is a perfect square. The construction of gaussian diophantine quadruple (A,B,C,D) is illustrated through employing the non-zero distinct integer solutions of the system of double diophantine equations. The repetition of the above process leads to the generation of sequences of gaussian diophantine quadruples with the given property.

Keywords: *Diophantine Quadruple; Double Diophantine Equations; Gaussian Diophantine Quadruples; Integer Solutions; Pell Equations.*

1. Introduction

The construction of the sets with property that the product of any two of its distinct elements is one less than a square has a very long history and such sets were studied by Diophantus. A set of m distinct non-zero integers $\{a_1,a_2,.....,a_m\}$ is called a Diophantine m-tuple with property $D(n)$ if a_ia_j+n is a perfect square for all $1\le i\le j\le m$ [1]. Various mathematicians discussed the construction of different formulations of Diophantine triple and diophantine quadruples with property $D(n)$ for any arbitrary integer n and also for polynomials in n [2-15].

A set $\{a_1,a_2,.....,a_m\}\subset Z(i)-\{0\}$ is said to have this property $D(z)$ if the product of its any two distinct elements increased by z is a square of a Gaussian integer. If the set $\{a_1,a_2,.....,a_m\}$ is a complex diophantine quadruple then the same is true for the set $\{-a_1,-a_2,.....,-a_m\}$. Particularly in [16], the authors have analyzed the problem of the existence of the complex diophantine quadruples. In this context, one may refer [17-25].

In this communication, we construct sequences of gaussian diophantine quadruples with properties $D(1)$ and $D(4)$.

2. Method of analysis

2.1. Problem 1:

Let A, B be two gaussian integers represented by

$$A=kp\pm k+ikq\ ,\ B=kp\pm(2n+k)+ikq$$

where k, p, q and n are non-zero integers.
Note that

$$AB+n^2=\left(kp\pm(n+k)+ikq\right)^2=r^2\ \text{(say)}$$

Therefore, the pair (A,B) is a gaussian diophantine 2-tuple with property $D(n^2)$

Consider C to be a gaussian integer such that

$$AC+n^2=\alpha^2 \qquad (1)$$

$$BC+n^2=\beta^2 \qquad (2)$$

Assume

$$\alpha=A+r\ ,\ \beta=B+r \qquad (3)$$

Substituting (3) in (1) and (2) and subtracting one from the other, observe that

$$C=A+B+2r=4kp\pm4(n+k)+i\,4kq$$

It is observed that the triple (A,B,C) is a gaussian diophantine 3-tuple with property $D(n^2)$. When $n=1,n=2$, the above triple (A,B,C) can be extended to diophantine quadruple with their corresponding properties.

2.1.1. Diophantine quadruple with property $D(1)$:

Let $n=1$. Then the triple (A,B,C) is given by

$$A=kp\pm k+ikq$$

$$B=kp\pm(2+k)+ikq$$

$$C=4kp\pm4(k+1)+i\,4kq$$

which is a gaussian diophantine triple with property $D(1)$.

If (A,B,C) is a diophantine triple with property $D(1)$ then the fourth tuple D is given by

Copyright ©2018 S.Vidhyalakshmi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

$$D=(A+B+C)+2(ABC+\alpha\beta r) \quad (4)$$

where

$$AC+1=\alpha^2\ ,\ BC+1=\beta^2\ ,\ AB+1=r^2$$

Employing the above formula (4), the fourth tuple D is given by

$$D=4(kp\pm(k+1)+ikq)\left(4(kp\pm(k+1)+ikq)^2-1\right)$$

Note that (A,B,C,D) is a gaussian diophantine quadruple with property $D(1)$.
Now, consider

$$B=kp\pm(2+k)+ikq$$

$$C=4kp\pm4(k+1)+i\,4kq$$

$$D=4(kp\pm(k+1)+ikq)\left(4(kp\pm(k+1)+ikq)^2-1\right)$$

which is a gaussian diophantine triple with property $D(1)$.
Using the above formula (4), the fourth tuple E is given by

$$\begin{aligned}E=&\,256(kp+ikq)^5\pm(1280k+1536)(kp+ikq)^4+(2560k^2+6144k+3552)(kp+ikq)\\&\pm(2560k^3+9216k^2+10656k+3936)(kp+ikq)^2\\&+(1280k^4+6144k^3+10656k^2+7872k+2081)(kp+ikq)\\&\pm(256k^5+1536k^4+3552k^3+3936k^2+2081k+420)\end{aligned}$$

Observe that (B,C,D,E) is a gaussian diophantine quadruple with property $D(1)$. Proceeding in this way, one may generate a sequence of gaussian diophantine quadruples (A,B,C,D), (B,C,D,E),.....with property $D(1)$.
Some numerical examples of gaussian diophantine quadruples for the choices of $(k,p,q)=(1,2,1),(2,1,2),(2,2,3),(3,3,2)$ are respectively given below in Table: 1.

Table1: Numerical Examples

(A,B,C,D)	(B,C,D,E)
$(3+i\ ,5+i\ ,16+4i\ ,816+748i)$,	$(5+i\ ,16+4i\ ,816+748i\ ,142023+346401i)$,
$(1+i\ ,-1+i\ ,4i\ ,-20i)$	$(-1+i\ ,4i\ ,-20i\ ,-323+289i)$
$(4+4i\ ,6+4i\ ,20+16i\ ,-1860+3760i)$,	$(6+4i\ ,20+16i\ ,-1860+3760i\ ,-3067352-459644i)$,
$(-4i\ ,-2+4i\ ,-4+16i\ ,756-848i)$	$(-2+4i\ ,-4+16i\ ,756-848i\ ,-330660+43140i)$
$(6+6i\ ,8+6i\ ,28+24i\ ,-6636+10632i)$,	$(8+6i\ ,28+24i\ ,-6636+10632i\ ,-17446806-6132282i)$,
$(2+6i\ ,6i\ ,4+24i\ ,-1716-3192i)$	$(6i\ ,4+24i\ ,-1716-3192i\ ,1291422+1667526i)$
$(12+6i\ ,14+6i\ ,52+24i\ ,12636+45192i)$,	$(14+6i\ ,52+24i\ ,12636+45192i\ ,-87594576+138411462i)$,
$(6+6i\ ,4+6i\ ,20+24i\ ,-6660+3720i)$	$(4+6i\ ,20+24i\ ,-6660+3720i\ ,-1522398-6699066i)$

2.1.2. Diophantine quadruple with property $D(4)$:

Let $n=2$. Then the triple (A,B,C) is given by

$$A=kp\pm k+ikq$$

$$B=kp\pm(4+k)+ikq$$

$$C=4kp\pm4(2+k)+i\,4kq$$

which is a gaussian diophantine triple with property $D(4)$.
If (A,B,C) is a diophantine triple with property $D(4)$ then the fourth tuple D is given by

$$D=(A+B+C)+\frac{1}{2}(ABC+\alpha\beta r) \quad (5)$$

where

$$AC+4=\alpha^2\ ,\ BC+4=\beta^2\ ,\ AB+4=r^2$$

Employing the above formula (5), the fourth tuple D is given by

$$D=4(kp\pm(k+2)+ikq)\left((kp\pm(k+2)+ikq)^2-1\right)$$

Note that (A,B,C,D) is a gaussian diophantine quadruple with property $D(4)$.
Now, consider

$$B=kp\pm(4+k)+ikq$$

$$C=4kp\pm4(2+k)+i\,4kq$$

$$D=4(kp\pm(k+2)+ikq)\left((kp\pm(k+2)+ikq)^2-1\right)$$

which is a gaussian diophantine triple with property $D(4)$.
Using the above formula (5), the fourth tuple E is given by

$$\begin{aligned}E=&\,16(kp+ikq)^5\pm(80k+192)(kp+ikq)^4+(160k^2+768k+888)(kp+ikq)^3\\&\pm(160k^3+1152k^2+2664k+1968)(kp+ikq)^2\\&+(80k^4+768k^3+2664k^2+3936k+2081)(kp+ikq)\\&\pm(16k^5+192k^4+888k^3+1968k^2+2081k+840)\end{aligned}$$

Observe that (B,C,D,E) is a gaussian diophantine quadruple with property $D(4)$. Proceeding in this way, one may generate a sequence of gaussian diophantine quadruples (A,B,C,D), (B,C,D,E), ...with property $D(4)$.
Some numerical examples of gaussian diophantine quadruples for the choices of $(k,p,q)=(1,2,1),(2,1,2),(2,2,3),(3,3,2)$ are respectively given below in Table: 2.

Table 2: Numerical Examples

(A,B,C,D)	(B,C,D,E)

$(3+i\ ,7+i\ ,20+4i\ ,420+292i)$,	$(7+i\ ,20+4i\ ,420+292i\ ,43995+60465i)$,
$(1+i\ ,-3+i\ ,-4+4i\ ,12+4i)$	$(-3+i\ ,-4+4i\ ,12+4i\ ,169-143i)$
$(4+4i\ ,8+4i\ ,24+16i\ ,-312+1456i)$,	$(8+4i\ ,24+16i\ ,-312+1456i\ ,-366644+119428i)$,
$(4i\ ,-4+4i\ ,-8+16i\ ,360-80i)$	$(-4+4i\ ,-8+16i\ ,360-80i\ ,-18504-32124i)$
$(6+6i\ ,10+6i\ ,32+24i\ ,-1440+3720i)$,	$(10+6i\ ,32+24i\ ,-1440+3720i\ ,-1998976+40134i)$,
$(2+6i\ ,-2+6i\ ,24i\ ,-888i)$	$(-2+6i\ ,24i\ ,-888i\ ,-42630+126150i)$
$(12+6i\ ,16+6i\ ,56+24i\ ,4872+13224i)$,	$(16+6i\ ,56+24i\ ,4872+13224i\ ,-5847660+13478790i)$,
$(6+6i\ ,2+6i\ ,16+24i\ ,-1488+264i)$	$(2+6i\ ,16+24i\ ,-1488+264i\ ,125694-243258i)$

2.2. Problem 2:

Let A, B be two gaussian integers represented by

$A=p+q+i\ ,\ B=p-q+i$

where p and q are non-zero integers.
Note that

$AB+q^2=(p+i)^2=r^2$ (say)

Therefore, the pair (A,B) is a gaussian diophantine 2-tuple with property $D(q^2)$
Consider C to be a gaussian integer such that

$AC+q^2=\alpha^2$ (6)

$BC+q^2=\beta^2$ (7)

Assume

$\alpha=A+r\ ,\ \beta=B+r$ (8)

Substituting (8) in (6) and (7) and subtracting one from the other, observe that

$C=A+B+2r=4(p+i)$

It is observed that the triple (A,B,C) is a gaussian diophantine 3-tuple with property $D(q^2)$. When $q=1,q=2$, the above triple (A,B,C) can be extended to diophantine quadruple with their corresponding properties.

2.2.1. Diophantine quadruple with property $D(1)$:

Let $q=1$. Then the triple (A,B,C) is given by

$A=p+1+i$

$B=p-1+i$

$C=4(p+i)$

which is a gaussian diophantine triple with property $D(1)$.
If (A,B,C) is a diophantine triple with property $D(1)$ then the fourth tuple D is given by

$D=(A+B+C)+2(ABC+\alpha\beta r)$ (9)

where

$AC+1=\alpha^2\ ,\ BC+1=\beta^2\ ,\ AB+1=r^2$

Employing the above formula (9), the fourth tuple D is given by

$D=4(p+i)\left(4(p+i)^2-1\right)$

Note that (A,B,C,D) is a gaussian diophantine quadruple with property $D(1)$.
Now, consider

$B=p-1+i$

$C=4(p+i)$

$D=4(p+i)\left(4(p+i)^2-1\right)$

which is a gaussian diophantine triple with property $D(1)$.
Using the above formula (9), the fourth tuple E is given by

$E=256(p+i)^5-256(p+i)^4-32(p+i)^3+64(p+i)^2+(p+i)-3$

Observe that (B,C,D,E) is a gaussian diophantine quadruple with property $D(1)$. Proceeding in this way, one may generate a sequence of gaussian diophantine quadruples (A,B,C,D), (B,C,D,E), ...with property $D(1)$.
Some numerical examples are given below:

Table 3: Numerical Examples

p	(A,B,C,D)	(B,C,D,E)
2	$(3+i\ ,1+i\ ,8+4i\ ,24+172i)$	$(1+i\ ,8+4i\ ,24+172i\ ,-7809+4257i)$
3	$(4+i\ ,2+i\ ,12+4i\ ,276+412i)$	$(2+i\ ,12+4i\ ,276+412i\ ,-10304+55873i)$
4	$(5+i\ ,3+i\ ,16+4i\ ,816+748i)$	$(3+i\ ,16+4i\ ,816+748i\ ,61505+224545i)$
5	$(6+i\ ,4+i\ ,20+4i\ ,1740+1180i)$	$(4+i\ ,20+4i\ ,1740+1180i\ ,372802+587073i)$

2.2.2. Diophantine quadruple with property $D(4)$:

Let $q=2$. Then the triple (A,B,C) is given by

$A=p+2+i$

$B=p-2+i$

$C=4(p+i)$

which is a gaussian diophantine triple with property $D(4)$.

If (A,B,C) is a diophantine triple with property $D(4)$ then the fourth tuple D is given by

$$D=(A+B+C)+\frac{1}{2}(ABC+\alpha\beta r) \quad (10)$$

where

$AC+4=\alpha^2$, $BC+4=\beta^2$, $AB+4=r^2$

Employing the above formula (10), the fourth tuple D is given by

$$D=4(p+i)\left((p+i)^2-1\right)$$

Note that (A,B,C,D) is a gaussian diophantine quadruple with property $D(4)$.
Now, consider

$B=p-2+i$

$C=4(p+i)$

$D=4(p+i)\left((p+i)^2-1\right)$

which is a gaussian diophantine triple with property $D(4)$.
Using the above formula (10), the fourth tuple E is given by

$$E=16(p+i)^5-32(p+i)^4-8(p+i)^3+32(p+i)^2+(p+i)-6$$

Observe that (B,C,D,E) is a gaussian diophantine quadruple with property $D(4)$. Proceeding in this way, one may generate a sequence of gaussian diophantine quadruples (A,B,C,D), (B,C,D,E),.....with property $D(4)$.
Some numerical examples are given below:

Table4: Numerical Examples

p	(A,B,C,D)	(B,C,D,E)
2	$(4+i\ ,\ i\ ,\ 8+4i\ ,\ 40i)$	$(i\ ,\ 8+4i\ ,\ 40i\ ,\ -308-71i)$
3	$(5+i\ ,\ 1+i\ ,\ 12+4i\ ,\ 60+100i)$	$(1+i\ ,\ 12+4i\ ,\ 60+100i\ ,\ -979+1969i)$
4	$(6+i\ ,\ 2+i\ ,\ 16+4i\ ,\ 192+184i)$	$(2+i\ ,\ 16+4i\ ,\ 192+184i\ ,\ 1374+10137i)$
5	$(7+i\ ,\ 3+i\ ,\ 20+4i\ ,\ 420+292i)$	$(3+i\ ,\ 20+4i\ ,\ 420+292i\ ,\ 15055+30385i)$

References

[1] L.E. Dickson, History of the theory of numbers, Chelsea Publishing Company, Newyork, Vol.2, 1966, 513-520.

[2] E. Brown, Sets in which $xy+k$ is always a square, Math. Comp., 45, (1985), 613-620.https://doi.org/10.2307/2008150.

[3] J. Morgado, Generalization of a result of Hoggatt and Bergum on Fibonacci numbers Portugaliae Math, 42, (1983-1984), 441-445.

[4] J. Morgado, note on a Shannon's theorem concerning the Fibonacci numbers and Diophantine quadruples, Portugaliae Math, 48, (1991), 429-439.

[5] G. Srividhya, Diophantine quadruples for Fibonacci numbers with property $D(1)$, Indian Journal of Mathematics and Mathematical Sciences, 5, (2009), 57-59.

[6] M.A. Gopalan, G. Srividhya, Diophantine quadruples for Fibonacci and Lucas numbers with property $D(4)$, Diophantus J. Math., 1, (2012), 15-18.

[7] A. Dujella, Generalization of a problem of Diophantus, Acta Arith, 65, (1993), 15-27.https://doi.org/10.4064/aa-65-1-15-27.

[8] A. Dujella, A problem of Diophantus and pell numbers, Application of Fibonacci Numbers, Vol.7, (G.E. Bergum, A.N. Philippou, A.F. Horadan eds.) Khowet Dordrecht, (1998), 61-68.https://doi.org/10.1007/978-94-011-5020-0_9.

[9] Y. Fujita, The extensibility of Diophantine pairs $[k-1,k+1]$, J. Number Theory, 128, (2008), 322-353.

[10] Y. Fujita, A. Togbe, Uniqueness of the extension of the $D(4k^2)$-triple $(k^2-4\ ,\ k^2\ ,\ 4k^2-4)$, Notes Number Theory, Discrete Math, 17, (2011), 42-49.

[11] Z. Cerin, on extended Euler quadruples, AttiSemin Mat Fis. Univ Modena Reggio Emilia 58, (2011), 101-120.

[12] Y. Zhang, Diophantine triples and extendibility of [1, 2, 5] and [1, 5, 10], Master Thesis, Central Michigan University, 2011.

[13] Lj. Bacic, A. Filipin, The extendibility of D(4)-pairs Math, Commun, 18, (2013), 447-456.

[14] K. Meena, S. Vidhyalakshmi, M.A. Gopalan, G. Akila, R. Presenna, Formation of special diophantine quadruples with property $D(6kpq)^2$, The international journal of Science and Technology 2, (2014) , 11-14.https://doi.org/10.9790/5728-10321214.

[15] Andrej Dujella and Florian Luca, Diophantine m-tuples for primes Intern. Math.Research, Notes 47, (2005), 2913-2940.https://doi.org/10.1155/IMRN.2005.2913.

[16] M.A. Gopalan, K. Lakshmi, S. Vidhyalakshmi, Gaussian Diophantine quadruples with property D(1), IOSR, Vol 10(3), version II, (2014), 12-14.https://doi.org/10.9790/5728-10321214.

[17] M.A. Gopalan, S. Mallika, S. Vidhyalakshmi, On the extendibility of the Gaussian 2-tuple to Gaussian Diophantine quadruple with property D(1), International Journal of Innovation in Science and Mathematics, 2(3), (2014), 310-311.https://doi.org/10.9790/5728-10321214.

[18] M.A. Gopalan, S. Mallika, S. Vidhyalakshmi, On Diophantine quadruple with property $D(P^2)$ where P is prime and $P^2\equiv 1(\mathrm{mod}\,6)$, International Journal of Current Research, 6(5), (2014), 6810-6813.

[19] K. Meena, S. Vidhyalakshmi, M.A. Gopalan, R. Presenna, Sequences of special dio-triples, IJMTT, 10(1), (2014), 43-46.https://doi.org/10.14445/22315373/IJMTT-V10P508.

[20] M.A. Gopalan, S. Mallika, S. Vidhyalakshmi, On sequences of irrational diophantine quadruples, International Journal of Mathematical Sciences and Applications, 4(1), (2014), 128-135.

[21] M.A. Gopalan, S. Vidhyalakshmi, N. Thiruniraiselvi, On strong rational diophantine quadruples with equal members, SJMPS, 1(2), (2014), 88-95.

[22] M.A. Gopalan, S. Vidhyalakshmi, N. Thiruniraiselvi, On strong diophantine quadruples with property D(1), The IJST, 2(7), (2014), 17-22.https://doi.org/10.14445/22315373/IJMTT-V20P514.

[23] M.A. Gopalan, S. Vidhyalakshmi, N. Thiruniraiselvi, "A special diophantine quadruple with property D (17)", IJMTT, 20(2), (April 2015), 110-112.https://doi.org/10.14445/22315373/IJMTT-V20P514.

[24] M.A. Gopalan, S. Vidhyalakshmi, N. Thiruniraiselvi, "Construction of irrational gaussian diophantine quadruples", IJETMR, 1(1), (April 2015), 1-7.

[25] P. Saranya and G. Janaki, "Construction of gaussian diophantine triples with the property D(25)", International Journal of Statistics and Applied Mathematics, 2(6), (2017), 301-302.