International Journal of Medicine, 14 (2) (2026) 36-44

International Journal of Medicine

Website: www.sciencepubco.com/index.php/IJM
https://doi.org/10.14419/v0qj6v63
Research paper

The Chemistry of Love: Exploring The Roles of Dopamine, Serotonin, and Oxytocin in Human Attraction And Emotional Bonding

Prof. Dr. Ravuri Hema Krishna *

Department of Chemistry, Amrita Sai Institute of Science and Technology
Bathinapadu, Andhra Pradesh, India 521180
**Corresponding author E-mail: hkravuri32@gmail.com*

Received: 21 Sep 2026. Accepted: 25 Oct 2026, Published: 8 Nov 2026

Abstract

Love is a complex human experience that involves a combination of emotional, psychological, neurological, and biochemical processes. Although it is commonly described as an emotion, scientific research demonstrates that several neurotransmitters and hormones contribute to the feelings of attraction, affection, pleasure, and attachment associated with love. Among these chemical messengers, dopamine, serotonin, and oxytocin play significant roles in influencing emotional responses and interpersonal relationships. Dopamine is primarily associated with the brain's reward and motivation systems, contributing to feelings of pleasure, excitement, and the desire to maintain meaningful interactions. Serotonin participates in mood regulation, emotional processing, and cognitive functions, potentially influencing attention and thought patterns during romantic attraction. Oxytocin, often studied in the context of social bonding, contributes to attachment, trust, and affiliative behaviours, particularly during supportive social interactions and close relationships. These substances do not operate independently; rather, they participate in interconnected neural pathways that interact with other hormones, brain regions, and environmental experiences. Their effects may vary according to individual differences, relationship stages, social circumstances, and personal experiences. Understanding the chemistry of love provides valuable insights into how biological mechanisms contribute to human emotions and social connections. This article explores the physiological functions of dopamine, serotonin, and oxytocin, examines their involvement in attraction and emotional attachment, and discusses how their interactions influence motivation, mood, attention, and interpersonal behavior. It also highlights the importance of distinguishing simplified popular explanations from the complexity of scientific evidence, recognizing that love cannot be explained by a single chemical or reaction. Overall, studying the biochemical foundations of love offers a scientific perspective on the relationship between brain chemistry, emotional experiences, and human bonding, while emphasizing the integrated nature of biological and psychological influences in shaping meaningful relationships.

Keywords: *Love; Dopamine; Serotonin; Oxytocin; Neurotransmitters; Hormones; Brain Chemistry; Attachment.*

1. Introduction

Love is one of the most complex and meaningful experiences of human life, influencing emotions, thoughts, motivation, and social behavior. It can be associated with happiness, excitement, affection, emotional security, attraction, and a strong desire to maintain close relationships. From a scientific perspective, love is not simply an emotion produced by a single part of the brain or a single chemical reaction [1]. Instead, it is a multifaceted phenomenon involving interconnected neurological, biochemical, psychological, and social processes. These processes work together to shape how individuals experience attraction, develop emotional bonds, respond to affection, and maintain interpersonal relationships [2].

The human brain plays a central role in the experience of love by coordinating several systems responsible for reward, motivation, emotional regulation, memory, attention, and social interaction. When individuals experience romantic attraction or emotional closeness, particular neural pathways may become active, influencing their feelings and behavior. Chemical messengers known as neurotransmitters and hormones help transmit signals within the nervous system and between the brain and other parts of the body. Their activity contributes to the physiological and psychological changes that can accompany attraction, affection, and attachment [3-4]. Among the chemical messengers frequently investigated in the scientific study of love are dopamine, serotonin, and oxytocin. Each has distinct biological functions, although their effects can overlap and depend on the wider neural systems in which they operate. Dopamine is involved in reward processing, motivation, learning, and the reinforcement of behaviors. Serotonin participates in mood regulation, cognition, and emotional processing. Oxytocin is associated with social bonding, affiliative behavior, and aspects of interpersonal attachment. Examining these chemicals provides a useful foundation for understanding how biological mechanisms contribute to emotional experiences and social relationships [5].

Copyright © Prof. Dr. Ravuri Hema Krishna. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Romantic love can involve changes in the way the brain processes rewards, interprets social information, and directs attention toward a particular individual. For example, interactions with a loved person may become especially rewarding, encouraging individuals to seek further contact and develop shared experiences. Emotional memories, personal expectations, communication, and environmental circumstances can also influence the development of relationships. Consequently, the experience of love differs between individuals and may change over time as relationships develop from initial attraction toward familiarity, trust, companionship, and emotional attachment [6]. It is also important to recognize that dopamine, serotonin, and oxytocin do not independently determine whether a person experiences love. Their effects depend on interactions with other neurotransmitters, hormones, brain regions, and psychological factors. Popular explanations sometimes describe love as a simple chemical reaction or attribute specific feelings entirely to one substance. However, such explanations can oversimplify the available scientific evidence. Human relationships involve considerably more than biochemical activity alone, including personal experiences, social learning, communication, cultural influences, and individual differences [7].

The scientific investigation of love therefore requires an interdisciplinary approach combining neuroscience, biochemistry, psychology, and behavioural research. Studying its biological foundations can help explain why emotional connections may influence motivation, attention, mood, and social behavior while also improving our understanding of human bonding [8-9]. The present article examines the roles of dopamine, serotonin, and oxytocin in the chemistry of love, focusing on their biological functions, their involvement in attraction and attachment, and their interactions within the nervous system. It also considers the relationship between biochemical mechanisms and psychological experiences, emphasizing that love is a complex phenomenon emerging from the coordinated activity of biological, emotional, and social processes [10-11].

2. Dopamine: The Chemistry of Reward and Motivation

Dopamine is a neurotransmitter, meaning that it helps nerve cells communicate with one another. It plays important roles in reward, motivation, learning, movement, and attention. During romantic attraction, activity in brain reward pathways can increase when a person thinks about or interacts with someone, they find important. Dopamine helps create feelings of anticipation and motivation [12]. This can partly explain why a person may become highly interested in seeing or communicating with someone they care about as shown in Figure1.




Fig. 1: Dopamine: The Chemistry of Reward, Motivation, and Romantic Attraction.

Dopamine is therefore associated with the "wanting" and motivational aspects of attraction. It is important to note that dopamine is not simply a chemical that produces happiness. Its functions are broader and include learning which experiences or cues are important and worth pursuing. Dopamine belongs to a group of compounds called catecholamines. Its molecular formula is: $C_8H_{11}NO_2$ [13]. Its chemical structure allows it to interact with specific dopamine receptors in the nervous system.

3. Serotonin: Mood and Emotional Regulation

Serotonin, chemically known as 5-hydroxytryptamine (5-HT), is another important neurotransmitter. It is involved in mood, sleep, appetite, learning, and many other physiological processes. Serotonin is sometimes popularly described as the "happiness chemical," but this is an oversimplification [14]. Its role is much more complicated. The serotonin system interacts with many regions of the brain and influences several aspects of emotional and behavioural regulation.




Fig. 2: Serotonin: The Chemistry of Mood Regulation, Emotional Processing, and Romantic Love.

Research on romantic love has found differences in serotonin-related measures in some people experiencing intense early romantic attraction. However, these findings should not be interpreted as meaning that serotonin alone causes love. Human emotions arise from interactions among many biological and psychological systems. Serotonin has the molecular formula: $C_{10}H_{12}N_2O$ [15]. It is synthesized from the amino acid tryptophan through a series of biochemical reactions as shown in Figure 2.

4. Oxytocin: A Chemical Messenger of Social Bonding

Oxytocin is a hormone and neuropeptide that has important roles in social behavior, reproduction, childbirth, and lactation. It is also involved in social bonding and interactions between individuals. Oxytocin is sometimes called the "love hormone" or "bonding hormone." These names are useful for introducing its role, but they do not describe the whole scientific picture. Oxytocin can influence social behavior depending on the person, situation, and broader neural context [16-17].




Fig. 3: Oxytocin: Its Chemical Structure, Neurotransmission, and Role in Social Bonding and Emotional Attachment.

Positive social interactions and physical affection can be associated with oxytocin release. The chemical may contribute to feelings of social connection and attachment, particularly in established relationships. Unlike dopamine and serotonin, which are commonly discussed as neurotransmitters, oxytocin is a peptide hormone and neuropeptide. It is produced in the hypothalamus and released through the posterior pituitary, while also acting within the brain as shown in Figure 3.

5. Review Methodology

This article was prepared as a narrative review examining the roles of dopamine, serotonin, and oxytocin in romantic attraction, emotional bonding, and interpersonal relationships. The review focuses on relevant literature from neuroscience, neuroendocrinology, behavioral science, psychology, and behavioral medicine. Literature identification should be documented using the databases actually searched, such as PubMed, Scopus, Web of Science, or Google Scholar, together with the search date, search terms, and any applicable publication-date or language restrictions. Relevant search concepts include "romantic love," "dopamine and reward," "serotonin and emotional regulation," "oxytocin and social bonding," "pair bonding," and "neurobiology of attachment."

Studies should be selected according to their relevance to the review objectives, methodological quality, and contribution to understanding the biological and psychological processes associated with attraction and attachment. Human neuroimaging studies, experimental investigations, observational studies, animal research relevant to bonding mechanisms, and authoritative reviews may be considered, while their different evidentiary contributions should be distinguished during synthesis. Studies should not be described as having been systematically identified or screened unless a documented search and selection process was performed. The findings are synthesized narratively, with attention to established mechanisms, emerging evidence, conflicting findings, methodological limitations, and implications for human relationships and behavioral medicine [18].

6. How these Chemicals Work Together

Love cannot be explained by one molecule. Instead, several biological systems interact. A simplified model can be described as follows: Attraction → Reward and motivation → Emotional interaction → Bonding and attachment.




Fig. 4: Synergistic Roles of Dopamine, Serotonin, and Oxytocin in the Neurobiology of Love, Attraction, and Emotional Bonding.

Dopamine can contribute to motivation and reward associated with attraction. Serotonin participates in mood and emotional regulation. Oxytocin contributes to social bonding and attachment. Other biological systems are also involved, including brain regions associated with reward, emotion, memory, and social cognition. Additional chemicals, including norepinephrine and endogenous opioids, can also contribute to aspects of attraction, stress, pleasure, and attachment. Therefore, describing love as simply "dopamine + serotonin + oxytocin" would be scientifically incomplete, as shown in Figure 4. Table. 1 compares dopamine, serotonin, and oxytocin based on their main biological functions and their involvement in love. It highlights how dopamine contributes to reward and motivation, serotonin participates in mood and emotional regulation, and oxytocin supports social bonding and attachment. Together, these chemical messengers contribute to the complex biological processes underlying attraction, affection, and emotional relationships.

Table 1: Dopamine, Serotonin and Oxytocin Compared

Chemical	Main biological role	Connection to love
Dopamine	Reward, motivation, learning and movement	Can contribute to attraction, motivation and reward
Serotonin	Mood, sleep, appetite and other functions	Participates in emotional regulation and has been studied in romantic attraction
Oxytocin	Social bonding, reproduction and other physiological functions	Can contribute to bonding and social attachment

7. Critical Evaluation of Neurochemical Evidence in Romantic Love

Although dopamine, serotonin, and oxytocin are frequently discussed in relation to romantic attraction and emotional bonding, their contributions should not be interpreted as simple or independent causal mechanisms. Dopamine has well-established roles in motivation, reward learning, and incentive salience. However, evidence that a person experiences romantic attraction does not establish that dopamine

alone causes the associated emotional experience. Human neuroimaging findings identify activity in brain regions associated with reward and motivation, but these findings do not directly measure the release or concentration of a specific neurotransmitter. The evidence concerning serotonin and romantic attraction requires similar caution. Studies of serotonin-related measures have reported associations with aspects of early romantic attraction, but peripheral measures, such as platelet serotonin transporter measures, cannot be assumed to represent serotonin activity throughout the human brain. Differences between studies and the complexity of serotonin signalling limit attempts to assign a single emotional function to this neurotransmitter.

Oxytocin has been investigated extensively in relation to social recognition, affiliation, attachment, and stress-related processes. Nevertheless, its behavioral effects are context-dependent and may vary with individual characteristics, social circumstances, and the methods used to measure or administer it. Findings from animal models can provide valuable mechanistic insights but cannot automatically be generalized to human romantic relationships. Likewise, peripheral oxytocin measurements do not necessarily provide a direct indication of oxytocin activity in specific brain circuits. Taken together, these limitations indicate that romantic love is better understood as an emergent phenomenon involving interconnected neural networks, neurochemical signalling, memory, cognition, individual psychology, and social experience. Future research should combine appropriately designed longitudinal, experimental, neuroimaging, and behavioral studies to clarify how these systems interact over time [19].

8. Chemistry Behind the Feeling of Love

The experience of love involves a complex interaction between biological, neurological, psychological, and social processes. It can influence attention, motivation, emotional responses, memory, and behavior. For example, a person may frequently think about someone important to them, feel excitement before meeting them, or experience comfort and emotional security during positive social interactions. These experiences are not produced by one chemical alone; they emerge from the coordinated activity of multiple systems in the brain and body [20].

8.1. Neurotransmitters

Neurotransmitters are chemical messengers that allow nerve cells to communicate. Dopamine participates in reward, motivation, learning, and attention, while serotonin contributes to mood regulation and emotional processing. Other neurotransmitters, including norepinephrine, can influence alertness and physiological arousal. Their combined activity helps explain some of the changes in attention, excitement, and motivation that may accompany romantic attraction [21].

8.2. Hormones and neuropeptides

Hormones and neuropeptides help regulate bodily functions and can also influence behavior and social interactions. Oxytocin, for instance, is associated with social bonding and attachment, while vasopressin has been studied in relation to social behavior and partner bonding. Their effects depend on the situation, individual differences, and interactions with other biological systems [22].

8.3. Brain reward pathways

Brain reward pathways help individuals learn which experiences are meaningful and motivating. During attraction, interactions or reminders involving a significant person may become especially noticeable or rewarding. Dopamine-related signalling contributes to these processes by influencing motivation, reward learning, and the anticipation of meaningful experiences. Reward activity, however, does not explain every aspect of love [23].

8.4. Memory systems

Memory systems help individuals remember important people, shared experiences, conversations, and emotionally meaningful events. The hippocampus and other brain regions involved in memory contribute to the formation and retrieval of these experiences. Positive memories can strengthen feelings of familiarity and connection, while past experiences may influence how individuals interpret and respond to relationships [24].

8.5. Emotional processing

Brain regions involved in emotional processing help individuals recognize, interpret, and respond to emotional information. The amygdala, prefrontal cortex, and other interconnected regions participate in evaluating social situations, regulating emotional responses, and making decisions. These processes can influence how people experience affection, excitement, uncertainty, and emotional security [25].

8.6. Social experiences

Love develops within a social environment. Communication, mutual understanding, emotional support, trust, shared activities, and positive interactions can contribute to the development and maintenance of close relationships. Social experiences may also influence the activity of biological systems associated with stress regulation, reward, and attachment [26].

8.7. Individual psychology

Individual psychological characteristics also shape the experience of love. Personality, expectations, previous experiences, emotional learning, cultural background, and personal values can influence how individuals understand attraction and develop relationships. Consequently, the same social interaction may produce different emotional responses in different people [27].

8.8. Integration of biological and psychological processes

These systems interact continuously rather than operating as separate mechanisms. Neurotransmitters influence neural communication, hormones and neuropeptides contribute to physiological and behavioural regulation, and brain networks integrate information from memories, emotions, and social experiences [28]. The resulting experience depends on the combined activity of these systems and the individual's circumstances. A simplified conceptual model of love is shown in Figure 5.




Fig. 5: A Simplified Conceptual Model of Love: from Attraction and Social Interaction to Reward Processing, Emotional Experiences, and the Development of Bonding and Attachment.

The chemistry behind love demonstrates how biological mechanisms contribute to complex human experiences. Dopamine, serotonin, oxytocin, and other chemical messengers participate in neural systems associated with reward, mood, motivation, and social bonding. However, their actions are influenced by memory, emotional processing, social experiences, and individual psychology. Therefore, love cannot be accurately described as a simple combination of three chemicals. It is better understood as a dynamic experience emerging from the interaction of biological processes with psychological and social factors. This integrated perspective provides a more scientifically meaningful explanation of how humans develop attraction, affection, and lasting emotional connections [29].

9. Love is Really A Chemical Reaction

Love is not merely a chemical reaction. Chemicals are essential for the brain and body to experience emotions, but love also involves memories, experiences, thoughts, relationships, culture, personality, and social circumstances. A useful way to understand this is: Chemistry provides part of the biological machinery; the human experience emerges from the interaction of that machinery with thoughts, experiences, and relationships. Thus, studying the chemistry of love does not reduce love to a single molecule. Instead, it helps scientists understand how the brain and body participate in one of the most important aspects of human social life [30].

10. Consequences of Excessive Neurochemical Activity

Dopamine, serotonin, and oxytocin are essential chemical messengers involved in brain function, emotional regulation, motivation, and social bonding. However, excessive or abnormal activity in these systems, particularly when associated with medications or other substances, can produce adverse effects. The consequences differ depending on the chemical involved, the amount and duration of exposure, individual susceptibility, and the surrounding biological conditions [31].

10.1. Dopamine

Abnormal dopamine signalling has been associated with psychotic symptoms, including hallucinations, delusions, and disturbances in the interpretation of information. However, dopamine activity is regulated by complex brain circuits, and excessive dopamine signalling does not automatically cause schizophrenia. Schizophrenia is a multifactorial disorder involving genetic, environmental, developmental, and neurobiological influences [32-33].

10.2. Serotonin

Excessive serotonin activity caused by certain medications, drug interactions, or overdoses can lead to a potentially serious condition called serotonin syndrome. Symptoms may include agitation, confusion, sweating, tremors, increased heart rate, and muscle rigidity. Severe cases can involve high fever, seizures, and loss of consciousness, requiring emergency medical treatment.

10.3. Oxytocin

Oxytocin is involved in social bonding and several physiological processes. When administered medically, particularly during childbirth, excessive doses can cause dangerous uterine contractions, abnormal heart rhythms, and water intoxication. Severe complications can include seizures and loss of consciousness. These risks concern excessive medical administration and should not be interpreted as evidence that ordinary social affection or naturally occurring oxytocin release is dangerous. The Consequences of Excessive Neurochemical Activity: Dopamine Dysregulation, Serotonin Syndrome, and Adverse Effects of Excessive Oxytocin Administration, as shown in Figure 6.




Fig. 6: Consequences of Excessive Neurochemical Activity: Dopamine Dysregulation, Serotonin Syndrome, and Adverse Effects of Excessive Oxytocin Administration.

11. Strengths and Limitations of The Available Evidence

The evidence discussed in this review can be considered in four broad categories. First, established neurobiological research supports the involvement of dopamine in reward learning and motivation, serotonin in multiple aspects of neural and behavioral regulation, and oxytocin in reproductive physiology and aspects of social behavior. These general functions are better established than any claim that a particular chemical independently produces romantic love.

Second, human neuroimaging and behavioral studies provide evidence of associations between romantic experiences and activity in brain networks involved in motivation, reward, attention, and social cognition. Such associations are informative but do not, by themselves, establish the causal contribution of a particular neurotransmitter. Third, animal studies provide important evidence concerning the neural and hormonal mechanisms of pair bonding and social attachment. However, species differences and experimental conditions must be considered when applying these findings to human relationships. Fourth, conceptual models that integrate neurochemistry with memory, personal experience, and social interaction are useful for education and hypothesis generation. They should be identified as explanatory frameworks rather than experimentally validated representations of every individual's experience of love [34]. Throughout the manuscript, these distinctions will guide the interpretation of findings and prevent established physiological functions, preliminary associations, and theoretical explanations from being presented as equivalent forms of evidence.

12. Clinical and Behavioral Medicine Implications

The neurobiology of attraction and emotional bonding is relevant to behavioural medicine because reward processing, emotional regulation, stress responses, memory, and social interaction are important components of psychological well-being. Research on these systems may contribute to a broader understanding of how interpersonal experiences influence motivation, emotional responses, and social functioning. Alterations in reward processing and emotional regulation have been investigated across several psychiatric and neurological conditions. However, these processes are not specific to romantic relationships, and the involvement of dopamine or serotonin in a disorder should not be interpreted as evidence that the disorder is caused by a single chemical imbalance. Mental health conditions generally involve multiple interacting genetic, developmental, environmental, psychological, and neurobiological factors.

Research on oxytocin and social bonding may also inform the study of affiliation, social cognition, and interpersonal stress. Nevertheless, the relationship between oxytocin and attachment is complex, and current findings do not justify treating oxytocin as a universal indicator of trust or relationship quality. Similarly, attachment patterns should not be treated as psychiatric diagnoses in themselves. From a clinical perspective, the most useful contribution of neurochemical research is to provide one part of a broader biopsychosocial understanding of human behavior [35]. The present review does not propose a diagnostic test or treatment based on measuring dopamine, serotonin, or oxytocin. Further translational research is needed before specific findings can be converted into reliable clinical applications.

13. Conclusions

The chemistry of love demonstrates the extraordinary complexity of the human brain. Dopamine contributes to reward and motivation, serotonin participates in mood and emotional regulation, and oxytocin plays an important role in social bonding and attachment. These chemicals interact with numerous other biological systems rather than acting alone. Love therefore represents a fascinating connection between chemistry, neuroscience, psychology, and human relationships. Studying these processes can help us understand how biological mechanisms contribute to attraction, affection, attachment, and social connection. Romantic love cannot be represented by a single chemical formula or a simple additive combination of neurotransmitters. Instead, it emerges from the dynamic interaction of neurochemical signalling, neural networks, emotional processing, memory, motivation, and social experiences. The roles of dopamine, serotonin, and oxytocin should therefore be understood as interconnected contributions to a complex biopsychosocial phenomenon rather than as independent or sufficient explanations of love. The effects of excessive neurochemical activity depend on the specific substance, its biological context, and the circumstances involved. It is scientifically inaccurate to claim that overdosing on dopamine, serotonin, or oxytocin

invariably causes schizophrenia, extreme paranoia, or insanity. A more accurate explanation distinguishes abnormal neurotransmitter signalling from medically documented drug toxicity and recognizes that mental health conditions generally arise through multiple interacting factors.

Acknowledgements

The author is deeply grateful to Almighty God and my parents for the wisdom, grace, and strength to complete this manuscript. Special thanks are extended to Dr. M. Sasidhar- Principal, Dr. K. Sai Manoj- CEO, Sri K. Rama Mohana Rao- Secretary and Correspondent, Sri K. Lakshmi Karthik- President, and Sri K. Ramesh Babu- Industrialist and Chairman of Amrita Sai Institute of Science and Technology, whose Candor, patience, understanding, and constant encouragement have been a source of inspiration throughout this challenging journey of writing the manuscript. The author also gratefully acknowledges the support and cooperation of all the members of the S&H, CRT departments.

Disclosure Statement

No potential conflict of interest was reported by the authors.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or non-profit sectors.

Authors' Contributions

The authors have contributed to data analysis, drafting, and revising of the article and agreed to be responsible for all aspects of this work.

References

[1] Bartels A, Zeki S. The neural basis of romantic love. Neuroreport. 2000;11(17):3829–3834. https://doi.org/10.1097/00001756-200011270-00046.
[2] Fisher HE, Aron A, Brown LL. Romantic love: a mammalian brain system for mate choice. Philosophical Transactions of the Royal Society B: Biological Sciences. 2006;361(1476):2173–2186. https://doi.org/10.1098/rstb.2006.1938.
[3] Aron A, Fisher H, Mashek DJ, Strong G, Li H, Brown LL. Reward, motivation, and emotion systems associated with early-stage intense romantic love. Journal of Neurophysiology. 2005;94(1):327–337. https://doi.org/10.1152/jn.00838.2004.
[4] Fisher H, Aron A, Brown LL. Romantic love: an fMRI study of a neural mechanism for mate choice. Journal of Comparative Neurology. 2005;493(1):58–62. https://doi.org/10.1002/cne.20772.
[5] de Boer A, van Buel EM, ter Horst GJ. Love is more than just a kiss: a neurobiological perspective on love and affection. Neuroscience. 2012; 201:114–124. https://doi.org/10.1016/j.neuroscience.2011.11.017.
[6] Marazziti D, Akiskal HS, Rossi A, Cassano GB. Alteration of the platelet serotonin transporter in romantic love. Psychological Medicine. 1999;29(3):741–745. https://doi.org/10.1017/S0033291798007946.
[7] Young LJ, Wang Z. The neurobiology of pair bonding. Nature Neuroscience. 2004; 7:1048–1054. https://doi.org/10.1038/nn1327.
[8] Ross HE, Young LJ. Oxytocin and the neural mechanisms regulating social cognition and affiliative behavior. Frontiers in Neuroendocrinology. 2009;30(4):534–547. https://doi.org/10.1016/j.yfrne.2009.05.004.
[9] Olff M, Frijling JL, Kubzansky LD, Bradley B, Ellenbogen MA, Cardoso C, et al. The role of oxytocin in social bonding, stress regulation and mental health: an update on the moderating effects of context and interindividual differences. Psychoneuroendocrinology. 2013;38(9):1883–1894. https://doi.org/10.1016/j.psyneuen.2013.06.019.
[10] Bosch OJ, Young LJ. Oxytocin and social relationships: from attachment to bond disruption. Current Topics in Behavioral Neurosciences. 2018; 35:97–117. https://doi.org/10.1007/7854_2017_10.
[11] Szymanska M, Schneider M, Chateau-Smith C, et al. Psychophysiological effects of oxytocin on parent–child interactions: a literature review on oxytocin and parent–child interactions. Psychiatry and Clinical Neurosciences. 2017;71(10):690–705. https://doi.org/10.1111/pcn.12544.
[12] Cong X, et al. Oxytocin and early parent-infant interactions: a systematic review. International Journal of Nursing Sciences. 2019;6(4):445–453. https://doi.org/10.1016/j.ijnss.2019.09.009.
[13] Campbell A. Attachment, aggression and affiliation: the role of oxytocin in female social behavior. Biological Psychology. 2008;77(1):1–10. https://doi.org/10.1016/j.biopsycho.2007.09.001.
[14] Froemke RC, Young LJ. Oxytocin, neural plasticity, and social behavior. Annual Review of Neuroscience. 2021; 44: 359–381. https://doi.org/10.1146/annurev-neuro-102320-102847.
[15] Loth MK, Donaldson ZR. Oxytocin, dopamine, and opioid interactions underlying pair bonding: highlighting a potential role for microglia. Endocrinology. 2021;162(2): bqaa223. https://doi.org/10.1210/endocr/bqaa223.
[16] Bromberg-Martin ES, Matsumoto M, Hikosaka O. Dopamine in motivational control: rewarding, aversive, and alerting. Neuron. 2010;68(5):815–834. https://doi.org/10.1016/j.neuron.2010.11.022.
[17] Wise RA. Dopamine, learning and motivation. Nature Reviews Neuroscience. 2004; 5: 483–494. https://doi.org/10.1038/nrn1406.
[18] Ikemoto S, Yang C, Tan A. Basal ganglia circuit loops, dopamine and motivation: a review and enquiry. Behavioural Brain Research. 2015; 290: 17–31. https://doi.org/10.1016/j.bbr.2015.04.018.
[19] Rappeneau V, Castillo Díaz F. Convergence of oxytocin and dopamine signalling in neuronal circuits: insights into the neurobiology of social interactions across species. Neuroscience & Biobehavioral Reviews. 2024; 161: 105675. https://doi.org/10.1016/j.neubiorev.2024.105675.
[20] Palacios-Ariza MA, Pradilla IF, Acosta-Quevedo DC. Neurobiología del amor romántico. Revista Médica Sanitas. 2021;24(4):197–209.
[21] Marazziti D, Palermo S, Mucci F. The science of love: state of the art. Advances in Experimental Medicine and Biology. 2021; 1331: 249–254. https://doi.org/10.1007/978-3-030-74046-7_16.
[22] Stein DJ, Vythilingum B. Love and attachment: the psychobiology of social bonding. CNS Spectrums. 2009;14(5):239–242. https://doi.org/10.1017/S1092852900025384.
[23] Donaldson ZR, Nautiyal KM, Ahmari SE, Hen R. Genetic approaches for understanding the role of serotonin receptors in mood and behavior. Current Opinion in Neurobiology. 2013;23(3):399–406. https://doi.org/10.1016/j.conb.2013.01.011.
[24] Berendzen KM, Manoli DS. Rethinking the architecture of attachment: new insights into the role for oxytocin signaling. Affective Science. 2022; 3: 734–748. https://doi.org/10.1007/s42761-022-00142-5.

[25] Weinstein AM. Reward, motivation and brain imaging in human healthy participants – a narrative review. Frontiers in Behavioral Neuroscience. 2023; 17: 1123733. https://doi.org/10.3389/fnbeh.2023.1123733.
[26] Berridge KC, Robinson TE. What is the role of dopamine in reward: hedonic impact, reward learning, or incentive salience? Brain Research Reviews. 1998;28(3):309–369. https://doi.org/10.1016/S0165-0173(98)00019-8.
[27] Schultz W, Dayan P, Montague PR. A neural substrate of prediction and reward. Science. 1997; 275(5306): 1593–1599. https://doi.org/10.1126/science.275.5306.1593.
[28] Love TM. The brain and romantic love. Journal of Neuroscience, Psychology, and Economics. 2015; 8(3): 201–208. https://doi.org/10.1037/npe0000055.
[29] Blumenthal SA, Young LJ. The neurobiology of love and pair bonding from human and animal perspectives. Biology (Basel). 2023 Jun 12;12(6):844. https://doi.org/10.3390/biology12060844.
[30] The neurobiological basis of love: a meta-analysis of human functional neuroimaging studies of maternal and passionate love. Brain Sciences. 2022;12(7):830. https://doi.org/10.3390/brainsci12070830.
[31] Toda M, Abi-Dargham A. Dopamine hypothesis of schizophrenia: making sense of it all. Current Psychiatry Reports. 2007;9(4):329–336. https://doi.org/10.1007/s11920-007-0041-7.
[32] Howes OD, Kapur S. The dopamine hypothesis of schizophrenia: version III the final common pathway. Schizophrenia Bulletin. 2009;35(3):549–562. https://doi.org/10.1093/schbul/sbp006.
[33] Scotton WJ, Hill LJ, Williams AC, Barnes NM. Serotonin syndrome: pathophysiology, clinical features, management, and potential future directions. International Journal of Tryptophan Research. 2019; 12: 1178646919873925. https://doi.org/10.1177/1178646919873925.
[34] Mikkelsen N, Damkier P, Pedersen SA. Serotonin syndrome a focused review. Basic & Clinical Pharmacology & Toxicology. 2023; 133(2): 124–129. https://doi.org/10.1111/bcpt.13912.
[35] Chiew AL, Isbister GK. Management of serotonin syndrome (toxicity). British Journal of Clinical Pharmacology. 2025. https://doi.org/10.1111/bcp.16152.