International Journal of Physical Research, 10 (1) (2022) 39-44

International Journal of Physical Research

Website: www.sciencepubco.com/index.php/IJPR

Research paper



Modified Kudryashov method and its applications to the (2+1)-dimensional cubic Klein-Gordon and (3+1)-dimensional Zakharov-Kuznetsov equations

K. M. Abdul Al Woadud [1] *, Md. Jahirul Islam [1], Aminur Rahman Khan [2]

[1] *Department of Electrical & Electronic Engineering, Uttara University, Dhaka-1230, Bangladesh*
[2] *Department of Mathematics, Jahangirnagar University, Savar, Dhaka-1342,Bangladesh*
**Corresponding author E-mail: woadudru50@gmail.com*

Abstract

The exact solutions of nonlinear evolution equations (NLEEs) play a vital role to reveal the central mechanism of complex physical phenomenon. More precisely, in this paper, we acquired new exact solutions to the (2+1)-dimensional cubic Klein–Gordon (cKG) and (3+1)-dimensional Zakharov–Kuznetsov (ZK) equations by using the modified Kudraysov method. As results, a portion of the new accurate voyaging wave answers for the situations above is officially delivered. All arrangements are plotted in the perspective on three-dimensional (3D) and two-dimensional (2D) line shape through the MATLAB programming for exploring the genuine meaning of the concentrated on conditions. The periodic type of solution is created using a modified Kudryashov approach, which is distinct from the other methods investigated.

Keywords: (2+1)-Dimensional Cubic Klein-Gordon Equation; (3+1)-Dimensional Zakharov–Kuznetsov Equation; Modified Kudryashov Method; Traveling Wave Solutions; Symbolic Computation

1. Introduction

Nonlinear phenomena are found in many domains of science and engineering, including fluid mechanics, plasma physics, optical fibers, biology, solid state physics, chemical kinematics, chemical physics, and so on. It is well acknowledged that these complex physical processes can be described using NLEEs. So, study to find exact solutions of NLEEs is very important. A wide collection of researchers is eager to discover effective methods for discovering nonlinear equations analytic and numerical solutions. To find traveling wave solutions to nonlinear scientific phenomena like, the Hirota's bilinear transformation method [1], [2], the tanh-function method [3], the (G'/G)-expansion method [4-12], the Exp-function method [13-16], the homogeneous balance method [17], [18], the F-expansion method [19], the Adomian decomposition method [20], the homotopy perturbation method [21], the extended tanh-method [22], the auxiliary equation method [23], the Jacobi elliptic function method [24], Weierstrass elliptic function method [25], modified Exp-function method [26], the modified simple equation method [27-30], the extended multiple Riccati equations expansion method [31-34] they found many powerful and efficient methods and techniques.
Recently, utilizing computational software packages such as Maple, Mathematica, and MATLAB, researchers were able to readily build and apply these strategies for minimizing computational difficulties. New traveling wave solutions are being investigated, Firstly, we consider the (2+1) dimensional cubic Klein-Gordon equation is given by

$$\psi_{xx} + \psi_{yy} - \psi_{tt} + \alpha\psi + \beta\psi^3 = 0, \tag{1}$$

Where $\psi = \psi(x, y, t)$ and α, β are non zero constants. It's a specific form of nonlinear evolution equation that can be found in a wide range of applications.
Secondly, we consider the (3+1)-dimensional Zakharov–Kuznetsov (ZK) equation is given by

$$\psi_t + a\psi\psi_x + \psi_{xx} + \psi_{yy} + \psi_{zz} = 0, \tag{2}$$

Where $\psi = \psi(x, y, z, t)$.
The major goal of this study is to use an efficient method known as the modified Kudryashov method to generate new accurate traveling wave solutions of the (2+1)-dimensional cubic Klein-Gordon and the (3+1)-dimensional Zakharov–Kuznetsov (ZK) equations [10], [35], [36].

Copyright © K. M. Abdul Al Woadud et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The remainder of this document is organized as follows: The modified Kudryashov method approach is described in Sections 2 and 3. In Section 3, we will use the methods described above to obtain novel accurate traveling wave solutions for the (2+1) dimensional cubic Klein-Gordon equation and the (3+1) dimensional Zakharov–Kuznetsov (ZK) equation. Finally, section 4 concludes with a statement summarizing the findings.

2. Outline of the modified kudryashov method

The modified Kudryashov method is a reliable problem-solving methodology that has gotten a lot of attention in the search for new exact solutions to nonlinear differential equations in mathematical physics. To begin, we will offer a quick overview of the modified Kudryashov method approach [10], which is used to find new exact solutions to a nonlinear partial differential equation. We use a non-linear partial differential equation to achieve this goal

$$F(u, u_x, u_t, u_{xx}, u_{xt}, u_{tt}, \dots) = 0, \tag{3}$$

Where the function $u = u(x,t)$ is unknown and F is a polynomial. The main steps are as follows:
Step 1: Introducing the transformation $u(x,t) = U(\xi)$ where $\xi = kx + \omega t$, varies according to Eq. (3). This reduces to the following non-linear ordinary differential equation

$$P(U, U', U'', \dots) = 0, \tag{4}$$

Where P is a polynomial of U and its derivatives, and the superscripts denote the ordinary derivatives with respect to ξ.
Step 2: Let us assume that the solution $U(\xi)$ of the nonlinear Eq. (4) can be written as

$$U(\xi) = A_0 + \sum_{i=1}^{N} A_i Q^i(\xi), A_N \neq 0, \tag{5}$$

Where A_i $(i = 0,1 \dots N)$ are later determined, N is a positive integer that may be obtained using the balancing principle on Eq. (5), and $Q(\xi)$ satisfies the following new equation

$$Q'(\xi) = \left(Q^2(\xi) - Q(\xi)\right)\ln(A), \tag{6}$$

With the exact solution $Q(\xi) = \frac{1}{1+dA^{\xi}}$, where $A \neq 0,1$.
Step 3: We get a system of algebraic equations in parameters A_i $(i = 0,1 \dots N)$, k, and ω by inserting Eq. (5) and Eq. (6) into Eq. (4) and performing basic mathematical operations. By setting the obtained values in Eq. (5), finally generates new exact solutions for the Eq. (3).

3. Mathematical analysis

3.1. The (2+1)-dimensional cubic Klein-Gordon (cKG) equation

By considering the following transformation:

$$\psi(x,y,t) = U(\xi), \xi = x + y - \lambda t \tag{7}$$

Using Eq. (7) into the (2+1) dimensional cubic Klein-Gordon equation (1), which can be reduced to a nonlinear ordinary differential equation as shown below

$$(2 - \lambda^2)U'' + \alpha U + \beta U^3 = 0 \tag{8}$$

N=1 is found using the homogeneous balancing principle. The answer to Eq. (8) then takes the form

$$U(\xi) = A_0 + A_1 Q(\xi) \tag{9}$$

By substituting Eq. (9) along with its second derivative into Eq. (8), and equating the like powers of $Q(\xi)$, then we get in the following system of equations:

$$-2\,(\lambda^2 - 2)\ln(A)^2 A_1 + \beta A_1^3 = 0,$$

$$3\,(\lambda^2 - 2)\,\ln(A)^2 A_1 + 3\beta A_0 A_1^2 = 0,$$

$$-(\lambda^2 - 2)\,\ln(A)^2 A_1 + \alpha A_1 + 3\beta A_0^2 A_1 = 0,$$

$$\beta A_0^3 + \alpha A_0 = 0.$$

By solving the above system, we receive the following different solution sets:

Set–I: $\lambda = \frac{\sqrt{2\ln(A)^2 - 2\alpha}}{\ln(A)}$, $A_0 = \frac{\sqrt{-\alpha\beta}}{\beta}$, and $A_1 = \frac{2\alpha}{\sqrt{-\alpha\beta}}$.

Set-I possesses the following exact solution of the (2+1) dimensional cubic Klein-Gordon equations extracted:

$$\psi_1(x,y,t) = \frac{\sqrt{-\alpha\beta}}{\beta} + \frac{2\alpha}{\sqrt{-\alpha\beta}\left(1+dA^{-\frac{\sqrt{2\ln(A)^2-2\alpha}\,t}{2\ln(A)}+x+y}\right)} \tag{10}$$

Set–II: $\lambda = \frac{\sqrt{2\ln(A)^2-2\alpha}}{\ln(A)}, A_0 = -\frac{\sqrt{-\alpha\beta}}{\beta}$, and $A_1 = -\frac{2\alpha}{\sqrt{-\alpha\beta}}$.

Set-II possesses to the following exact solution of the (2+1) dimensional cubic Klein-Gordon equations determined:

$$\psi_2(x,y,t) = -\frac{\sqrt{-\alpha\beta}}{\beta} - \frac{2\alpha}{\sqrt{-\alpha\beta}\left(1+dA^{-\frac{\sqrt{2\ln(A)^2-2\alpha}\,t}{2\ln(A)}+x+y}\right)} \tag{11}$$

Set–III: $\lambda = -\frac{\sqrt{2\ln(A)^2-2\alpha}}{\ln(A)}, A_0 = \frac{\sqrt{-\alpha\beta}}{\beta}$, and $A_1 = \frac{2\alpha}{\sqrt{-\alpha\beta}}$.

Set-II possesses to the following exact solution of the (2+1) dimensional cubic Klein-Gordon equations determined:

$$\psi_3(x,y,t) = \frac{\sqrt{-\alpha\beta}}{\beta} + \frac{2\alpha}{\sqrt{-\alpha\beta}\left(1+dA^{\frac{\sqrt{2\ln(A)^2-2\alpha}\,t}{2\ln(A)}+x+y}\right)} \tag{12}$$

Set–IV: $\lambda = -\frac{\sqrt{2\ln(A)^2-2\alpha}}{\ln(A)}, A_0 = -\frac{\sqrt{-\alpha\beta}}{\beta}$, and $A_1 = -\frac{2\alpha}{\sqrt{-\alpha\beta}}$.

Set-II possesses to the following exact solution of the (2+1) dimensional cubic Klein-Gordon equations determined:

$$\psi_4(x,y,t) = -\frac{\sqrt{-\alpha\beta}}{\beta} - \frac{2\alpha}{\sqrt{-\alpha\beta}\left(1+dA^{\frac{\sqrt{2\ln(A)^2-2\alpha}\,t}{2\ln(A)}+x+y}\right)} \tag{13}$$

Fig.1, Fig.2, Fig.3, and Fig.4 show the three-dimensional (3D) and two-dimensional (2D) plots of Eq. (10), Eq. (11), Eq. (12), and Eq. (13) correspondingly. The plots of Eqs. (10), (11), (12), and (13) appear to have a periodic characteristic that is distinct from others.




Fig. 1: (a)-(b)-(c) 3D illustrations of the Eq. (10) for the choosing arbitrary parametr $A = -1, \alpha = 5, \beta = 2, y = 0, d = 2$ within $-10 \le x \le 10$, $-10 \le t \le 10$, and (d)-(e)-(f) 2D line illustrations of (a)-(b)-(c) at $t = 1$, respectively.




Fig. 2: (a)-(b)-(c) 3D illustrations of the Eq. (11) for the choosing arbitrary parameters $A = -1, \alpha = 5, \beta = 2, y = 0, d = 2$ within $-10 \le x \le 10$, $-10 \le t \le 10$, and (d)-(e)-(f) 2D line illustrations of (a)-(b)-(c) at $t = 1$, respectively.




Fig. 3: (a)-(b)-(c) 3D illustrations of the Eq. (12) for the choosing arbitrary parameters $A = -1, \alpha = 5, \beta = 2, y = 0, d = 2$ within $-10 \le x \le 10$, $-10 \le t \le 10$, and (d)-(e)-(f) 2D line illustrations of (a)-(b)-(c) at $t = 1$, respectively.




Fig. 4: (a)-(b)-(c) 3D illustrations of the Eq. (13) for the choosing arbitrary parameters $A = -1, \alpha = 5, \beta = 2, y = 0, d = 2$ within $-10 \le x \le 10, -10 \le t \le 10$, and (d)-(e)-(f) 2D line illustrations of (a)-(b)-(c) at $t = 1$, respectively.

3.2. The (3+1)-dimensional Zakharov –Kuznetsov (ZK) equation

By introducing the following transformation:

$$\psi(x, y, z, t) = U(\xi), \xi = x + y + z - \lambda t \tag{14}$$

By employing the transformation Eq. (14) into the (3+1)-dimensional Zakharov–Kuznetsov (ZK) equation Eq. (2) can be converted to a nonlinear ordinary differential equation as follows

$$-\lambda u + \frac{1}{2} a u^2 + 3u' = 0 \tag{15}$$

N=1 is found using the homogeneous balancing principle. Then, the solution of Eq. (15) takes the form

$$U(\xi) = A_0 + A_1 Q(\xi) \tag{16}$$

By substituting Eq. (16) along with its first derivative into Eq. (15), and equating the like powers of (ξ) , then we get in the following system of equations:

$$\frac{1}{2} A_1 \left(a\, A_1 + 6\, ln(A)\right) = 0,$$

$$A_1 \left(a\, A_0 - \lambda - 3\, ln(A)\right) = 0,$$

$$\frac{1}{2} A_0 (a\, A_0 - 2\lambda) = 0.$$

By solving the above system, we receive the following different solution sets:

Set–I: $\lambda = -3ln(A), A_0 = 0, A_1 = -\frac{6\, ln(A)}{a}$

Set-I corresponds the following exact solution of the (3+1)-dimensional Zakharov–Kuznetsov (ZK) equations derived:

$$\psi_1(\mathrm{x}, \mathrm{y}, \mathrm{z}, \mathrm{t}) = \frac{6 \ln(\mathrm{A})}{\mathrm{a}\left(1 + \mathrm{dA}^{3 \ln(\mathrm{A})\mathrm{t} + \mathrm{x} + \mathrm{y} + \mathrm{z}}\right)} \tag{17}$$

Set-II: $\lambda = 3\ln(\mathrm{A}), \mathrm{A}_0 = \frac{6 \ln(\mathrm{A})}{\mathrm{a}}, \mathrm{A}_1 = -\frac{6 \ln(\mathrm{A})}{\mathrm{a}}$

Set-II corresponds the following exact solution of of the (3+1)-dimensional Zakharov–Kuznetsov (ZK) equations explored:

$$\psi_2(x,y,z,t) = \frac{6\ln(A)}{a} - \frac{6\ln(A)}{a\left(1+dA^{-3\ln(A)t+x+y+z}\right)} \tag{18}$$

In Fig.5 and Fig.6, the three-dimensional (3D) and two-dimensional (2D) plots of Eq. (17) and Eq. (18) are shown, respectively. The plots appear to have a periodic activity that is distinct from others.




Fig. 5: (a)-(b)-(c) 3D illustrations of the Eq. (18) for the choosing arbitrary parameters $A = -18, a = -7, y = 0, z = 0, d = 8$ within $-10 \le x \le 10, -10 \le t \le 10$, and (d)-(e)-(f) 2D line illustrations of (a)-(b)-(c) at $y = 0, z = 0, t = 1$, respectively.




Fig. 6: (a)-(b)-(c) 3D illustrations of the Eq. (17) for the choosing arbitrary parameters $A = -18, a = -7, y = 0, z = 0, d = 8$ within $-10 \le x \le 10, -10 \le t \le 10$, and (d)-(e)-(f) 2D Line Illustrations of (a)-(b)-(c) at $y = 0, z = 0, t = 1$, respectively.

4. Conclusions

The modified Kudraysov method is used to solve the (2+1)-D cKG and (3+1)-D ZK equations in this study as a robust strategy. A number of new accurate traveling wave solutions for the nonlinear equations are explicitly derived as a result of this work. It should be noted that the validity of the results presented in this study is tested by re-entering each new solution into its original equation.

Acknowledgment

The authors would like to express their gratitude to Md. Moiful Alam and Mohammed Khurshed Alam of Uttara University's Department of Electrical and Electronic Engineering for their helpful ideas in the preparation of the manuscript.

References

[1] Hirota, R. (1973). Exact envelope-soliton solutions of a nonlinear wave equation. *Journal of Mathematical Physics*, *14*(7), 805-809. https://doi.org/10.1063/1.1666399.

[2] Hirota, R., & Satsuma, J. (1981). Soliton solutions of a coupled Korteweg-de Vries equation. *Physics Letters A*, *85*(8-9), 407-408. https://doi.org/10.1016/0375-9601(81)90423-0.

[3] Abdel-All, N. H., Abdel-Razek, M. A. A., & Seddeek, A. A. K. (2011). Expanding the tanh-function method for solving nonlinear equations. *Applied Mathematics*, *2*(09), 1096. https://doi.org/10.4236/am.2011.29151.

[4] Wang, M., Li, X., & Zhang, J. (2008). The (G′ G)-expansion method and travelling wave solutions of nonlinear evolution equations in mathematical physics. *Physics Letters A*, *372*(4), 417-423. https://doi.org/10.1016/j.physleta.2007.07.051.

[5] Zayed, E. M. E. (2010). TRAVELING WAVE SOLUTIONS FOR HIGHER DIMENSIONAL NONLINEAR EVOLUTION EQUATIONS USING THE $(\frac {G'} {G}) $-EXPANSION METHOD. *Journal of Applied Mathematics & Informatics*, *28*(1_2), 383-395.

[6] Zayed, E. M. E., & Gepreel, K. A. (2009). The (G′/G)-expansion method for finding traveling wave solutions of nonlinear partial differential equations in mathematical physics. *Journal of Mathematical Physics*, *50*(1), 013502. https://doi.org/10.1063/1.3033750.

[7] Akbar, M. A., Ali, N. H. M., & Zayed, E. M. E. (2012). Abundant exact traveling wave solutions of generalized bretherton equation via improved (G′/G)-expansion method. *Communications in Theoretical Physics*, *57*(2), 173. https://doi.org/10.1088/0253-6102/57/2/01.

[8] Akbar, M. A., Ali, N. H., & Zayed, E. M. E. (2012). A generalized and improved-expansion method for nonlinear evolution equations. *Mathematical Problems in Engineering*, *2012*. https://doi.org/10.1155/2012/459879.

[9] Akbar, M. A., Ali, N. H. M., & Mohyud-Din, S. T. (2012). The alternative (G′/G)-expansion method with generalized Riccati equation: Application to fifth order (1+1)-dimensional Caudrey-Dodd-Gibbon equation. *Int J Phys Sci*, *7*(5), 743-752. https://doi.org/10.5897/IJPS11.1669.

[10] Joardar, A. K., Kumar, D., & Al Woadud, K. A. (2018). New exact solutions of the combined and double combined sinh-cosh-Gordon equations via modified Kudryashov method. *Int. J. Phys. Res*, *1*, 25-30.Akbar, M. A., & Ali, N. H. M. (2011). The alternative (G′/G)-expansion method and its applications to nonlinear partial differential equations. *Int J Phys Sci*, *6*(35), 7910-7920. https://doi.org/10.14419/ijpr.v6i1.9261.

[11] Shehata, A. R. (2010). The traveling wave solutions of the perturbed nonlinear Schrödinger equation and the cubic–quintic Ginzburg Landau equation using the modified (G′/G)-expansion method. *Applied Mathematics and Computation*, *217*(1), 1-10. https://doi.org/10.1016/j.amc.2010.03.047.

[12] He, J. H., & Wu, X. H. (2006). Exp-function method for nonlinear wave equations. *Chaos, Solitons & Fractals*, *30*(3), 700-708. https://doi.org/10.1016/j.chaos.2006.03.020.

[13] Akbar, M. A., & Ali, N. H. M. (2011). Exp-function method for Duffing Equation and new solutions of (2+1) dimensional dispersive long wave equations. *Progress in Applied Mathematics*, *1*(2), 30-42.

[14] Naher, H., Abdullah, F. A., & Akbar, M. A. (2011). The Exp-function method for new exact solutions of the nonlinear partial differential equations. *International Journal of Physical Sciences*, *6*(29), 6706-6716. https://doi.org/10.5897/IJPS11.1026.

[15] Naher, H., Abdullah, F. A., & Akbar, M. A. (2012). New traveling wave solutions of the higher dimensional nonlinear partial differential equation by the Exp-function method. *Journal of Applied Mathematics*, *2012*. https://doi.org/10.1155/2012/575387.

[16] Wang, M. (1995). Solitary wave solutions for variant Boussinesq equations. *Physics letters A*, *199*(3-4), 169-172. https://doi.org/10.1016/0375-9601(95)00092-H.

[17] Zayed, E. M. E., Zedan, H. A., & Gepreel, K. A. (2004). On the solitary wave solutions for nonlinear Hirota–Satsuma coupled KdV of equations. *Chaos, Solitons & Fractals*, *22*(2), 285-303. https://doi.org/10.1016/j.chaos.2003.12.045.

[18] Zhou, Y., Wang, M., & Wang, Y. (2003). Periodic wave solutions to a coupled KdV equations with variable coefficients. *Physics Letters A*, *308*(1), 31-36. https://doi.org/10.1016/S0375-9601(02)01775-9.

[19] Adomian, G. (1994). Solving Frontier Problems Decomposition Method. https://doi.org/10.1007/978-94-015-8289-6.

[20] Mohyud-Din, S. T., & Noor, M. A. (2007). Homotopy perturbation method for solving fourth-order boundary value problems. *Mathematical Problems in Engineering*, *2007*. https://doi.org/10.1155/2007/98602.

[21] Abdou, M. A. (2007). The extended tanh method and its applications for solving nonlinear physical models. *Applied mathematics and computation*, *190*(1), 988-996. https://doi.org/10.1016/j.amc.2007.01.070.

[22] Sirendaoreji. (2004). New exact travelling wave solutions for the Kawahara and modified Kawahara equations. *Chaos Solitons & Fractals*, *19*(1), 147-150. https://doi.org/10.1016/S0960-0779(03)00102-4.

[23] Ali, A. T. (2011). New generalized Jacobi elliptic function rational expansion method. *Journal of computational and applied mathematics*, *235*(14), 4117-4127. https://doi.org/10.1016/j.cam.2011.03.002.

[24] Shi, L. M., Zhang, L. F., Meng, H., Zhao, H. W., & Zhou, S. P. (2011). A method to construct Weierstrass elliptic function solution for nonlinear equations. *International Journal of Modern Physics B*, *25*(14), 1931-1939. https://doi.org/10.1142/S0217979211100436.

[25] He, Y., Li, S., & Long, Y. (2012). Exact solutions of the Klein-Gordon equation by modified Exp-function method. In *Int. Math. Forum* (Vol. 7, No. 4, pp. 175-182).

[26] Jawad, A. J. A. M., Petković, M. D., & Biswas, A. (2010). Modified simple equation method for nonlinear evolution equations. *Applied Mathematics and Computation*, *217*(2), 869-877. https://doi.org/10.1016/j.amc.2010.06.030.

[27] Zayed, E. M. (2011). A note on the modified simple equation method applied to Sharma–Tasso–Olver equation. *Applied Mathematics and Computation*, *218*(7), 3962-3964. https://doi.org/10.1016/j.amc.2011.09.025.

[28] Zayed, E. M. E., & Ibrahim, S. H. (2012). Exact solutions of nonlinear evolution equations in mathematical physics using the modified simple equation method. *Chinese Physics Letters*, *29*(6), 060201. https://doi.org/10.1088/0256-307X/29/6/060201.

[29] Zayed, E. M. E., & Arnous, A. H. (2012, September). Exact solutions of the nonlinear ZK-MEW and the potential YTSF equations using the modified simple equation method. In *AIP Conference Proceedings* (Vol. 1479, No. 1, pp. 2044-2048). AIP. https://doi.org/10.1063/1.4756591.

[30] Gepreel, K. A., & Shehata, A. R. (2012). Exact complexiton soliton solutions for nonlinear partial differential equations in mathematical physics. *Scientific Research and Essays*, *7*(2), 149-157.

[31] Gepreel, K. A., & Shehata, A. R. (2012). Exact complexiton soliton solutions for nonlinear partial differential equations in mathematical physics. *Scientific Research and Essays*, *7*(2), 149-157.

[32] Zayed, E. M. E., & Gepreel, K. A. (2011). A series of complexiton soliton solutions for nonlinear Jaulent—Miodek PDEs using the Riccati equations method. *Proceedings of the Royal Society of Edinburgh Section A: Mathematics*, *141*(5), 1001-1015. https://doi.org/10.1017/S0308210510000405.

[33] Gepreel, K. A. (2011). A Generalized (G G)-expansion method to find the traveling wave solutions of nonlinear evolution equations. *Journal of Partial Differential Equations*, *24*(1), 55-69. https://doi.org/10.4208/jpde.v24.n1.4.

[34] Gepreel, K. A. (2011). Exact solutions for nonlinear PDEs with the variable coefficients in mathematical physics. *Journal of Information and Computing Science*, *6*(1), 003-014.

[35] Kudryashov, N. A. (2012). One method for finding exact solutions of nonlinear differential equations. *Communications in Nonlinear Science and Numerical Simulation*, *17*(6), 2248-2253. https://doi.org/10.1016/j.cnsns.2011.10.016.

[36] Al Woadud, K. A., Kumar, D., Islam, M. J., Kayes, M. I., & Kundu, A. K. (2019). Extraction of Solitary Wave Features to the Heisenberg Ferromagnetic Spin Chain and the Complex Klein–Gordon Equations. *International Journal of Applied and Computational Mathematics*, *5*(3), 57. https://doi.org/10.1007/s40819-019-0642-y.