

Multiple Mellin Convolution and I-function Transformation involving r variables

Arif M. Khan

Head & Asso. Professor, JIET Jodhpur, Raj. India – 342008
 Email: khanarif76@gmail.com

Abstract

An expression is obtained for an I-function of r variables, transform of the Multiple Mellin convolution of the two functions in terms of the Multiple Mellin convolution of I-function transforms of the function involving r variables, it generalize the result given earlier by Kumbhat and Khan [5].

Keywords: Convolution , Multivariable I-Function, Multiple Mellin Convolution, Mellin Transformation.

1 Introduction

Kumbhat and Khan [5] have obtained an interesting relation involving a convolution of the Mellin type in connection with the I-function transformation. The aim of the present paper is to generalize their result to r variables, I-function transformation.

Definition 1.1 : The multivariable I-function represented by Prasad [7] as

$$I[z_1, \dots, z_r] = I_{\left\{ \begin{matrix} \{0, n_i\}_{2,r} : \{ (m^{(i)}, n^{(i)}) \}_{1,r} \\ \{p_i, q_i\}_{2,r} : \{ (p^{(i)}, q^{(i)}) \}_{1,r} \end{matrix} \right\}} \left[\begin{matrix} z_1 \\ \vdots \\ z_r \end{matrix} \middle| \begin{matrix} A : B \\ C : D \end{matrix} \right]$$

$$= \frac{1}{(2\pi\omega)^r} \int_{L_1} \dots \int_{L_r} \psi(\zeta_1, \dots, \zeta_r) \prod_{i=1}^r \{ \phi_i(\zeta_i) z_i^{\zeta_i} \} d\zeta_1 \dots d\zeta_r \quad (1.1)$$

where $\omega = \sqrt{-1}$,

$$\psi(\zeta_1, \dots, \zeta_r) = \frac{\prod_{k=2}^r \left[\prod_{j=1}^{n_k} \Gamma(1 - a_{kj} + \sum_{i=1}^k \alpha_{kj}^{(i)} \zeta_i) \right]}{\prod_{k=2}^r \left[\prod_{j=n_k+1}^{p_k} \Gamma(a_{kj} - \sum_{i=1}^k \alpha_{kj}^{(i)} \zeta_i) \right]}$$

$$\times \frac{1}{\prod_{k=2}^r \left[\prod_{j=1}^{q_k} \Gamma(1 - b_{kj} + \sum_{i=1}^k \beta_{kj}^{(i)} \zeta_i) \right]} \quad (1.2)$$

$$\phi_i(\zeta_i) = \frac{\left[\prod_{k=1}^{m^{(i)}} \Gamma(b_k^{(i)} - \beta_k^{(i)} \zeta_i) \right] \left[\prod_{j=1}^{n^{(i)}} \Gamma(1 - a_j^{(i)} + \alpha_j^{(i)} \zeta_i) \right]}{\left[\prod_{j=n^{(i)}+1}^{p^{(i)}} \Gamma(a_j^{(i)} - \alpha_j^{(i)} \zeta_i) \right] \left[\prod_{k=m^{(i)}+1}^{q^{(i)}} \Gamma(1 - b_k^{(i)} + \beta_k^{(i)} \zeta_i) \right]} \quad (1.3)$$

$\forall i \in \{1, \dots, r\}$. Also,

$$\begin{aligned} \{0, n_i\}_{2,r} &:= 0, n_2 : \dots : 0, n_r, \\ \{p_i, q_i\}_{2,r} &:= p_2, q_2 : \dots : p_r, q_r, \\ \{(m^{(i)}, n^{(i)})\}_{1,r}^{l,r} &:= (m^{(1)}, n^{(1)}) ; \dots ; (m^{(r)}, n^{(r)}), \\ \{(p^{(i)}, q^{(i)})\}_{1,r}^{l,r} &:= (p^{(1)}, q^{(1)}) ; \dots ; (p^{(r)}, q^{(r)}), \\ \\ A &:= \left\{ (a_{ij}; \alpha_{ij}^{(1)}, \dots, \alpha_{ij}^{(i)})_{1,p_i}^{2,r} \right\} &:= (a_{2j}; \alpha_{2j}^{(1)}, \alpha_{2j}^{(2)})_{1,p_2} ; \dots ; (a_{rj}; \alpha_{rj}^{(1)}, \alpha_{rj}^{(r)})_{1,p_r} \\ B &:= \left\{ (a_j^{(i)}; \alpha_j^{(i)})_{1,p^{(i)}}^{l,r} \right\} &:= (a_j^{(1)}, \alpha_j^{(1)})_{1,p^{(1)}} ; \dots ; (a_j^{(r)}, \alpha_j^{(r)})_{1,p^{(r)}} \\ C &:= \left\{ (b_{ij}; \beta_{ij}^{(1)}, \dots, \beta_{ij}^{(i)})_{1,q_i}^{2,r} \right\} &:= (b_{2j}; \beta_{2j}^{(1)}, \beta_{2j}^{(2)})_{1,q_2} ; \dots ; (b_{rj}; \beta_{rj}^{(1)}, \beta_{rj}^{(r)})_{1,q_r} \\ D &:= \left\{ (b_j^{(i)}; \beta_j^{(i)})_{1,q^{(i)}}^{l,r} \right\} &:= (b_j^{(1)}, \beta_j^{(1)})_{1,q^{(1)}} ; \dots ; (b_j^{(r)}, \beta_j^{(r)})_{1,q^{(r)}} \end{aligned} \quad (1.4)$$

such that $n_i, p_i, q_i, m^{(i)}, n^{(i)}, p^{(i)}, q^{(i)}$ are non-negative integers and all $a_{ij}, b_{ij}, \alpha_{ij}, \beta_{ij}, a_j^{(i)}, b_j^{(i)}, \alpha_j^{(i)}, \beta_j^{(i)}$ are complex numbers and the empty product denotes unity.

The contour integral (1.1) converges, if

$$|\arg z_i| < \frac{1}{2} U_i \pi, U_i > 0, i = 1, \dots, r \quad (1.5)$$

where

$$\begin{aligned} U_i &= \sum_{j=1}^{n^{(i)}} \alpha_j^{(i)} - \sum_{j=n^{(i)}+1}^{p^{(i)}} \alpha_j^{(i)} + \sum_{j=1}^{m^{(i)}} \beta_j^{(i)} - \sum_{j=m^{(i)}+1}^{q^{(i)}} \beta_j^{(i)} + \left(\sum_{j=1}^{n_2} \alpha_{2j}^{(i)} - \sum_{j=n_2+1}^{p_2} \alpha_{2j}^{(i)} \right) \\ &+ \dots + \left(\sum_{j=1}^{n_r} \alpha_{rj}^{(i)} - \sum_{j=n_r+1}^{p_r} \alpha_{rj}^{(i)} \right) - \left(\sum_{j=1}^{q_2} \beta_{2j}^{(i)} + \dots + \sum_{j=1}^{q_r} \beta_{rj}^{(i)} \right) \end{aligned} \quad (1.6)$$

and $I[z_1, \dots, z_r] = O(|z_1|^{\alpha_1}, \dots, |z_r|^{\alpha_r}), \max \{|z_1|, \dots, |z_r|\} \rightarrow 0,$

where $\alpha_i = \min_{1 \leq j \leq m^{(i)}} \Re(b_j^{(i)} / \beta_j^{(i)})$, and $\beta_i = \max_{1 \leq j \leq n^{(i)}} \Re(a_j^{(i)} - 1 / \alpha_j^{(i)}), i = 1, \dots, r.$

For the condition of convergence and analyticity of multivariable I-function we refer [7, 8]

Definition 1.2 : Generalized Mellin convolution can be taken in the form, given by Datta & Arora [1]

$$(k * f)(y_1, y_2, \dots, y_r) = \int_0^\infty \dots \int_0^\infty \prod_{i=1}^r x_i^{-1} k\left(\frac{y_1}{x_1}, \dots, \frac{y_r}{x_r}\right) f(x_1, \dots, x_r) dx_1 dx_2 \dots dx_r \quad (1.7)$$

Further the Mellin convolution of two I functions involving r-variables given by Datta & Arora [1] as

Theorem : Suppose the condition (9) to be satisfied and $\mu_i > 0, \nu_i > 0, n_i = 0, n_i' = 0, i \in \{1, \dots, r\}$. Then the Multiple Mellin transform of the product of two I-function of r variable.

$$\begin{aligned} (\Re I \cdot I)(s) &= \int_0^\infty \dots \int_0^\infty \prod_{i=1}^r t_i^{s_i-1} I[z_1 t_1^{-1}, \dots, z_r t_r^{-1}] I[\eta_1 t_1, \dots, \eta_r t_r] dt_1 \dots dt_r \\ &= \prod_{i=1}^r \left\{ \eta_i^{-s_i} \right\} \left\{ (0, 0)_{2,r} : \left(\left(m^{(i)} + m'^{(i)}, n^{(i)} + n'^{(i)} \right) \right)_{1,r}^{l,r} \right\} \left[\begin{matrix} z_1 \eta_1^{s_1} \\ M \\ z_r \eta_r^{s_r} \end{matrix} \middle| \begin{matrix} A : B \\ C : D \end{matrix} \right] \end{aligned} \quad (1.8)$$

where

$$\begin{aligned} & - \min_{1 \leq j \leq n^{(i)}} \Re \left(\frac{1 - a_j^{(i)}}{\alpha_j^{(i)}} \right) - \min_{1 \leq j \leq m'^{(i)}} \Re \left(\frac{b_j^{(i)}}{\beta_j^{(i)}} \right) < \Re(s_i) < \\ & \min_{1 \leq j \leq m^{(i)}} \Re \left(\frac{b_j^{(i)}}{\beta_j^{(i)}} \right) + \min_{1 \leq j \leq n'^{(i)}} \Re \left(\frac{1 - a_j^{(i)}}{\alpha_j^{(i)}} \right) \end{aligned}$$

$i = 1, \dots, r$ and

$$\begin{aligned} A &\equiv \left\{ \left(a_{ij}; \alpha_{ij}^{(1)}, \dots, \alpha_{ij}^{(i)} \right)_{1,p_i}^{2,r}; \left(a'_{ij} + \sum_{k=1}^i s_k \alpha_{ij}^{(k)} \alpha_{ij}^{(1)}, \dots, \alpha_{ij}^{(i)} \right)_{1,p_i}^{2,r} \right\}, \\ B &\equiv \left\{ \left(a_j^{(i)}, \alpha_j^{(i)} \right)_{1,p^{(i)}}^{l,r}; \left(a_j^{(i)} + s_i \alpha_j^{(i)}, \alpha_j^{(i)} \right)_{1,p^{(i)}}^{l,r} \right\} \\ C &\equiv \left\{ \left(b_{ij}; \beta_{ij}^{(1)}, \dots, \beta_{ij}^{(i)} \right)_{1,q_i}^{2,r}; \left(b_{ij} + \sum_{k=1}^i s_k \beta_{ij}^{(k)} \beta_{ij}^{(1)}, \dots, \beta_{ij}^{(i)} \right)_{1,q_i}^{2,r} \right\}, \\ D &\equiv \left\{ \left(b_j^{(i)}, \beta_j^{(i)} \right)_{1,q^{(i)}}^{l,r}; \left(b_j^{(i)} + s_i \beta_j^{(i)}, \beta_j^{(i)} \right)_{1,q^{(i)}}^{l,r} \right\} \end{aligned}$$

2 The I-Function Transformation Involving R-Variables

The I-function Transformation of function f involving r-variables is given as

$$\hat{f}(y_1, y_2, \dots, y_r) = I[f(x_1, x_2, \dots, x_r); y_1, y_2, \dots, y_r]$$

=

$$\int_0^\infty \dots \int_0^\infty \prod_{i=1}^r x_i^{-1} I[y_1 x_1, y_2 x_2, \dots, y_r x_r] f(x_1, \dots, x_r) dx_1 \dots dx_r$$

provided that the integral exists. (2.1)

3 The Convolution Property

$$\left. \begin{array}{l} \text{Let } \hat{k}(s) = I[k(x_1, \dots, x_r); s_1, \dots, s_r] \\ \text{and } \hat{f}(s) = I[f(x_1, \dots, x_r); s_1, \dots, s_r] \end{array} \right\} \quad (3.1)$$

Now we obtain an interesting result which expresses the Mellin convolution of I function transforms of two functions of r variables as the I function transform of multiple Mellin convolution of the functions.

$$\begin{aligned} (\hat{k} * \hat{f})(y_1, y_2, \dots, y_r) &= \int_0^\infty \dots \int_0^\infty \prod_{i=1}^r z_i^{-1} I[y_1 z_1, y_2 z_2, \dots, y_r z_r] \\ &\quad \cdot \left\{ \int_0^\infty \dots \int_0^\infty \prod_{i=1}^r x_i^{-1} k\left(\frac{z_1}{x_1}, \dots, \frac{z_r}{x_r}\right) f(x_1, \dots, x_r) dx_1 \dots dx_r \right\} \cdot dz_1 \dots dz_r \end{aligned} \quad (3.2)$$

provided that each side exists.

Proof : The Multiple Mellin transform of $\phi(x_1 \dots x_r)$ is given as follows

$$M\{\phi(x_1, x_2, \dots, x_r); s_i\} = \int_0^\infty \dots \int_0^\infty \prod_{i=1}^r x_i^{s_i-1} \phi(x_1, \dots, x_r) dx_1 \dots dx_r \quad (3.3)$$

with the assumptions of absolute convergence of the integrals exist and

$$\text{let } K(s) = M\{k(x_1 \dots x_r) : s_1 \dots s_r\}, F(s) = M\{f(x_1 \dots x_r) : s_1 \dots s_r\} \quad (3.4)$$

Take Mellin transform of the left hand side of (3.2)

$$\begin{aligned} \Delta(s) &= M\{(\hat{k} * \hat{f}); s_1 \dots s_r\} \\ &= \\ \int_0^\infty \dots \int_0^\infty \prod_{i=1}^r y_i^{s_i-1} &\left\{ \int_0^\infty \dots \int_0^\infty \prod_{i=1}^r x_i^{-1} k\left(\frac{y_1}{x_1}, \dots, \frac{y_r}{x_r}\right) f(x_1, \dots, x_r) dx_1 \dots dx_r \right\} dy_1 \dots dy_r \end{aligned} \quad (3.5)$$

On suitable changing the order of two sets of integrations on the right hand side of (3.5) we get

$$\begin{aligned}\Delta(s) &= M\left\{\left(\hat{k} * \hat{f}\right): s_1 \dots s_r\right\} \\ &= \int_0^\infty \dots \int_0^\infty \prod_{i=1}^r x_i^{-1} f(x_1, \dots, x_r) \left\{ \int_0^\infty \dots \int_0^\infty \prod_{i=1}^r y_i^{s_i-1} k\left(\frac{y_1}{x_1} \dots \frac{y_r}{x_r}\right) dy_1 \dots dy_r \right\} dx_1 \dots dx_r \\ &= M\{f(x_1 \dots x_r): s_1 \dots s_r\} \quad M\{k(y_1 \dots y_r): s_1 \dots s_r\} \quad (3.6)\end{aligned}$$

$$= F(s) \cdot K(s) \quad (3.7)$$

Now using the following result,

$$\begin{aligned}M\left\{\hat{k}(y_1 \dots y_r): s_1 \dots s_r\right\} &= M\left\{I(y_1 \dots y_r): s_1 \dots s_r\right\} K(-s) \\ M\left\{\hat{f}(y_1 \dots y_r): s_1 \dots s_r\right\} &= M\left\{I(y_1 \dots y_r): s_1 \dots s_r\right\} F(-s) \text{ in (3.7)}\end{aligned}$$

which can easily proved as

$$\begin{aligned}M\left\{\hat{k}(y_1 \dots y_r)\right\} &= \int_0^\infty \dots \int_0^\infty \prod_{i=1}^r y_i^{s_i-1} \left\{ \int_0^\infty \dots \int_0^\infty \prod_{i=1}^r x_i^{-1} I(y_1 x_1, \dots, y_r x_r) k(x_1, \dots, x_r) dx_1 \dots dx_r \right\} dy_1 \dots dy_r \\ &= \int_0^\infty \dots \int_0^\infty \prod_{i=1}^r x_i^{-1} k(x_1, \dots, x_r) dx_1 \dots dx_r \int_0^\infty \dots \int_0^\infty \prod_{i=1}^r y_i^{s_i-1} I(y_1 x_1, \dots, y_r x_r) dy_1 \dots dy_r\end{aligned}$$

$$\begin{aligned}\text{Let } \prod_{i=1}^r y_i x_i &= \prod_{i=1}^r z_i \\ \prod_{i=1}^r dy_i &= \frac{\prod_{i=1}^r dz_i}{\prod_{i=1}^r x_i} \\ &= \int_0^\infty \dots \int_0^\infty \prod_{i=1}^r x_i^{-s_i-1} k(x_1, \dots, x_r) dx_1 \dots dx_r \int_0^\infty \dots \int_0^\infty \prod_{i=1}^r z_i^{s_i-1} I(z_1, \dots, z_r) dz_1 \dots dz_r \\ M\left\{\hat{k}(y_1 \dots y_r)\right\} &= M\left\{I(y_1 \dots y_r)\right\} K(-s) \quad (3.8)\end{aligned}$$

Similarly

$$M\left\{\hat{f}(y_1 \dots y_r)\right\} = M\left\{I(y_1 \dots y_r)\right\} F(-s) \quad (3.9)$$

Hence from (3.7) we get

$$\Delta(s) = M\{\hat{k} * \hat{f}(y_1 \dots y_r)\} = M\{\hat{k}(y_1 \dots y_r)\} M\{\hat{f}(y_1 \dots y_r)\}$$

By using (3.8) & (3.9) we get

$$\Delta(s) = M\{I_{p,q;r}^{m,n}(y_1 \dots y_r)\} K(-s) \cdot M\{I_{p,q;r}^{m',n'}(y_1 \dots y_r)\} F(-s) \quad (3.10)$$

On the other hand Mellin convolution of the right hand side of (3.2) is

$$\begin{aligned} \Delta'(s) &= M\left\{\int_0^\infty \dots \int_0^\infty \prod_{i=1}^r z_i^{-1} = I_{p+p',q+q'}^{m+m',n+n'}[y_1 z_1, \dots, y_r z_r] \cdot (k * f)(z) dz\right\} \\ \Delta'(s) &= \int_0^\infty \dots \int_0^\infty \prod_{i=1}^r y_i^{s_i-1} \left\{ \int_0^\infty \dots \int_0^\infty \prod_{i=1}^r z_i^{-1} I[y_1 z_1, \dots, y_r z_r] (k * f)(z) dz_1 \dots dz_r \right\} dy_1 \dots dy_r \end{aligned} \quad (3.11)$$

Changing the order of two sets of integrations in (3.11), we get

$$\Delta'(s) = M\{I(y_1 z_1 \dots y_r z_r)\} \int_0^\infty \dots \int_0^\infty \prod_{i=1}^r z_i^{-s_i-1} (k * f)(z) dz_1 \dots dz_r$$

Now applying (1.8) and (3.7)

$$\Delta'(s) = M\{I_{p,q;r}^{m,n}[\xi] s_1 \dots s_r\} \cdot M\{I_{p,q;r}^{m',n'}[\eta] s'_1, s'_2 \dots s'_r\} K(-s) F(-s) \quad (3.12)$$

Equation (3.10) and (3.12) together exhibit the fact that the Mellin transforms of the two sides of (3.2) are equal, and so the two sides of (3.2) must also be equal.

References

- [1] Dutta B.K., Arora L.K., Multiple and Laplace Transforms of I-function of r variables, Journal of Fractal Calculus and Applications, Vol-1 (2011), 1-8.
- [2] Erdelyi A., et al., Table of integral transforms, Vols. I and II, Mc Graw-Hill, New York, London and Toronto, 1954.
- [3] Gupta, K.C. and Goyal, S.P., On the two dimensional Mellin convolution, Indian Journal of Mathematics, Vol. 21, No. 3(1979), 177-186.
- [4] Gupta, K.C. and Jain, U.C., The H-function II, Proc. Nat. Acad. Sci. India Sect. A 36(1966), 594-609.

- [5] Kumbhat, R.K. and Khan, A.M., Mellin convolution and I-function transformation, Proc. Int. Conf. SSFA, Vol. II, 2001, pp. 93-98.
- [6] Prasad T., Singh N.P., Mellin and Laplace transform of the multivariable I-function, Ganita Sandesh, 4(2) (1990), 66-70.
- [7] Prasad Y.N., Multivariable I-function, Vijanana Parishad Anusandhan Patrika, 29(4) (1986), 231-235.
- [8] Prasad Y.N., Yadav G.S., Proc. math. Soc. B.H.U., I (1985), 127-136.
- [9] Sharma, C.K. and Shrivastava, S., Some expansion formula for the I-function, Nat, Acad, Sci. India, 62 A (1992), 236-239.
- [10] Shrivastava, H.M. and Buschman, R.G., Mellin convolution and H-function transformations, Rocky Mountain J. Math. 6(1976), 331-343.
- [11] Saxena R.K., Singh Y., On the derivatives of the multivariable I-function, Vijananan Parishad Anusandhan Patrika, 36(2) (1993), 93-38.
- [12] Vaishya G.D., Jain R. and Varma R.C., Certain Properties of the I-function, Proc. Nat., Acad. Sci. India. 59(A) II. 1989.