

A modified Adomian decomposition method for singular initial value Emden-Fowler type equations

J. Biazar¹, K. Hosseini^{1, 2, 3*}

¹ Department of Mathematics, Rasht Branch, Islamic Azad University, Rasht, Iran

² Department of Mathematics, Guilan Science and Research Branch, Islamic Azad University, Rasht, Iran

³ Department of Applied Mathematics, Ahrar Institute of Technology and Higher Education, Rasht, Iran

*Corresponding author E-mail: kamyar_hosseini@yahoo.com

Abstract

Traditional Adomian decomposition method (ADM) usually fails to solve singular initial value problems of Emden-Fowler type. To overcome this shortcoming, a new and effective modification of ADM that only requires calculation of the first Adomian polynomial is formally proposed in the present paper. Three singular initial value problems of Emden-Fowler type with $\alpha = 1, 2$, and > 2 , and have been selected to demonstrate the efficiency of the method.

Keywords: Singular Initial Value Problems; Emden-Fowler Type; Adomian Decomposition Method; New Modification.

1. Introduction

Singular initial value problems play a fundamental role in a wide range of scientific disciplines. In this article, a special type of singular initial value problems which can be expressed as the following form is investigated.

$$u'' + \frac{\alpha}{x} u' + g(x)h(u) = k(x), \quad u(0) = a, \quad u'(0) = b.$$

Singularity behaviour that occurs at $x = 0$ is the main difficulty of this type of initial value problems. In recent years, a variety of methods have been adopted to handle this type of initial value problems. For example, Wazwaz employed a general approach for constructing the exact and series solution of this problem by means of the variational iteration method [1]. Parand et al. applied an approximation algorithm for the solution of this problem using Hermite functions, as basis functions, and collocation method [2]. Parand et al. also adopted a pseudospectral technique based on the rational Legendre functions and Gauss-Radau integration to handle this problem [3]. For further methods, the reader is referred to the references [4-19]. In the present article, ADM is modified effectively to solve presented singular initial value problem. The rest of this article is arranged as follows:

In Section 2, the basic ideas of the method are expressed with details. In Section 3, the proposed method is implemented to solve three singular initial value problems of Emden-Fowler type with $\alpha = 1, 2$, and > 2 . Finally, Section 4 is devoted to presenting conclusion.

2. Modified ADM

Let's consider the following nonlinear equation

$$u'' + R(u) + N(u) = f(x), \quad (1)$$

with the following initial conditions

$$u(0) = \xi_1, \quad u'(0) = \xi_2,$$

where R is a linear operator, N is a nonlinear operator, and $f(x)$ is a known function. It is assumed that the unknown function $u(x)$ can be presented by an infinite series, say

$$u(x) = \sum_{n=0}^{+\infty} u_n(x), \quad (2)$$

and the nonlinear term $N(u)$ can be expressed as an infinite series of polynomials given by

$$N(u) = \sum_{n=0}^{+\infty} A_n(u_0, u_1, \dots, u_n), \quad (3)$$

where $A_n, n = 0, 1, \dots$ are called the Adomian polynomials and are defined by [20]

$$A_n = \begin{cases} N(u_0) & n = 0, \\ \frac{1}{n} \sum_{i=0}^{n-1} (i+1) u_{i+1} \frac{dA_{n-1-i}}{du_0}, & n = 1, 2, \dots \end{cases}$$

Applying the inverse operator $L^{-1} = \int_0^x \int_0^x (\cdot) dx dx$, on both sides of Eq. (1) and considering (2) and (3), leads to

$$\begin{aligned} \sum_{n=0}^{+\infty} u_n &= u(0) + u'(0)x + L^{-1}[f(x)] \\ &- L^{-1}[R(\sum_{n=0}^{+\infty} u_n) + \sum_{n=0}^{+\infty} A_n(u_0, u_1, \dots, u_n)]. \end{aligned} \quad (4)$$

We rewrite (4) as follows [21], [22]

$$\sum_{n=0}^{+\infty} u_n = u(0) + u'(0)x$$

$$+L^{-1}[\sum_{n=0}^{+\infty} a_n x^n] - pL^{-1}[\sum_{n=0}^{+\infty} a_n x^n] + L^{-1}[f(x)]$$

$$-L^{-1}[R(\sum_{n=0}^{+\infty} u_n) + \sum_{n=0}^{+\infty} A_n(u_0, u_1, \dots, u_n)],$$

where p is an artificial parameter and $a_i, i=0,1,\dots$ are unknown coefficients. We now define

$$u_0 = \xi_1 + \xi_2 x + L^{-1}[\sum_{n=0}^{+\infty} a_n x^n],$$

$$u_1 = L^{-1}[f(x)] - pL^{-1}[\sum_{n=0}^{+\infty} a_n x^n] - L^{-1}[R(u_0) + A_0(u_0)],$$

$$u_2 = -L^{-1}[R(u_1) + A_1(u_0, u_1)],$$

$$u_3 = -L^{-1}[R(u_2) + A_2(u_0, u_1, u_2)],$$

$\vdots$

To avoid calculation of $A_n, n=1,2,\dots$, let determine $a_i, i=0,1,\dots$ such that $u_1 = 0$. This implies that

$$u_2 = u_3 = \dots = 0.$$

Setting $p=1$, yields the solution of Eq. (1) with the initial conditions as follows

$$u(x) = \xi_1 + \xi_2 x + L^{-1}[\sum_{n=0}^{+\infty} a_n x^n].$$

3. Application

In this section, three singular initial value Emden-Fowler type equations, including a homogeneous nonlinear Emden-Fowler equation with $\alpha=1$, and two inhomogeneous Emden-Fowler equations, with $\alpha=2$ and $\alpha>2$ will be solved to illustrate the efficiency of the method. The computations associated with these examples have been performed by Maple package.

Example 3.1: Consider the homogeneous nonlinear Emden-Fowler equation with $\alpha=1$ [1]

$$u'' + \frac{1}{x}u' - u^3 + 3u^5 = 0,$$

with the following initial conditions

$$u(0)=1, \quad u'(0)=0.$$

Traditional ADM. As we know in the traditional ADM, we will reach the following expression

$$\sum_{n=0}^{+\infty} u_n = u(0) + u'(0)x$$

$$-L^{-1}[R(\sum_{n=0}^{+\infty} u_n) + \sum_{n=0}^{+\infty} A_n(u_0, u_1, \dots, u_n)], \quad (5)$$

where $L^{-1} = \int_0^x \int_0^x (\cdot) dx dx$, $R(u) = (1/x)u'$, and $A_n, n=0,1,\dots$ are as the following

$$A_0 = -u_0^3 + 3u_0^5,$$

$$A_1 = -3u_0^2 u_1(x) + 15u_0^4 u_1(x),$$

$\vdots$

We can define

$$u_0 = u(0) + u'(0)x,$$

$$u_n = -L^{-1}[R(u_{n-1}) + A_{n-1}(u_0, u_1, \dots, u_{n-1})], \quad n=1,2,\dots$$

Therefore

$$u_0(x) = 1, \quad u_1(x) = -x^2,$$

$$u_2(x) = x^2 + x^4, \quad u_3(x) = -x^2 - \frac{4}{3}x^4 - \frac{13}{10}x^6,$$

$\vdots$

Now, the series solution derived by the traditional ADM can be written as follows

$$u(x) = 1 - x^2 - \frac{1}{3}x^4 - \frac{13}{10}x^6 + \dots$$

Modified ADM. To solve the problem by the modified ADM, let us rewrite (5) as follows

$$\sum_{n=0}^{+\infty} u_n = u(0) + u'(0)x$$

$$+L^{-1}[\sum_{n=0}^{+\infty} a_n x^n] - pL^{-1}[\sum_{n=0}^{+\infty} a_n x^n]$$

$$-L^{-1}[R(\sum_{n=0}^{+\infty} u_n) + \sum_{n=0}^{+\infty} A_n(u_0, u_1, \dots, u_n)],$$

where p is an artificial parameter and $a_i, i=0,1,\dots$ are unknown coefficients. We now define

$$u_0 = u(0) + u'(0)x + L^{-1}[\sum_{n=0}^{+\infty} a_n x^n] = 1 + \frac{1}{2}a_0 x^2$$

$$+ \frac{1}{6}a_1 x^3 + \frac{1}{12}a_2 x^4 + \frac{1}{20}a_3 x^5 + \dots,$$

$$u_1 = -pL^{-1}[\sum_{n=0}^{+\infty} a_n x^n] - L^{-1}[R(u_0) + A_0(u_0)],$$

$$u_2 = -L^{-1}[R(u_1) + A_1(u_0, u_1)],$$

$\vdots$

To avoid calculation of $A_n, n=1,2,\dots$, let determine $a_i, i=0,1,\dots$ such that $u_1 = 0$. Thus

$$(-1 - \frac{1}{2}a_0 - \frac{1}{2}pa_0)x^2 + (-\frac{1}{12}a_1 - \frac{1}{6}pa_1)x^3$$

$$+ (-\frac{1}{2}a_0 - \frac{1}{36}a_2 - \frac{1}{12}pa_2)x^4 + \dots = 0.$$

It can be easily shown that

$$a_0 = -\frac{2}{1+p}, \quad a_1 = 0, \quad a_2 = \frac{36}{(1+p)(1+3p)}, \quad a_3 = 0, \quad \dots$$

Setting $p=1$, results in

$$u(x) = 1 - \frac{1}{2}x^2 + \frac{3}{8}x^4 - \frac{5}{16}x^6 + \dots = \frac{1}{\sqrt{1+x^2}},$$

which is the exact solution of the problem.

Example 3.2: Consider the inhomogeneous Emden-Fowler equation with $\alpha = 2$ [1]

$$u'' + \frac{2}{x}u' - (6 + 4x^2)u = 6 - 6x^2 - 4x^4,$$

subject to the following initial conditions

$$u(0) = 1, \quad u'(0) = 0.$$

Traditional ADM. Applying the traditional ADM yields

$$u_0(x) = 1 + 3x^2 - \frac{1}{2}x^4 - \frac{2}{15}x^6,$$

$$u_1(x) = -3x^2 + \frac{13}{6}x^4 + \frac{53}{150}x^6 - \frac{1}{20}x^8 - \frac{4}{675}x^{10},$$

$$u_2(x) = 6x^2 - \frac{53}{18}x^4 - \frac{27}{250}x^6 + \frac{869}{4200}x^8 + \frac{1663}{121500}x^{10}$$

$$- \frac{53}{29700}x^{12} - \frac{8}{61425}x^{14},$$

⋮

Consequently, the series solution obtained by the traditional ADM is as follows

$$u(x) = 1 + 6x^2 - \frac{23}{18}x^4 + \frac{14}{125}x^6 + \frac{659}{4200}x^8 + \frac{943}{121500}x^{10} - \frac{53}{29700}x^{12} - \frac{8}{61425}x^{14} + \dots$$

Modified ADM. A similar procedure, described in previous example, leads to

$$u_0 = u(0) + u'(0)x + L^{-1}[\sum_{n=0}^{+\infty} a_n x^n] = 1 + \frac{1}{2}a_0 x^2$$

$$+ \frac{1}{6}a_1 x^3 + \frac{1}{12}a_2 x^4 + \frac{1}{20}a_3 x^5 + \dots,$$

$$u_1 = L^{-1}[6 - 6x^2 - 4x^4] - pL^{-1}[\sum_{n=0}^{+\infty} a_n x^n] - L^{-1}[R(u_0)],$$

where $L^{-1} = \int_0^x \int_0^x (\cdot) dx dx$ and $R(u) = (2/x)u' - (6 + 4x^2)u$. Now, by setting $u_1(x) = 0$, we find

$$(6 - a_0 - \frac{1}{2}pa_0)x^2 + (-\frac{1}{6}a_1 - \frac{1}{6}pa_1)x^3 + (-\frac{1}{6} + \frac{1}{4}a_0 - \frac{1}{18}a_2 - \frac{1}{12}pa_2)x^4 + \dots = 0.$$

It can be shown that

$$a_0 = \frac{12}{2+p}, \quad a_1 = 0, \quad a_2 = -\frac{6(p-16)}{(2+p)(2+3p)}, \quad a_3 = 0, \quad \dots$$

Setting $p=1$, yields

$$u(x) = 1 + 2x^2 + \frac{1}{2}x^4 + \frac{1}{6}x^6 + \dots = x^2$$

$$+ (1 + x^2 + \frac{1}{2}x^4 + \frac{1}{6}x^6 + \dots),$$

which converges to the exact solution

$$u(x) = x^2 + e^{x^2}.$$

Example 3.3: Consider the inhomogeneous Emden-Fowler equation with $\alpha > 2$ [1]

$$u'' + \frac{4}{x}u' - (18x + 9x^4)u = 20 - 36x^3 - 18x^6,$$

with the following initial conditions

$$u(0) = 1, \quad u'(0) = 0.$$

Traditional ADM. Using the traditional ADM leads to

$$u_0(x) = 1 + 10x^2 - \frac{9}{5}x^5 - \frac{9}{28}x^8,$$

$$u_1(x) = -40x^2 + 3x^3 + \frac{54}{5}x^5 + \frac{3}{10}x^6$$

$$+ \frac{297}{245}x^8 - \frac{1539}{7700}x^{11} - \frac{81}{5096}x^{14},$$

$$u_2(x) = 160x^2 - 6x^3 - \frac{234}{5}x^5 + \frac{39}{25}x^6$$

$$- \frac{12519}{3430}x^8 + \frac{9}{20}x^9 + \frac{156573}{134750}x^{11} + \frac{9}{440}x^{12}$$

$$+ \frac{2873799}{63763700}x^{14} - \frac{365229}{47647600}x^{17} - \frac{729}{1936480}x^{20},$$

⋮

Therefore, the series solution resulted from the traditional ADM is as follows

$$u(x) = 1 + 130x^2 - 3x^3 - \frac{189}{5}x^5 + \frac{93}{50}x^6$$

$$- \frac{18927}{6860}x^8 + \frac{9}{20}x^9 + \frac{23571}{24500}x^{11} + \frac{9}{440}x^{12}$$

$$+ \frac{3720573}{127527400}x^{14} - \frac{365229}{47647600}x^{17} - \frac{729}{1936480}x^{20} + \dots$$

Modified ADM. In a manner similar to that described in previous examples, we can define

$$u_0 = u(0) + u'(0)x + L^{-1}[\sum_{n=0}^{+\infty} a_n x^n] = 1 + \frac{1}{2}a_0 x^2$$

$$+ \frac{1}{6}a_1 x^3 + \frac{1}{12}a_2 x^4 + \frac{1}{20}a_3 x^5 + \dots,$$

$$u_1 = L^{-1}[20 - 36x^3 - 18x^6] - pL^{-1}[\sum_{n=0}^{+\infty} a_n x^n] - L^{-1}[R(u_0)],$$

where $L^{-1} = \int_0^x \int_0^x (\cdot) dx dx$ and $R(u) = (4/x)u' - (18x + 9x^4)u$. Now, if we set $u_1(x)$ equal to zero, then we obtain

$$(10 - 2a_0 - \frac{1}{2}pa_0)x^2 + (3 - \frac{1}{3}a_1 - \frac{1}{6}pa_1)x^3 \\ + (-\frac{1}{9}a_2 - \frac{1}{12}pa_2)x^4 + \dots = 0.$$

It can be easily shown that

$$a_0 = \frac{20}{4+p}, \quad a_1 = \frac{18}{2+p}, \quad a_2 = 0, \quad a_3 = -\frac{36(p-1)}{(1+p)(4+p)}, \dots$$

Setting $p=1$, results in

$$u(x) = 1 + 2x^2 + x^3 + \frac{1}{2}x^6 + \dots = 2x^2 \\ + (1 + x^3 + \frac{1}{2}x^6 + \dots) = 2x^2 + e^{x^3},$$

which is the exact solution of the problem.

4. Conclusion

In this article, a new and effective modification of Adomian decomposition method was proposed to solve singular, Emden-Fowler type, initial value problems with $\alpha=1$, 2, and >2 . As it was observed

- The method only requires the calculation of the first Adomian polynomial.
- The method provides the solution of the problems in the form of a convergent series, whereas the traditional ADM fails.
- The method overcomes the singularity at $x=0$.

It is worth mentioning that the proposed method can be extended for solving systems of ordinary differential equations of Emden-Fowler type as follows [23]

$$u_1'' + \frac{\alpha_1}{x} u_1' + f_1(u_1, u_2) = g_1(x), \quad u_1(0) = a_1, \quad u_1'(0) = 0,$$

$$u_2'' + \frac{\alpha_2}{x} u_2' + f_2(u_1, u_2) = g_2(x), \quad u_2(0) = a_2, \quad u_2'(0) = 0.$$

References

- [1] A.M. Wazwaz, A reliable treatment of singular Emden-Fowler initial value problems and boundary value problems, *Applied Mathematics and Computation*, 217 (2011), 10387–10395. <http://dx.doi.org/10.1016/j.amc.2011.04.084>.
- [2] K. Parand, M. Dehghan, A.R. Rezaei, S.M. Ghaderi, An approximation algorithm for the solution of the nonlinear Lane-Emden type equations arising in astrophysics using Hermite functions collocation method, *Computer Physics Communications*, 181 (2010), 1096–1108. <http://dx.doi.org/10.1016/j.cpc.2010.02.018>.
- [3] K. Parand, M. Shahini, M. Dehghan, Rational Legendre pseudospectral approach for solving nonlinear differential equations of Lane-Emden type, *Journal of Computational Physics*, 228 (2009), 8830–8840. <http://dx.doi.org/10.1016/j.jcp.2009.08.029>.
- [4] A.M. Wazwaz, A new algorithm for solving differential equations of Lane-Emden type, *Applied Mathematics and Computation*, 118 (2001), 287–310. [http://dx.doi.org/10.1016/S0096-3003\(99\)00223-4](http://dx.doi.org/10.1016/S0096-3003(99)00223-4).
- [5] S.J. Liao, A new analytic algorithm of Lane-Emden type equations, *Applied Mathematics and Computation*, 142 (2003), 1–16. [http://dx.doi.org/10.1016/S0096-3003\(02\)00943-8](http://dx.doi.org/10.1016/S0096-3003(02)00943-8).
- [6] O.P. Singh, R.K. Pandey, V.K. Singh, An analytic algorithm of Lane-Emden type equations arising in astrophysics using modified homotopy analysis method, *Computer Physics Communications*, 180 (2009), 1116–1124. <http://dx.doi.org/10.1016/j.cpc.2009.01.012>.
- [7] J.I. Ramos, Linearization techniques for singular initial-value problems of ordinary differential equations, *Applied Mathematics and Computation*, 161 (2005), 525–542. <http://dx.doi.org/10.1016/j.amc.2003.12.047>.
- [8] A. Yildirim, T. Ozis, Solutions of singular IVPs of Lane-Emden type by homotopy perturbation method, *Physics Letters A*, 369 (2007), 70–76. <http://dx.doi.org/10.1016/j.physleta.2007.04.072>.
- [9] X. Shang, P. Wu, X. Shao, An efficient method for solving Emden-Fowler equations, *Journal of the Franklin Institute*, 346 (2009), 889–897. <http://dx.doi.org/10.1016/j.franklin.2009.07.005>.
- [10] J.H. He, Variational approach to the Lane-Emden equation, *Applied Mathematics and Computation*, 143 (2003), 539–541. [http://dx.doi.org/10.1016/S0096-3003\(02\)00382-X](http://dx.doi.org/10.1016/S0096-3003(02)00382-X).
- [11] J.I. Ramos, Series approach to the Lane-Emden equation and comparison with the homotopy perturbation method, *Chaos, Solitons & Fractals*, 38 (2008), 400–408. <http://dx.doi.org/10.1016/j.chaos.2006.11.018>.
- [12] S.A. Yousefi, Legendre wavelets method for solving differential equations of Lane-Emden type, *Applied Mathematics and Computation*, 181 (2006), 1417–1422. <http://dx.doi.org/10.1016/j.amc.2006.02.031>.
- [13] N.T. Shawagfeh, Nonperturbative approximate solution for Lane-Emden equation, *Journal of Mathematical Physics*, 34 (1993), 4364–4369. <http://dx.doi.org/10.1063/1.530005>.
- [14] A.S. Bataineh, M.S.M. Noorani, I. Hashim, Homotopy analysis method for singular IVPs of Emden-Fowler type, *Communications in Nonlinear Science and Numerical Simulation*, 14 (2009), 1121–1131. <http://dx.doi.org/10.1016/j.cnsns.2008.02.004>.
- [15] M.S.H. Chowdhury, I. Hashim, Solutions of Emden-Fowler equations by homotopy perturbation method, *Nonlinear Analysis: Real World Applications*, 10 (2009), 104–115. <http://dx.doi.org/10.1016/j.nonrwa.2007.08.017>.
- [16] H.R. Marzban, H.R. Tabrizdooz, M. Razzaghi, Hybrid functions for nonlinear initial-value problems with applications to Lane-Emden type equations, *Physics Letters A*, 372 (2008), 5883–5886. <http://dx.doi.org/10.1016/j.physleta.2008.07.055>.
- [17] M. Dehghan, F. Shakeri, Approximate solution of a differential equation arising in astrophysics using the variational iteration method, *New Astronomy*, 13 (2008), 53–59. <http://dx.doi.org/10.1016/j.newast.2007.06.012>.
- [18] R.K. Pandey, N. Kumar, Solution of Lane-Emden type equations using Bernstein matrix of differentiation, *New Astronomy*, 17 (2012), 303–308. <http://dx.doi.org/10.1016/j.newast.2011.09.005>.
- [19] B. Muatjetjeja, C.M. Khalique, Exact solutions of the generalized Lane-Emden equations of the first and second kind, *Pramana Journal of Physics*, 77 (2011), 545–554. <http://dx.doi.org/10.1007/s12043-011-0174-4>.
- [20] J.S. Duan, Convenient analytic recurrence algorithms for the Adomian polynomials, *Applied Mathematics and Computation*, 217 (2011), 6337–6348. <http://dx.doi.org/10.1016/j.amc.2011.01.007>.
- [21] H. Aminikhah, J. Biazar, A new HPM for ordinary differential equations, *Numerical Methods for Partial Differential Equations*, 26 (2009), 480–489. <http://dx.doi.org/10.1002/num.20413>.
- [22] K. Hosseini, J. Biazar, R. Ansari, P. Gholamin, A new algorithm for solving differential equations, *Mathematical Methods in the Applied Sciences*, 35 (2012), 993–999. <http://dx.doi.org/10.1002/mma.1601>.
- [23] B. Muatjetjeja, C.M. Khalique, Lagrangian approach to a generalized coupled Lane-Emden system: Symmetries and first integrals, *Communications in Nonlinear Science and Numerical Simulation*, 15 (2010), 1166–1171. <http://dx.doi.org/10.1016/j.cnsns.2009.06.002>.